

CHAPTER P. PRELIMINARIES

Section P.1 Real Numbers and the Real Line (page 10)

- $\frac{2}{9} = 0.22222222 \dots = 0.\overline{2}$
- $\frac{1}{11} = 0.09090909 \dots = 0.\overline{09}$
- If $x = 0.121212 \dots$, then $100x = 12.121212 \dots = 12 + x$. Thus $99x = 12$ and $x = 12/99 = 4/33$.
- If $x = 3.277777 \dots$, then $10x - 32 = 0.77777 \dots$ and $100x - 320 = 7 + (10x - 32)$, or $90x = 295$. Thus $x = 295/90 = 59/18$.
- $1/7 = 0.142857142857 \dots = 0.\overline{142857}$
 $2/7 = 0.285714285714 \dots = 0.\overline{285714}$
 $3/7 = 0.428571428571 \dots = 0.\overline{428571}$
 $4/7 = 0.571428571428 \dots = 0.\overline{571428}$
 note the same cyclic order of the repeating digits
 $5/7 = 0.714285714285 \dots = 0.\overline{714285}$
 $6/7 = 0.857142857142 \dots = 0.\overline{857142}$
- Two different decimal expansions can represent the same number. For instance, both $0.999999 \dots = 0.\overline{9}$ and $1.000000 \dots = 1.\overline{0}$ represent the number 1.
- $x \geq 0$ and $x \leq 5$ define the interval $[0, 5]$.
- $x < 2$ and $x \geq -3$ define the interval $[-3, 2)$.
- $x > -5$ or $x < -6$ defines the union $(-\infty, -6) \cup (-5, \infty)$.
- $x \leq -1$ defines the interval $(-\infty, -1]$.
- $x > -2$ defines the interval $(-2, \infty)$.
- $x < 4$ or $x \geq 2$ defines the interval $(-\infty, \infty)$, that is, the whole real line.
- If $-2x > 4$, then $x < -2$. Solution: $(-\infty, -2)$
- If $3x + 5 \leq 8$, then $3x \leq 8 - 5 - 3$ and $x \leq 1$. Solution: $(-\infty, 1]$
- If $5x - 3 \leq 7 - 3x$, then $8x \leq 10$ and $x \leq 5/4$. Solution: $(-\infty, 5/4]$
- If $\frac{6-x}{4} \geq \frac{3x-4}{2}$, then $6-x \geq 6x-8$. Thus $14 \geq 7x$ and $x \leq 2$. Solution: $(-\infty, 2]$
- If $3(2-x) < 2(3+x)$, then $0 < 5x$ and $x > 0$. Solution: $(0, \infty)$
- If $x^2 < 9$, then $|x| < 3$ and $-3 < x < 3$. Solution: $(-3, 3)$
- Given: $1/(2-x) < 3$.
 CASE I. If $x < 2$, then $1 < 3(2-x) = 6-3x$, so $3x < 5$ and $x < 5/3$. This case has solutions $x < 5/3$.
 CASE II. If $x > 2$, then $1 > 3(2-x) = 6-3x$, so $3x > 5$ and $x > 5/3$. This case has solutions $x > 2$.
 Solution: $(-\infty, 5/3) \cup (2, \infty)$.
- Given: $(x+1)/x \geq 2$.
 CASE I. If $x > 0$, then $x+1 \geq 2x$, so $x \leq 1$.
 CASE II. If $x < 0$, then $x+1 \leq 2x$, so $x \geq 1$. (not possible)
 Solution: $(0, 1]$.
- Given: $x^2 - 2x \leq 0$. Then $x(x-2) \leq 0$. This is only possible if $x \geq 0$ and $x \leq 2$. Solution: $[0, 2]$.
- Given $6x^2 - 5x \leq -1$, then $(2x-1)(3x-1) \leq 0$, so either $x \leq 1/2$ and $x \geq 1/3$, or $x \leq 1/3$ and $x \geq 1/2$. The latter combination is not possible. The solution set is $[1/3, 1/2]$.
- Given $x^3 > 4x$, we have $x(x^2 - 4) > 0$. This is possible if $x < 0$ and $x^2 < 4$, or if $x > 0$ and $x^2 > 4$. The possibilities are, therefore, $-2 < x < 0$ or $2 < x < \infty$. Solution: $(-2, 0) \cup (2, \infty)$.
- Given $x^2 - x \leq 2$, then $x^2 - x - 2 \leq 0$ so $(x-2)(x+1) \leq 0$. This is possible if $x \leq 2$ and $x \geq -1$ or if $x \geq 2$ and $x \leq -1$. The latter situation is not possible. The solution set is $[-1, 2]$.
- Given: $\frac{x}{2} \geq 1 + \frac{4}{x}$.
 CASE I. If $x > 0$, then $x^2 \geq 2x + 8$, so that $x^2 - 2x - 8 \geq 0$, or $(x-4)(x+2) \geq 0$. This is possible for $x > 0$ only if $x \geq 4$.
 CASE II. If $x < 0$, then we must have $(x-4)(x+2) \leq 0$, which is possible for $x < 0$ only if $x \geq -2$.
 Solution: $[-2, 0) \cup [4, \infty)$.
- Given: $\frac{3}{x-1} < \frac{2}{x+1}$.
 CASE I. If $x > 1$ then $(x-1)(x+1) > 0$, so that $3(x+1) < 2(x-1)$. Thus $x < -5$. There are no solutions in this case.
 CASE II. If $-1 < x < 1$, then $(x-1)(x+1) < 0$, so $3(x+1) > 2(x-1)$. Thus $x > -5$. In this case all numbers in $(-1, 1)$ are solutions.
 CASE III. If $x < -1$, then $(x-1)(x+1) > 0$, so that $3(x+1) < 2(x-1)$. Thus $x < -5$. All numbers $x < -5$ are solutions.
 Solutions: $(-\infty, -5) \cup (-1, 1)$.
- If $|x| = 3$ then $x = \pm 3$.
- If $|x-3| = 7$, then $x-3 = \pm 7$, so $x = -4$ or $x = 10$.
- If $|2t+5| = 4$, then $2t+5 = \pm 4$, so $t = -9/2$ or $t = -1/2$.
- If $|1-t| = 1$, then $1-t = \pm 1$, so $t = 0$ or $t = 2$.

31. If $|8 - 3s| = 9$, then $8 - 3s = \pm 9$, so $3s = -1$ or 17 , and $s = -1/3$ or $s = 17/3$.
32. If $\left|\frac{s}{2} - 1\right| = 1$, then $\frac{s}{2} - 1 = \pm 1$, so $s = 0$ or $s = 4$.
33. If $|x| < 2$, then x is in $(-2, 2)$.
34. If $|x| \leq 2$, then x is in $[-2, 2]$.
35. If $|s - 1| \leq 2$, then $1 - 2 \leq s \leq 1 + 2$, so s is in $[-1, 3]$.
36. If $|t + 2| < 1$, then $-2 - 1 < t < -2 + 1$, so t is in $(-3, -1)$.
37. If $|3x - 7| < 2$, then $7 - 2 < 3x < 7 + 2$, so x is in $(5/3, 3)$.
38. If $|2x + 5| < 1$, then $-5 - 1 < 2x < -5 + 1$, so x is in $(-3, -2)$.
39. If $\left|\frac{x}{2} - 1\right| \leq 1$, then $1 - 1 \leq \frac{x}{2} \leq 1 + 1$, so x is in $[0, 4]$.
40. If $\left|2 - \frac{x}{2}\right| < \frac{1}{2}$, then $x/2$ lies between $2 - (1/2)$ and $2 + (1/2)$. Thus x is in $(3, 5)$.
41. The inequality $|x + 1| > |x - 3|$ says that the distance from x to -1 is greater than the distance from x to 3 , so x must be to the right of the point half-way between -1 and 3 . Thus $x > 1$.
42. $|x - 3| < 2|x| \Leftrightarrow x^2 - 6x + 9 = (x - 3)^2 < 4x^2 \Leftrightarrow 3x^2 + 6x - 9 > 0 \Leftrightarrow 3(x + 3)(x - 1) > 0$. This inequality holds if $x < -3$ or $x > 1$.
43. $|a| = a$ if and only if $a \geq 0$. It is false if $a < 0$.
44. The equation $|x - 1| = 1 - x$ holds if $|x - 1| = -(x - 1)$, that is, if $x - 1 < 0$, or, equivalently, if $x < 1$.
45. The triangle inequality $|x + y| \leq |x| + |y|$ implies that
- $$|x| \geq |x + y| - |y|.$$
- Apply this inequality with $x = a - b$ and $y = b$ to get
- $$|a - b| \geq |a| - |b|.$$
- Similarly, $|a - b| = |b - a| \geq |b| - |a|$. Since $||a| - |b||$ is equal to either $|a| - |b|$ or $|b| - |a|$, depending on the sizes of a and b , we have
- $$|a - b| \geq \left| |a| - |b| \right|.$$
- Section P.2 Cartesian Coordinates in the Plane (page 16)**
1. From $A(0, 3)$ to $B(4, 0)$, $\Delta x = 4 - 0 = 4$ and $\Delta y = 0 - 3 = -3$. $|AB| = \sqrt{4^2 + (-3)^2} = 5$.
2. From $A(-1, 2)$ to $B(4, -10)$, $\Delta x = 4 - (-1) = 5$ and $\Delta y = -10 - 2 = -12$. $|AB| = \sqrt{5^2 + (-12)^2} = 13$.
3. From $A(3, 2)$ to $B(-1, -2)$, $\Delta x = -1 - 3 = -4$ and $\Delta y = -2 - 2 = -4$. $|AB| = \sqrt{(-4)^2 + (-4)^2} = 4\sqrt{2}$.
4. From $A(0.5, 3)$ to $B(2, 3)$, $\Delta x = 2 - 0.5 = 1.5$ and $\Delta y = 3 - 3 = 0$. $|AB| = 1.5$.
5. Starting point: $(-2, 3)$. Increments $\Delta x = 4$, $\Delta y = -7$. New position is $(-2 + 4, 3 + (-7))$, that is, $(2, -4)$.
6. Arrival point: $(-2, -2)$. Increments $\Delta x = -5$, $\Delta y = 1$. Starting point was $(-2 - (-5), -2 - 1)$, that is, $(3, -3)$.
7. $x^2 + y^2 = 1$ represents a circle of radius 1 centred at the origin.
8. $x^2 + y^2 = 2$ represents a circle of radius $\sqrt{2}$ centred at the origin.
9. $x^2 + y^2 \leq 1$ represents points inside and on the circle of radius 1 centred at the origin.
10. $x^2 + y^2 = 0$ represents the origin.
11. $y \geq x^2$ represents all points lying on or above the parabola $y = x^2$.
12. $y < x^2$ represents all points lying below the parabola $y = x^2$.
13. The vertical line through $(-2, 5/3)$ is $x = -2$; the horizontal line through that point is $y = 5/3$.
14. The vertical line through $(\sqrt{2}, -1.3)$ is $x = \sqrt{2}$; the horizontal line through that point is $y = -1.3$.
15. Line through $(-1, 1)$ with slope $m = 1$ is $y = 1 + 1(x + 1)$, or $y = x + 2$.
16. Line through $(-2, 2)$ with slope $m = 1/2$ is $y = 2 + (1/2)(x + 2)$, or $x - 2y = -6$.
17. Line through $(0, b)$ with slope $m = 2$ is $y = b + 2x$.
18. Line through $(a, 0)$ with slope $m = -2$ is $y = 0 - 2(x - a)$, or $y = 2a - 2x$.
19. At $x = 2$, the height of the line $2x + 3y = 6$ is $y = (6 - 4)/3 = 2/3$. Thus $(2, 1)$ lies above the line.
20. At $x = 3$, the height of the line $x - 4y = 7$ is $y = (3 - 7)/4 = -1$. Thus $(3, -1)$ lies on the line.
21. The line through $(0, 0)$ and $(2, 3)$ has slope $m = (3 - 0)/(2 - 0) = 3/2$ and equation $y = (3/2)x$ or $3x - 2y = 0$.
22. The line through $(-2, 1)$ and $(2, -2)$ has slope $m = (-2 - 1)/(2 + 2) = -3/4$ and equation $y = 1 - (3/4)(x + 2)$ or $3x + 4y = -2$.
23. The line through $(4, 1)$ and $(-2, 3)$ has slope $m = (3 - 1)/(-2 - 4) = -1/3$ and equation $y = 1 - \frac{1}{3}(x - 4)$ or $x + 3y = 7$.

24. The line through $(-2, 0)$ and $(0, 2)$ has slope $m = (2 - 0)/(0 + 2) = 1$ and equation $y = 2 + x$.
25. If $m = -2$ and $b = \sqrt{2}$, then the line has equation $y = -2x + \sqrt{2}$.
26. If $m = -1/2$ and $b = -3$, then the line has equation $y = -(1/2)x - 3$, or $x + 2y = -6$.
27. $3x + 4y = 12$ has x -intercept $a = 12/3 = 4$ and y -intercept $b = 12/4 = 3$. Its slope is $-b/a = -3/4$.




Fig. P.2.27

28. $x + 2y = -4$ has x -intercept $a = -4$ and y -intercept $b = -4/2 = -2$. Its slope is $-b/a = 2/(-4) = -1/2$.




Fig. P.2.28

29. $\sqrt{2}x - \sqrt{3}y = 2$ has x -intercept $a = 2/\sqrt{2} = \sqrt{2}$ and y -intercept $b = -2/\sqrt{3}$. Its slope is $-b/a = 2/\sqrt{6} = \sqrt{2/3}$.




Fig. P.2.29

30. $1.5x - 2y = -3$ has x -intercept $a = -3/1.5 = -2$ and y -intercept $b = -3/(-2) = 3/2$. Its slope is $-b/a = 3/4$.




Fig. P.2.30

31. line through $(2, 1)$ parallel to $y = x + 2$ is $y = x - 1$; line perpendicular to $y = x + 2$ is $y = -x + 3$.
32. line through $(-2, 2)$ parallel to $2x + y = 4$ is $2x + y = -2$; line perpendicular to $2x + y = 4$ is $x - 2y = -6$.
33. We have

$$\begin{aligned} 3x + 4y = -6 &\implies 6x + 8y = -12 \\ 2x - 3y = 13 &\implies 6x - 9y = 39. \end{aligned}$$

Subtracting these equations gives $17y = -51$, so $y = -3$ and $x = (13 - 9)/2 = 2$. The intersection point is $(2, -3)$.

34. We have

$$\begin{aligned} 2x + y = 8 &\implies 14x + 7y = 56 \\ 5x - 7y = 1 &\implies 5x - 7y = 1. \end{aligned}$$

Adding these equations gives $19x = 57$, so $x = 3$ and $y = 8 - 2x = 2$. The intersection point is $(3, 2)$.

35. If $a \neq 0$ and $b \neq 0$, then $(x/a) + (y/b) = 1$ represents a straight line that is neither horizontal nor vertical, and does not pass through the origin. Putting $y = 0$ we get $x/a = 1$, so the x -intercept of this line is $x = a$; putting $x = 0$ gives $y/b = 1$, so the y -intercept is $y = b$.
36. The line $(x/2) - (y/3) = 1$ has x -intercept $a = 2$, and y -intercept $b = -3$.




Fig. P.2.36

37. The line through $(2, 1)$ and $(3, -1)$ has slope $m = (-1 - 1)/(3 - 2) = -2$ and equation $y = 1 - 2(x - 2) = 5 - 2x$. Its y -intercept is 5.

38. The line through $(-2, 5)$ and $(k, 1)$ has x -intercept 3, so also passes through $(3, 0)$. Its slope m satisfies

$$\frac{1-0}{k-3} = m = \frac{0-5}{3+2} = -1.$$

Thus $k-3 = -1$, and so $k = 2$.

39. $C = Ax + B$. If $C = 5,000$ when $x = 10,000$ and $C = 6,000$ when $x = 15,000$, then

$$\begin{aligned} 10,000A + B &= 5,000 \\ 15,000A + B &= 6,000 \end{aligned}$$

Subtracting these equations gives $5,000A = 1,000$, so $A = 1/5$. From the first equation, $2,000 + B = 5,000$, so $B = 3,000$. The cost of printing 100,000 pamphlets is $\$100,000/5 + 3,000 = \$23,000$.

40. -40° and -40° is the same temperature on both the Fahrenheit and Celsius scales.




Fig. P.2.40

41. $A = (2, 1)$, $B = (6, 4)$, $C = (5, -3)$
- $$|AB| = \sqrt{(6-2)^2 + (4-1)^2} = \sqrt{25} = 5$$
- $$|AC| = \sqrt{(5-2)^2 + (-3-1)^2} = \sqrt{25} = 5$$
- $$|BC| = \sqrt{(6-5)^2 + (4+3)^2} = \sqrt{50} = 5\sqrt{2}.$$
- Since $|AB| = |AC|$, triangle ABC is isosceles.

42. $A = (0, 0)$, $B = (1, \sqrt{3})$, $C = (2, 0)$
- $$|AB| = \sqrt{(1-0)^2 + (\sqrt{3}-0)^2} = \sqrt{4} = 2$$
- $$|AC| = \sqrt{(2-0)^2 + (0-0)^2} = \sqrt{4} = 2$$
- $$|BC| = \sqrt{(2-1)^2 + (0-\sqrt{3})^2} = \sqrt{4} = 2.$$
- Since $|AB| = |AC| = |BC|$, triangle ABC is equilateral.

43. $A = (2, -1)$, $B = (1, 3)$, $C = (-3, 2)$
- $$|AB| = \sqrt{(1-2)^2 + (3+1)^2} = \sqrt{17}$$
- $$|AC| = \sqrt{(-3-2)^2 + (2+1)^2} = \sqrt{34} = \sqrt{2}\sqrt{17}$$
- $$|BC| = \sqrt{(-3-1)^2 + (2-3)^2} = \sqrt{17}.$$

Since $|AB| = |BC|$ and $|AC| = \sqrt{2}|AB|$, triangle ABC is an isosceles right-angled triangle with right angle at B . Thus $ABCD$ is a square if D is displaced from C by the same amount A is from B , that is, by increments $\Delta x = 2 - 1 = 1$ and $\Delta y = -1 - 3 = -4$. Thus $D = (-3 + 1, 2 + (-4)) = (-2, -2)$.

44. If $M = (x_m, y_m)$ is the midpoint of P_1P_2 , then the displacement of M from P_1 equals the displacement of P_2 from M :

$$x_m - x_1 = x_2 - x_m, \quad y_m - y_1 = y_2 - y_m.$$

Thus $x_m = (x_1 + x_2)/2$ and $y_m = (y_1 + y_2)/2$.

45. If $Q = (x_q, y_q)$ is the point on P_1P_2 that is two thirds of the way from P_1 to P_2 , then the displacement of Q from P_1 equals twice the displacement of P_2 from Q :

$$x_q - x_1 = 2(x_2 - x_q), \quad y_q - y_1 = 2(y_2 - y_q).$$

Thus $x_q = (x_1 + 2x_2)/3$ and $y_q = (y_1 + 2y_2)/3$.

46. Let the coordinates of P be $(x, 0)$ and those of Q be $(X, -2X)$. If the midpoint of PQ is $(2, 1)$, then

$$(x + X)/2 = 2, \quad (0 - 2X)/2 = 1.$$

The second equation implies that $X = -1$, and the second then implies that $x = 5$. Thus P is $(5, 0)$.

47. $\sqrt{(x-2)^2 + y^2} = 4$ says that the distance of (x, y) from $(2, 0)$ is 4, so the equation represents a circle of radius 4 centred at $(2, 0)$.

48. $\sqrt{(x-2)^2 + y^2} = \sqrt{x^2 + (y-2)^2}$ says that (x, y) is equidistant from $(2, 0)$ and $(0, 2)$. Thus (x, y) must lie on the line that is the right bisector of the line from $(2, 0)$ to $(0, 2)$. A simpler equation for this line is $x = y$.

49. The line $2x + ky = 3$ has slope $m = -2/k$. This line is perpendicular to $4x + y = 1$, which has slope -4 , provided $m = 1/4$, that is, provided $k = -8$. The line is parallel to $4x + y = 1$ if $m = -4$, that is, if $k = 1/2$.

50. For any value of k , the coordinates of the point of intersection of $x + 2y = 3$ and $2x - 3y = -1$ will also satisfy the equation

$$(x + 2y - 3) + k(2x - 3y + 1) = 0$$

because they cause both expressions in parentheses to be 0. The equation above is linear in x and y , and so represents a straight line for any choice of k . This line will pass through $(1, 2)$ provided $1 + 4 - 3 + k(2 - 6 + 1) = 0$, that is, if $k = 2/3$. Therefore, the line through the point of intersection of the two given lines and through the point $(1, 2)$ has equation

$$x + 2y - 3 + \frac{2}{3}(2x - 3y + 1) = 0,$$

or, on simplification, $x = 1$.

Section P.3 Graphs of Quadratic Equations (page 22)

1. $x^2 + y^2 = 16$
2. $x^2 + (y - 2)^2 = 4$, or $x^2 + y^2 - 4y = 0$
3. $(x + 2)^2 + y^2 = 9$, or $x^2 + y^2 + 4x = 5$
4. $(x - 3)^2 + (y + 4)^2 = 25$, or $x^2 + y^2 - 6x + 8y = 0$.
5. $x^2 + y^2 - 2x = 3$
 $x^2 - 2x + 1 + y^2 = 4$
 $(x - 1)^2 + y^2 = 4$
 centre: $(1, 0)$; radius 2.
6. $x^2 + y^2 + 4y = 0$
 $x^2 + y^2 + 4y + 4 = 4$
 $x^2 + (y + 2)^2 = 4$
 centre: $(0, -2)$; radius 2.
7. $x^2 + y^2 - 2x + 4y = 4$
 $x^2 - 2x + 1 + y^2 + 4y + 4 = 9$
 $(x - 1)^2 + (y + 2)^2 = 9$
 centre: $(1, -2)$; radius 3.
8. $x^2 + y^2 - 2x - y + 1 = 0$
 $x^2 - 2x + 1 + y^2 - y + \frac{1}{4} = \frac{1}{4}$
 $(x - 1)^2 + (y - \frac{1}{2})^2 = \frac{1}{4}$
 centre: $(1, 1/2)$; radius $1/2$.
9. $x^2 + y^2 > 1$ represents all points lying outside the circle of radius 1 centred at the origin.
10. $x^2 + y^2 < 4$ represents the open disk consisting of all points lying inside the circle of radius 2 centred at the origin.
11. $(x + 1)^2 + y^2 \leq 4$ represents the closed disk consisting of all points lying inside or on the circle of radius 2 centred at the point $(-1, 0)$.
12. $x^2 + (y - 2)^2 \leq 4$ represents the closed disk consisting of all points lying inside or on the circle of radius 2 centred at the point $(0, 2)$.

13. Together, $x^2 + y^2 > 1$ and $x^2 + y^2 < 4$ represent annulus (washer-shaped region) consisting of all points that are outside the circle of radius 1 centred at the origin and inside the circle of radius 2 centred at the origin.
14. Together, $x^2 + y^2 \leq 4$ and $(x + 2)^2 + y^2 \leq 4$ represent the region consisting of all points that are inside or on both the circle of radius 2 centred at the origin and the circle of radius 2 centred at $(-2, 0)$.
15. Together, $x^2 + y^2 < 2x$ and $x^2 + y^2 < 2y$ (or, equivalently, $(x - 1)^2 + y^2 < 1$ and $x^2 + (y - 1)^2 < 1$) represent the region consisting of all points that are inside both the circle of radius 1 centred at $(1, 0)$ and the circle of radius 1 centred at $(0, 1)$.
16. $x^2 + y^2 - 4x + 2y > 4$ can be rewritten $(x - 2)^2 + (y + 1)^2 > 9$. This equation, taken together with $x + y > 1$, represents all points that lie both outside the circle of radius 3 centred at $(2, -1)$ and above the line $x + y = 1$.
17. The interior of the circle with centre $(-1, 2)$ and radius $\sqrt{6}$ is given by $(x + 1)^2 + (y - 2)^2 < 6$, or $x^2 + y^2 + 2x - 4y < 1$.
18. The exterior of the circle with centre $(2, -3)$ and radius 4 is given by $(x - 2)^2 + (y + 3)^2 > 16$, or $x^2 + y^2 - 4x + 6y > 3$.
19. $x^2 + y^2 < 2, \quad x \geq 1$
20. $x^2 + y^2 > 4, \quad (x - 1)^2 + (y - 3)^2 < 10$
21. The parabola with focus $(0, 4)$ and directrix $y = -4$ has equation $x^2 = 16y$.
22. The parabola with focus $(0, -1/2)$ and directrix $y = 1/2$ has equation $x^2 = -2y$.
23. The parabola with focus $(2, 0)$ and directrix $x = -2$ has equation $y^2 = 8x$.
24. The parabola with focus $(-1, 0)$ and directrix $x = 1$ has equation $y^2 = -4x$.
25. $y = x^2/2$ has focus $(0, 1/2)$ and directrix $y = -1/2$.




Fig. P.3.25

26. $y = -x^2$ has focus $(0, -1/4)$ and directrix $y = 1/4$.




Fig. P.3.26

27. $x = -y^2/4$ has focus $(-1, 0)$ and directrix $x = 1$.




Fig. P.3.27

28. $x = y^2/16$ has focus $(4, 0)$ and directrix $x = -4$.




Fig. P.3.28

29.




Fig. P.3.29

- a) has equation $y = x^2 - 3$.
- b) has equation $y = (x - 4)^2$ or $y = x^2 - 8x + 16$.
- c) has equation $y = (x - 3)^2 + 3$ or $y = x^2 - 6x + 12$.
- d) has equation $y = (x - 4)^2 - 2$, or $y = x^2 - 8x + 14$.
30. a) If $y = mx$ is shifted to the right by amount x_1 , the equation $y = m(x - x_1)$ results. If (a, b) satisfies this equation, then $b = m(a - x_1)$, and so $x_1 = a - (b/m)$. Thus the shifted equation is $y = m(x - a + (b/m)) = m(x - a) + b$.
- b) If $y = mx$ is shifted vertically by amount y_1 , the equation $y = mx + y_1$ results. If (a, b) satisfies this equation, then $b = ma + y_1$, and so $y_1 = b - ma$. Thus the shifted equation is $y = mx + b - ma = m(x - a) + b$, the same equation obtained in part (a).
31. $y = \sqrt{(x/3) + 1}$
32. $4y = \sqrt{x + 1}$
33. $y = \sqrt{(3x/2) + 1}$
34. $(y/2) = \sqrt{4x + 1}$
35. $y = 1 - x^2$ shifted down 1, left 1 gives $y = -(x + 1)^2$.
36. $x^2 + y^2 = 5$ shifted up 2, left 4 gives $(x + 4)^2 + (y - 2)^2 = 5$.
37. $y = (x - 1)^2 - 1$ shifted down 1, right 1 gives $y = (x - 2)^2 - 2$.
38. $y = \sqrt{x}$ shifted down 2, left 4 gives $y = \sqrt{x + 4} - 2$.

39. $y = x^2 + 3$, $y = 3x + 1$. Subtracting these equations gives $x^2 - 3x + 2 = 0$, or $(x - 1)(x - 2) = 0$. Thus $x = 1$ or $x = 2$. The corresponding values of y are 4 and 7. The intersection points are (1, 4) and (2, 7).

40. $y = x^2 - 6$, $y = 4x - x^2$. Subtracting these equations gives $2x^2 - 4x - 6 = 0$, or $2(x - 3)(x + 1) = 0$. Thus $x = 3$ or $x = -1$. The corresponding values of y are 3 and -5 . The intersection points are (3, 3) and $(-1, -5)$.

41. $x^2 + y^2 = 25$, $3x + 4y = 0$. The second equation says that $y = -3x/4$. Substituting this into the first equation gives $25x^2/16 = 25$, so $x = \pm 4$. If $x = 4$, then the second equation gives $y = -3$; if $x = -4$, then $y = 3$. The intersection points are (4, -3) and $(-4, 3)$. Note that having found values for x , we substituted them into the linear equation rather than the quadratic equation to find the corresponding values of y . Had we substituted into the quadratic equation we would have got more solutions (four points in all), but two of them would have failed to satisfy $3x + 4y = 12$. When solving systems of nonlinear equations you should always verify that the solutions you find do satisfy the given equations.

42. $2x^2 + 2y^2 = 5$, $xy = 1$. The second equation says that $y = 1/x$. Substituting this into the first equation gives $2x^2 + (2/x^2) = 5$, or $2x^4 - 5x^2 + 2 = 0$. This equation factors to $(2x^2 - 1)(x^2 - 2) = 0$, so its solutions are $x = \pm 1/\sqrt{2}$ and $x = \pm\sqrt{2}$. The corresponding values of y are given by $y = 1/x$. Therefore, the intersection points are $(1/\sqrt{2}, \sqrt{2})$, $(-1/\sqrt{2}, -\sqrt{2})$, $(\sqrt{2}, 1/\sqrt{2})$, and $(-\sqrt{2}, -1/\sqrt{2})$.

43. $(x^2/4) + y^2 = 1$ is an ellipse with major axis between $(-2, 0)$ and $(2, 0)$ and minor axis between $(0, -1)$ and $(0, 1)$.




Fig. P.3.43

44. $9x^2 + 16y^2 = 144$ is an ellipse with major axis between $(-4, 0)$ and $(4, 0)$ and minor axis between $(0, -3)$ and $(0, 3)$.




Fig. P.3.44

45. $\frac{(x - 3)^2}{9} + \frac{(y + 2)^2}{4} = 1$ is an ellipse with centre at $(3, -2)$, major axis between $(0, -2)$ and $(6, -2)$ and minor axis between $(3, -4)$ and $(3, 0)$.




Fig. P.3.45

46. $(x - 1)^2 + \frac{(y + 1)^2}{4} = 4$ is an ellipse with centre at $(1, -1)$, major axis between $(1, -5)$ and $(1, 3)$ and minor axis between $(-1, -1)$ and $(3, -1)$.




Fig. P.3.46

47. $(x^2/4) - y^2 = 1$ is a hyperbola with centre at the origin and passing through $(\pm 2, 0)$. Its asymptotes are $y = \pm x/2$.




Fig. P.3.47

48. $x^2 - y^2 = -1$ is a rectangular hyperbola with centre at the origin and passing through $(0, \pm 1)$. Its asymptotes are $y = \pm x$.




Fig. P.3.48

49. $xy = -4$ is a rectangular hyperbola with centre at the origin and passing through $(2, -2)$ and $(-2, 2)$. Its asymptotes are the coordinate axes.




Fig. P.3.49

50. $(x - 1)(y + 2) = 1$ is a rectangular hyperbola with centre at $(1, -2)$ and passing through $(2, -1)$ and $(0, -3)$. Its asymptotes are $x = 1$ and $y = -2$.




Fig. P.3.50

51. a) Replacing x with $-x$ replaces a graph with its reflection across the y -axis.
 b) Replacing y with $-y$ replaces a graph with its reflection across the x -axis.
52. Replacing x with $-x$ and y with $-y$ reflects the graph in both axes. This is equivalent to rotating the graph 180° about the origin.
53. $|x| + |y| = 1$.
 In the first quadrant the equation is $x + y = 1$.
 In the second quadrant the equation is $-x + y = 1$.
 In the third quadrant the equation is $-x - y = 1$.
 In the fourth quadrant the equation is $x - y = 1$.




Fig. P.3.53

Section P.4 Functions and Their Graphs (page 31)

- $f(x) = 1 + x^2$; domain \mathbb{R} , range $[1, \infty)$
- $f(x) = 1 - \sqrt{x}$; domain $[0, \infty)$, range $(-\infty, 1]$
- $G(x) = \sqrt{8 - 2x}$; domain $(-\infty, 4]$, range $[0, \infty)$
- $F(x) = 1/(x - 1)$; domain $(-\infty, 1) \cup (1, \infty)$, range $(-\infty, 0) \cup (0, \infty)$

5. $h(t) = \frac{t}{\sqrt{2-t}}$; domain $(-\infty, 2)$, range \mathbb{R} . (The equation $y = h(t)$ can be squared and rewritten as $t^2 + y^2t - 2y^2 = 0$, a quadratic equation in t having real solutions for every real value of y . Thus the range of h contains all real numbers.)

6. $g(x) = \frac{1}{1 - \sqrt{x-2}}$; domain $(2, 3) \cup (3, \infty)$, range $(-\infty, 0) \cup (0, \infty)$. The equation $y = g(x)$ can be solved for $x = 2 - (1 - (1/y))^2$ so has a real solution provided $y \neq 0$.




Fig. P.4.7

Graph (ii) is the graph of a function because vertical lines can meet the graph only once. Graphs (i), (iii), and (iv) do not have this property, so are not graphs of functions.

8.




Fig. P.4.8

- a) is the graph of $x(1-x)^2$, which is positive for $x > 0$.
- b) is the graph of $x^2 - x^3 = x^2(1-x)$, which is positive if $x < 1$.
- c) is the graph of $x - x^4$, which is positive if $0 < x < 1$ and behaves like x near 0.
- d) is the graph of $x^3 - x^4$, which is positive if $0 < x < 1$ and behaves like x^3 near 0.

9.

x	$f(x) = x^4$
0	0
± 0.5	0.0625
± 1	1
± 1.5	5.0625
± 2	16




Fig. P.4.9

10.

x	$f(x) = x^{2/3}$
0	0
± 0.5	0.62996
± 1	1
± 1.5	1.3104
± 2	1.5874






Fig. P.4.10


11. $f(x) = x^2 + 1$ is even: $f(-x) = f(x)$
12. $f(x) = x^3 + x$ is odd: $f(-x) = -f(x)$
13. $f(x) = \frac{x}{x^2 - 1}$ is odd: $f(-x) = -f(x)$
14. $f(x) = \frac{1}{x^2 - 1}$ is even: $f(-x) = f(x)$
15. $f(x) = \frac{1}{x - 2}$ is odd about $(2, 0)$: $f(2 - x) = -f(2 + x)$
16. $f(x) = \frac{1}{x + 4}$ is odd about $(-4, 0)$:
 $f(-4 - x) = -f(-4 + x)$
17. $f(x) = x^2 - 6x$ is even about $x = 3$: $f(3 - x) = f(3 + x)$
18. $f(x) = x^3 - 2$ is odd about $(0, -2)$:
 $f(-x) + 2 = -(f(x) + 2)$
19. $f(x) = |x^3| = |x|^3$ is even: $f(-x) = f(x)$
20. $f(x) = |x + 1|$ is even about $x = -1$:
 $f(-1 - x) = f(-1 + x)$
21. $f(x) = \sqrt{2x}$ has no symmetry.
22. $f(x) = \sqrt{(x - 1)^2}$ is even about $x = 1$:
 $f(1 - x) = f(1 + x)$
- 23.




24.




25.




26.




27.




28.




29.




30.




31.




32.




33.




34.




35.




36.



37.



38.



39.




Fig. P.4.39(a) Fig. P.4.39(b)

40.






Fig. P.4.40(a) Fig. P.4.40(b)


41.




42.




43.




44.



45.



46.



47. Range is approximately $[-0.18, 0.68]$.




Fig. P.4.47

48. Range is approximately $(-\infty, 0.17]$.




Fig. P.4.48

49.




Fig. P.4.49

Apparent symmetry about $x = 1.5$.
This can be confirmed by calculating $f(3 - x)$, which turns out to be equal to $f(x)$.

50.




Fig. P.4.50

Apparent symmetry about $x = 1$.
This can be confirmed by calculating $f(2 - x)$, which turns out to be equal to $f(x)$.

51.




Fig. P.4.51

Apparent symmetry about $(2, 1)$, and about the lines $y = x - 1$ and $y = 3 - x$.

These can be confirmed by noting that $f(x) = 1 + \frac{1}{x-2}$, so the graph is that of $1/x$ shifted right 2 units and up one.

52.




Fig. P.4.52


Apparent symmetry about $(-2, 2)$.
This can be confirmed by calculating shifting the graph right by 2 (replace x with $x - 2$) and then down 2 (subtract 2). The result is $-5x/(1 + x^2)$, which is odd.

53. If f is both even and odd the $f(x) = f(-x) = -f(x)$, so $f(x) = 0$ identically.


Section P.5 Combining Functions to Make New Functions (page 37)

- $f(x) = x, g(x) = \sqrt{x-1}$.
 $\mathcal{D}(f) = \mathbb{R}, \mathcal{D}(g) = [1, \infty)$.
 $\mathcal{D}(f+g) = \mathcal{D}(f-g) = \mathcal{D}(fg) = \mathcal{D}(g/f) = [1, \infty)$,
 $\mathcal{D}(f/g) = (1, \infty)$.
 $(f+g)(x) = x + \sqrt{x-1}$
 $(f-g)(x) = x - \sqrt{x-1}$
 $(fg)(x) = x\sqrt{x-1}$
 $(f/g)(x) = x/\sqrt{x-1}$
 $(g/f)(x) = (\sqrt{1-x})/x$
- $f(x) = \sqrt{1-x}, g(x) = \sqrt{1+x}$.
 $\mathcal{D}(f) = (-\infty, 1], \mathcal{D}(g) = [-1, \infty)$.
 $\mathcal{D}(f+g) = \mathcal{D}(f-g) = \mathcal{D}(fg) = [-1, 1]$,
 $\mathcal{D}(f/g) = (-1, 1], \mathcal{D}(g/f) = [-1, 1)$.
 $(f+g)(x) = \sqrt{1-x} + \sqrt{1+x}$
 $(f-g)(x) = \sqrt{1-x} - \sqrt{1+x}$
 $(fg)(x) = \sqrt{1-x^2}$
 $(f/g)(x) = \sqrt{(1-x)/(1+x)}$
 $(g/f)(x) = \sqrt{(1+x)/(1-x)}$


3.




4.



5.



6.



7. $f(x) = x + 5, g(x) = x^2 - 3.$
 $f \circ g(0) = f(-3) = 2, \quad g(f(0)) = g(5) = 22$
 $f(g(x)) = f(x^2 - 3) = x^2 + 2$
 $g \circ f(x) = g(f(x)) = g(x + 5) = (x + 5)^2 - 3$
 $f \circ f(-5) = f(0) = 5, \quad g(g(2)) = g(1) = -2$
 $f(f(x)) = f(x + 5) = x + 10$
 $g \circ g(x) = g(g(x)) = (x^2 - 3)^2 - 3$

8. $f(x) = 2/x, g(x) = x/(1 - x).$
 $f \circ f(x) = 2/(2/x) = x; \quad \mathcal{D}(f \circ f) = \{x : x \neq 0\}$
 $f \circ g(x) = 2/(x/(1 - x)) = 2(1 - x)/x;$
 $\mathcal{D}(f \circ g) = \{x : x \neq 0, 1\}$
 $g \circ f(x) = (2/x)/(1 - (2/x)) = 2/(x - 2);$
 $\mathcal{D}(g \circ f) = \{x : x \neq 0, 2\}$
 $g \circ g(x) = (x/(1 - x))/(1 - (x/(1 - x))) = x/(1 - 2x);$
 $\mathcal{D}(g \circ g) = \{x : x \neq 1/2, 1\}$

9. $f(x) = 1/(1 - x), g(x) = \sqrt{x - 1}.$
 $f \circ f(x) = 1/(1 - (1/(1 - x))) = (x - 1)/x;$
 $\mathcal{D}(f \circ f) = \{x : x \neq 0, 1\}$
 $f \circ g(x) = 1/(1 - \sqrt{x - 1});$
 $\mathcal{D}(f \circ g) = \{x : x \geq 1, x \neq 2\}$
 $g \circ f(x) = \sqrt{1/(1 - x) - 1} = \sqrt{x/(1 - x)};$
 $\mathcal{D}(g \circ f) = [0, 1)$

$g \circ g(x) = \sqrt{\sqrt{x - 1} - 1}; \quad \mathcal{D}(g \circ g) = [2, \infty)$

10. $f(x) = (x + 1)/(x - 1) = 1 + 2/(x - 1), g(x) = \text{sgn}(x).$
 $f \circ f(x) = 1 + 2/(1 + (2/(x - 1) - 1)) = x;$
 $\mathcal{D}(f \circ f) = \{x : x \neq 1\}$
 $f \circ g(x) = \frac{\text{sgn } x + 1}{\text{sgn } x - 1} = 0; \quad \mathcal{D}(f \circ g) = (-\infty, 0)$
 $g \circ f(x) = \text{sgn} \left(\frac{x + 1}{x - 1} \right) = \begin{cases} 1 & \text{if } x < -1 \text{ or } x > 1; \\ -1 & \text{if } -1 < x < 1 \end{cases};$
 $\mathcal{D}(g \circ f) = \{x : x \neq -1, 1\}$
 $g \circ g(x) = \text{sgn}(\text{sgn}(x)) = \text{sgn}(x); \quad \mathcal{D}(g \circ g) = \{x : x \neq 0\}$

	$f(x)$	$g(x)$	$f \circ g(x)$
11.	x^2	$x + 1$	$(x + 1)^2$
12.	$x - 4$	$x + 4$	x
13.	\sqrt{x}	x^2	$ x $
14.	$2x^3 + 3$	$x^{1/3}$	$2x + 3$
15.	$(x + 1)/x$	$1/(x - 1)$	x
16.	$1/(x + 1)^2$	$x - 1$	$1/x^2$

17. $y = \sqrt{x}.$
 $y = 2 + \sqrt{x}:$ previous graph is raised 2 units.
 $y = 2 + \sqrt{3 + x}:$ previous graph is shifted left 3 units.
 $y = 1/(2 + \sqrt{3 + x}):$ previous graph turned upside down and shrunk vertically.




Fig. P.5.17

18.






Fig. P.5.18


19.




20.




21.




22.




23.




24.




25.



26.



27. $F(x) = Ax + B$
 (a) $F \circ F(x) = F(x)$
 $\Rightarrow A(Ax + B) + B = Ax + B$
 $\Rightarrow A[(A - 1)x + B] = 0$
 Thus, either $A = 0$ or $A = 1$ and $B = 0$.
 (b) $F \circ F(x) = x$
 $\Rightarrow A(Ax + B) + B = x$
 $\Rightarrow (A^2 - 1)x + (A + 1)B = 0$
 Thus, either $A = -1$ or $A = 1$ and $B = 0$.
28. $\lfloor x \rfloor = 0$ for $0 \leq x < 1$; $\lceil x \rceil = 0$ for $-1 \leq x < 0$.
29. $\lfloor x \rfloor = \lceil x \rceil$ for all integers x .
30. $\lceil -x \rceil = -\lfloor x \rfloor$ is true for all real x ; if $x = n + y$ where n is an integer and $0 \leq y < 1$, then $-x = -n - y$, so that $\lceil -x \rceil = -n$ and $\lfloor x \rfloor = n$.



31. $f(x)$ is called the integer part of x because $\lfloor f(x) \rfloor$ is the largest integer that does not exceed x ; i.e. $|x| = \lfloor f(x) \rfloor + y$, where $0 \leq y < 1$.




Fig. P.5.32

33. If f is even and g is odd, then: f^2 , g^2 , $f \circ g$, $g \circ f$, and $f \circ f$ are all even. fg , f/g , g/f , and $g \circ g$ are odd, and $f + g$ is neither even nor odd. Here are two typical verifications:

$$\begin{aligned} f \circ g(-x) &= f(g(-x)) = f(-g(x)) = f(g(x)) = f \circ g(x) \\ (fg)(-x) &= f(-x)g(-x) = f(x)[-g(x)] \\ &= -f(x)g(x) = -(fg)(x). \end{aligned}$$

The others are similar.

34. f even $\Leftrightarrow f(-x) = f(x)$
 f odd $\Leftrightarrow f(-x) = -f(x)$
 f even and odd $\Rightarrow f(x) = -f(x) \Rightarrow 2f(x) = 0$
 $\Rightarrow f(x) = 0$

35. a) Let $E(x) = \frac{1}{2}[f(x) + f(-x)]$.
 Then $E(-x) = \frac{1}{2}[f(-x) + f(x)] = E(x)$. Hence, $E(x)$ is even.
 Let $O(x) = \frac{1}{2}[f(x) - f(-x)]$.
 Then $O(-x) = \frac{1}{2}[f(-x) - f(x)] = -O(x)$ and $O(x)$ is odd.

$$\begin{aligned} E(x) + O(x) &= \frac{1}{2}[f(x) + f(-x)] + \frac{1}{2}[f(x) - f(-x)] \\ &= f(x). \end{aligned}$$

Hence, $f(x)$ is the sum of an even function and an odd function.

- b) If $f(x) = E_1(x) + O_1(x)$ where E_1 is even and O_1 is odd, then

$$E_1(x) + O_1(x) = f(x) = E(x) + O(x).$$

Thus $E_1(x) - E(x) = O(x) - O_1(x)$. The left side of this equation is an even function and the right side is an odd function. Hence both sides are both even and odd, and are therefore identically 0 by Exercise 36. Hence $E_1 = E$ and $O_1 = O$. This shows that f can be written in only one way as the sum of an even function and an odd function.

Section P.6 Polynomials and Rational Functions (page 43)

- $x^2 - 7x + 10 = (x + 5)(x + 2)$
The roots are -5 and -2 .
- $x^2 - 3x - 10 = (x - 5)(x + 2)$
The roots are 5 and -2 .
- If $x^2 + 2x + 2 = 0$, then $x = \frac{-2 \pm \sqrt{4 - 8}}{2} = -1 \pm i$.
The roots are $-1 + i$ and $-1 - i$.
 $x^2 + 2x + 2 = (x + 1 - i)(x + 1 + i)$.
- Rather than use the quadratic formula this time, let us complete the square.

$$\begin{aligned} x^2 - 6x + 13 &= x^2 - 6x + 9 + 4 \\ &= (x - 3)^2 + 2^2 \\ &= (x - 3 - 2i)(x - 3 + 2i). \end{aligned}$$

The roots are $3 + 2i$ and $3 - 2i$.

- $16x^4 - 8x^2 + 1 = (4x^2 - 1)^2 = (2x - 1)^2(2x + 1)^2$. There are two double roots: $1/2$ and $-1/2$.
- $x^4 + 6x^3 + 9x^2 = x^2(x^2 + 6x + 9) = x^2(x + 3)^2$. There are two double roots, 0 and -3 .

7. $x^3 + 1 = (x + 1)(x^2 - x + 1)$. One root is -1 . The other two are the solutions of $x^2 - x + 1 = 0$, namely

$$x = \frac{1 \pm \sqrt{1-4}}{2} = \frac{1}{2} \pm \frac{\sqrt{3}}{2}i.$$

We have

$$x^3 + 1 = (x + 1) \left(x - \frac{1}{2} - \frac{\sqrt{3}}{2}i \right) \left(x - \frac{1}{2} + \frac{\sqrt{3}}{2}i \right).$$

8. $x^4 - 1 = (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x - i)(x + i)$. The roots are $1, -1, i$, and $-i$.
9. $x^6 - 3x^4 + 3x^2 - 1 = (x^2 - 1)^3 = (x - 1)^3(x + 1)^3$. The roots are 1 and -1 , each with multiplicity 3 .

10. $x^5 - x^4 - 16x + 16 = (x - 1)(x^4 - 16)$
 $= (x - 1)(x^2 - 4)(x^2 + 4)$
 $= (x - 1)(x - 2)(x + 2)(x - 2i)(x + 2i)$.

The roots are $1, 2, -2, 2i$, and $-2i$.

11. $x^5 + x^3 + 8x^2 + 8 = (x^2 + 1)(x^3 + 8)$
 $= (x + 2)(x - i)(x + i)(x^2 - 2x + 4)$

Three of the five roots are $-2, i$ and $-i$. The remaining two are solutions of $x^2 - 2x + 4 = 0$, namely

$$x = \frac{2 \pm \sqrt{4-16}}{2} = 1 \pm \sqrt{3}i. \text{ We have}$$

$$x^5 + x^3 + 8x^2 + 8 = (x + 2)(x - i)(x + i)(x - 1 + \sqrt{3}i)(x - 1 - \sqrt{3}i).$$

12. $x^9 - 4x^7 - x^6 + 4x^4 = x^4(x^5 - x^2 - 4x^3 + 4)$
 $= x^4(x^3 - 1)(x^2 - 4)$
 $= x^4(x - 1)(x - 2)(x + 2)(x^2 + x + 1)$.

Seven of the nine roots are: 0 (with multiplicity 4), $1, 2$, and -2 . The other two roots are solutions of $x^2 + x + 1 = 0$, namely

$$x = \frac{-1 \pm \sqrt{1-4}}{2} = -\frac{1}{2} \pm \frac{\sqrt{3}}{2}i.$$

The required factorization of $x^9 - 4x^7 - x^6 + 4x^4$ is

$$x^4(x-1)(x-2)(x+2) \left(x - \frac{1}{2} + \frac{\sqrt{3}}{2}i \right) \left(x - \frac{1}{2} - \frac{\sqrt{3}}{2}i \right).$$

13. $\frac{x^3 - 1}{x^2 - 2} = \frac{x^3 - 2x + 2x - 1}{x^2 - 2}$
 $= \frac{x(x^2 - 2) + 2x - 1}{x^2 - 2}$
 $= x + \frac{2x - 1}{x^2 - 2}.$

14. $\frac{x^2}{x^2 + 5x + 3} = \frac{x^2 + 5x + 3 - 5x - 3}{x^2 + 5x + 3}$
 $= 1 + \frac{-5x - 3}{x^2 + 5x + 3}.$

15. $\frac{x^3}{x^2 + 2x + 3} = \frac{x^3 + 2x^2 + 3x - 2x^2 - 3x}{x^2 + 2x + 3}$
 $= \frac{x(x^2 + 2x + 3) - 2x^2 - 3x}{x^2 + 2x + 3}$
 $= x - \frac{2(x^2 + 2x + 3) - 4x - 6 - 3x}{x^2 + 2x + 3}$
 $= x - 2 + \frac{7x + 6}{x^2 + 2x + 3}.$

16. $\frac{x^4 + x^2}{x^3 + x^2 + 1} = \frac{x(x^3 + x^2 + 1) - x^3 - x + x^2}{x^3 + x^2 + 1}$
 $= x + \frac{-(x^3 + x^2 + 1) + x^2 + 1 - x + x^2}{x^3 + x^2 + 1}$
 $= x - 1 + \frac{2x^2 - x + 1}{x^3 + x^2 + 1}.$

17. Let $P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$, where $n \geq 1$. By the Factor Theorem, $x - 1$ is a factor of $P(x)$ if and only if $P(1) = 0$, that is, if and only if $a_n + a_{n-1} + \dots + a_1 + a_0 = 0$.

18. Let $P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$, where $n \geq 1$. By the Factor Theorem, $x + 1$ is a factor of $P(x)$ if and only if $P(-1) = 0$, that is, if and only if $a_0 - a_1 + a_2 - a_3 + \dots + (-1)^n a_n = 0$. This condition says that the sum of the coefficients of even powers is equal to the sum of coefficients of odd powers.

19. Let $P(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x + a_0$, where the coefficients $a_k, 0 \leq k \leq n$ are all real numbers, so that $\bar{a}_k = a_k$. Using the facts about conjugates of sums and products mentioned in the statement of the problem, we see that if $z = x + iy$, where x and y are real, then

$$\begin{aligned} \overline{P(z)} &= \overline{a_n z^n + a_{n-1} z^{n-1} + \dots + a_1 z + a_0} \\ &= a_n \bar{z}^n + a_{n-1} \bar{z}^{n-1} + \dots + a_1 \bar{z} + a_0 \\ &= P(\bar{z}). \end{aligned}$$

If z is a root of P , then $P(\bar{z}) = \overline{P(z)} = \bar{0} = 0$, and \bar{z} is also a root of P .

20. By the previous exercise, $\bar{z} = u - iv$ is also a root of P . Therefore $P(x)$ has two linear factors $x - u - iv$ and $x - u + iv$. The product of these factors is the real quadratic factor $(x - u)^2 - i^2 v^2 = x^2 - 2ux + u^2 + v^2$, which must also be a factor of $P(x)$.

21. By the previous exercise

$$\frac{P(x)}{x^2 - 2ux + u^2 + v^2} = \frac{P(x)}{(x - u - iv)(x - u + iv)} = Q_1(x),$$

where Q_1 , being a quotient of two polynomials with real coefficients, must also have real coefficients. If $z = u + iv$ is a root of P having multiplicity $m > 1$, then it must also be a root of Q_1 (of multiplicity $m - 1$), and so, therefore, \bar{z} must be a root of Q_1 , as must be the real quadratic $x^2 - 2ux + u^2 + v^2$. Thus

$$\frac{P(x)}{(x^2 - 2ux + u^2 + v^2)^2} = \frac{Q_1(x)}{x^2 - 2ux + u^2 + v^2} = Q_2(x),$$

where Q_2 is a polynomial with real coefficients. We can continue in this way until we get

$$\frac{P(x)}{(x^2 - 2ux + u^2 + v^2)^m} = Q_m(x),$$

where Q_m no longer has z (or \bar{z}) as a root. Thus z and \bar{z} must have the same multiplicity as roots of P .

Section P.7 The Trigonometric Functions (page 55)

$$1. \cos\left(\frac{3\pi}{4}\right) = \cos\left(\pi - \frac{\pi}{4}\right) = -\cos\frac{\pi}{4} = -\frac{1}{\sqrt{2}}$$

$$2. \tan\frac{-3\pi}{4} = -\tan\frac{3\pi}{4} = -1$$

$$3. \sin\frac{2\pi}{3} = \sin\left(\pi - \frac{\pi}{3}\right) = \sin\frac{\pi}{3} = \frac{\sqrt{3}}{2}$$

$$4. \sin\left(\frac{7\pi}{12}\right) = \sin\left(\frac{\pi}{4} + \frac{\pi}{3}\right) \\ = \sin\frac{\pi}{4}\cos\frac{\pi}{3} + \cos\frac{\pi}{4}\sin\frac{\pi}{3} \\ = \frac{1}{\sqrt{2}}\frac{1}{2} + \frac{1}{\sqrt{2}}\frac{\sqrt{3}}{2} = \frac{1 + \sqrt{3}}{2\sqrt{2}}$$

$$5. \cos\frac{5\pi}{12} = \cos\left(\frac{2\pi}{3} - \frac{\pi}{4}\right) \\ = \cos\frac{2\pi}{3}\cos\frac{\pi}{4} + \sin\frac{2\pi}{3}\sin\frac{\pi}{4} \\ = -\left(\frac{1}{2}\right)\left(\frac{1}{\sqrt{2}}\right) + \left(\frac{\sqrt{3}}{2}\right)\left(\frac{1}{\sqrt{2}}\right) \\ = \frac{\sqrt{3} - 1}{2\sqrt{2}}$$

$$6. \sin\frac{11\pi}{12} = \sin\frac{\pi}{12} \\ = \sin\left(\frac{\pi}{3} - \frac{\pi}{4}\right) \\ = \sin\frac{\pi}{3}\cos\frac{\pi}{4} - \cos\frac{\pi}{3}\sin\frac{\pi}{4} \\ = \left(\frac{\sqrt{3}}{2}\right)\left(\frac{1}{\sqrt{2}}\right) - \left(\frac{1}{2}\right)\left(\frac{1}{\sqrt{2}}\right) \\ = \frac{\sqrt{3} - 1}{2\sqrt{2}}$$

$$7. \cos(\pi + x) = \cos(2\pi - (\pi - x)) \\ = \cos(-(\pi - x)) \\ = \cos(\pi - x) = -\cos x$$

$$8. \sin(2\pi - x) = -\sin x$$

$$9. \sin\left(\frac{3\pi}{2} - x\right) = \sin\left(\pi - \left(x - \frac{\pi}{2}\right)\right) \\ = \sin\left(x - \frac{\pi}{2}\right) \\ = -\sin\left(\frac{\pi}{2} - x\right) \\ = -\cos x$$

$$10. \cos\left(\frac{3\pi}{2} + x\right) = \cos\frac{3\pi}{2}\cos x - \sin\frac{3\pi}{2}\sin x \\ = (-1)(-\sin x) = \sin x$$

$$11. \tan x + \cot x = \frac{\sin x}{\cos x} + \frac{\cos x}{\sin x} \\ = \frac{\sin^2 x + \cos^2 x}{\cos x \sin x} \\ = \frac{1}{\cos x \sin x}$$

$$12. \frac{\tan x - \cot x}{\tan x + \cot x} = \frac{\left(\frac{\sin x}{\cos x} - \frac{\cos x}{\sin x}\right)}{\left(\frac{\sin x}{\cos x} + \frac{\cos x}{\sin x}\right)} \\ = \frac{\left(\frac{\sin^2 x - \cos^2 x}{\cos x \sin x}\right)}{\left(\frac{\sin^2 x + \cos^2 x}{\cos x \sin x}\right)} \\ = \sin^2 x - \cos^2 x$$

$$13. \cos^4 x - \sin^4 x = (\cos^2 x - \sin^2 x)(\cos^2 x + \sin^2 x) \\ = \cos^2 x - \sin^2 x = \cos(2x)$$

$$14. (1 - \cos x)(1 + \cos x) = 1 - \cos^2 x = \sin^2 x \text{ implies} \\ \frac{1 - \cos x}{\sin x} = \frac{\sin x}{1 + \cos x}. \text{ Now} \\ \frac{1 - \cos x}{\sin x} = \frac{1 - \cos 2\left(\frac{x}{2}\right)}{\sin 2\left(\frac{x}{2}\right)} \\ = \frac{1 - \left(1 - 2\sin^2\left(\frac{x}{2}\right)\right)}{2\sin\frac{x}{2}\cos\frac{x}{2}} \\ = \frac{\sin\frac{x}{2}}{\cos\frac{x}{2}} = \tan\frac{x}{2}$$

$$15. \frac{1 - \cos x}{1 + \cos x} = \frac{2\sin^2\left(\frac{x}{2}\right)}{2\cos^2\left(\frac{x}{2}\right)} = \tan^2\left(\frac{x}{2}\right)$$

$$\begin{aligned}
 16. \quad \frac{\cos x - \sin x}{\cos x + \sin x} &= \frac{(\cos x - \sin x)^2}{(\cos x + \sin x)(\cos x - \sin x)} \\
 &= \frac{\cos^2 x - 2 \sin x \cos x + \sin^2 x}{\cos^2 x - \sin^2 x} \\
 &= \frac{1 - \sin(2x)}{\cos(2x)} \\
 &= \sec(2x) - \tan(2x)
 \end{aligned}$$




Fig. P.7.22

$$\begin{aligned}
 17. \quad \sin 3x &= \sin(2x + x) \\
 &= \sin 2x \cos x + \cos 2x \sin x \\
 &= 2 \sin x \cos^2 x + \sin x(1 - 2 \sin^2 x) \\
 &= 2 \sin x(1 - \sin^2 x) + \sin x - 2 \sin^3 x \\
 &= 3 \sin x - 4 \sin^3 x
 \end{aligned}$$

$$\begin{aligned}
 18. \quad \cos 3x &= \cos(2x + x) \\
 &= \cos 2x \cos x - \sin 2x \sin x \\
 &= (2 \cos^2 x - 1) \cos x - 2 \sin^2 x \cos x \\
 &= 2 \cos^3 x - \cos x - 2(1 - \cos^2 x) \cos x \\
 &= 4 \cos^3 x - 3 \cos x
 \end{aligned}$$

19. $\cos 2x$ has period π .




Fig. P.7.19

20. $\sin \frac{x}{2}$ has period 4π .




Fig. P.7.20

21. $\sin \pi x$ has period 2.






Fig. P.7.21

22. $\cos \frac{\pi x}{2}$ has period 4.

23.



24.



$$\begin{aligned}
 25. \quad \sin x &= \frac{3}{5}, \quad \frac{\pi}{2} < x < \pi \\
 \cos x &= -\frac{4}{5}, \quad \tan x = -\frac{3}{4}
 \end{aligned}$$




Fig. P.7.25

$$\begin{aligned}
 26. \quad \tan x &= 2 \text{ where } x \text{ is in } [0, \frac{\pi}{2}]. \text{ Then} \\
 \sec^2 x &= 1 + \tan^2 x = 1 + 4 = 5. \text{ Hence,} \\
 \sec x &= \sqrt{5} \text{ and } \cos x = \frac{1}{\sec x} = \frac{1}{\sqrt{5}}, \\
 \sin x &= \tan x \cos x = \frac{2}{\sqrt{5}}.
 \end{aligned}$$

$$27. \cos x = \frac{1}{3}, \quad -\frac{\pi}{2} < x < 0$$

$$\sin x = -\frac{\sqrt{8}}{3} = -\frac{2}{3}\sqrt{2}$$

$$\tan x = -\frac{\sqrt{8}}{1} = -2\sqrt{2}$$




Fig. P.7.27

$$28. \cos x = -\frac{5}{13} \text{ where } x \text{ is in } \left[\frac{\pi}{2}, \pi\right]. \text{ Hence,}$$

$$\sin x = \sqrt{1 - \cos^2 x} = \sqrt{1 - \frac{25}{169}} = \frac{12}{13},$$

$$\tan x = -\frac{12}{5}.$$

$$29. \sin x = -\frac{1}{2}, \quad \pi < x < \frac{3\pi}{2}$$

$$\cos x = -\frac{\sqrt{3}}{2}$$

$$\tan x = \frac{1}{\sqrt{3}}$$




Fig. P.7.29

$$30. \tan x = \frac{1}{2} \text{ where } x \text{ is in } \left[\pi, \frac{3\pi}{2}\right]. \text{ Then,}$$

$$\sec^2 x = 1 + \frac{1}{4} = \frac{5}{4}. \text{ Hence,}$$

$$\sec x = -\frac{\sqrt{5}}{2}, \quad \cos x = -\frac{2}{\sqrt{5}},$$

$$\sin x = \tan x \cos x = -\frac{1}{\sqrt{5}}.$$

$$31. c = 2, \quad B = \frac{\pi}{3}$$


$$a = c \cos B = 2 \times \frac{1}{2} = 1$$

$$b = c \sin B = 2 \times \frac{\sqrt{3}}{2} = \sqrt{3}$$

$$32. b = 2, \quad B = \frac{\pi}{3}$$

$$\frac{2}{a} = \tan B = \sqrt{3} \Rightarrow a = \frac{2}{\sqrt{3}}$$

$$\frac{2}{c} = \sin B = \frac{\sqrt{3}}{2} \Rightarrow c = \frac{4}{\sqrt{3}}$$



$$33. a = 5, \quad B = \frac{\pi}{6}$$

$$b = a \tan B = 5 \times \frac{1}{\sqrt{3}} = \frac{5}{\sqrt{3}}$$

$$c = \sqrt{a^2 + b^2} = \sqrt{25 + \frac{25}{3}} = \frac{10}{\sqrt{3}}$$

$$34. \sin A = \frac{a}{c} \Rightarrow a = c \sin A$$

$$35. \frac{a}{b} = \tan A \Rightarrow a = b \tan A$$

$$36. \cos B = \frac{a}{c} \Rightarrow a = c \cos B$$

$$37. \frac{b}{a} = \tan B \Rightarrow a = b \cot B$$

$$38. \sin A = \frac{a}{c} \Rightarrow c = \frac{a}{\sin A}$$

$$39. \frac{b}{c} = \cos A \Rightarrow c = b \sec A$$

$$40. \sin A = \frac{a}{c}$$

$$41. \sin A = \frac{a}{c} = \frac{\sqrt{c^2 - b^2}}{c}$$

$$42. \sin A = \frac{a}{c} = \frac{a}{\sqrt{a^2 + b^2}}$$

$$43. a = 4, b = 3, A = \frac{\pi}{4}$$


$$\sin B = b \frac{\sin A}{a} = \frac{3}{4} \frac{1}{\sqrt{2}} = \frac{3}{4\sqrt{2}}$$

$$44. \text{ Given that } a = 2, b = 2, c = 3.$$

$$\text{Since } a^2 = b^2 + c^2 - 2bc \cos A,$$

$$\cos A = \frac{a^2 - b^2 - c^2}{-2bc}$$

$$= \frac{4 - 4 - 9}{-2(2)(3)} = \frac{3}{4}.$$



$$45. a = 2, b = 3, c = 4$$

$$b^2 = a^2 + c^2 - 2ac \cos B$$

$$\text{Thus } \cos B = \frac{4 + 16 - 9}{2 \times 2 \times 4} = \frac{11}{16}$$

$$\sin B = \sqrt{1 - \frac{11^2}{16^2}} = \frac{\sqrt{256 - 121}}{16} = \frac{\sqrt{135}}{16}$$

46. Given that $a = 2$, $b = 3$, $C = \frac{\pi}{4}$.
 $c^2 = a^2 + b^2 - 2ab \cos C = 4 + 9 - 2(2)(3) \cos \frac{\pi}{4} = 13 - \frac{12}{\sqrt{2}}$.

Hence, $c = \sqrt{13 - \frac{12}{\sqrt{2}}} \approx 2.12479$.

47. $c = 3$, $A = \frac{\pi}{4}$, $B = \frac{\pi}{3}$ implies $C = \frac{5\pi}{12}$
 $\frac{a}{\sin A} = \frac{c}{\sin C} \Rightarrow a = \frac{1}{\sqrt{2}} \frac{3}{\sin\left(\frac{5\pi}{12}\right)}$

$$a = \frac{3}{\sqrt{2}} \frac{1}{\sin\left(\frac{7\pi}{12}\right)}$$

$$= \frac{3}{\sqrt{2}} \frac{2\sqrt{2}}{1 + \sqrt{3}} \text{ (by \#5)}$$

$$= \frac{6}{1 + \sqrt{3}}$$

48. Given that $a = 2$, $b = 3$, $C = 35^\circ$. Then
 $c^2 = 4 + 9 - 2(2)(3) \cos 35^\circ$, hence $c \approx 1.78050$.

49. $a = 4$, $B = 40^\circ$, $C = 70^\circ$
 Thus $A = 70^\circ$.
 $\frac{a}{\sin 40^\circ} = \frac{b}{\sin 70^\circ}$ so $b = 4 \frac{\sin 40^\circ}{\sin 70^\circ} = 2.736$

50. If $a = 1$, $b = \sqrt{2}$, $A = 30^\circ$, then $\frac{\sin B}{b} = \frac{\sin A}{a} = \frac{1}{2}$.
 Thus $\sin B = \frac{\sqrt{2}}{2} = \frac{1}{\sqrt{2}}$, $B = \frac{\pi}{4}$ or $\frac{3\pi}{4}$, and
 $C = \pi - \left(\frac{\pi}{4} + \frac{\pi}{6}\right) = \frac{7\pi}{12}$ or $C = \pi - \left(\frac{3\pi}{4} + \frac{\pi}{6}\right) = \frac{\pi}{12}$.
 Thus, $\cos C = \cos \frac{7\pi}{12} = \cos\left(\frac{\pi}{4} + \frac{\pi}{3}\right) = \frac{1 - \sqrt{3}}{2\sqrt{2}}$ or
 $\cos C = \cos \frac{\pi}{12} = \cos\left(\frac{\pi}{3} - \frac{\pi}{4}\right) = \frac{1 + \sqrt{3}}{2\sqrt{2}}$.

Hence,

$$c^2 = a^2 + b^2 - 2ab \cos C$$

$$= 1 + 2 - 2\sqrt{2} \cos C$$

$$= 3 - (1 - \sqrt{3}) \text{ or } 3 - (1 + \sqrt{3})$$

$$= 2 + \sqrt{3} \text{ or } 2 - \sqrt{3}.$$

Hence, $c = \sqrt{2 + \sqrt{3}}$ or $\sqrt{2 - \sqrt{3}}$.




Fig. P.7.50

51. Let h be the height of the pole and x be the distance from C to the base of the pole.
 Then $h = x \tan 50^\circ$ and $h = (x + 10) \tan 35^\circ$
 Thus $x \tan 50^\circ = x \tan 35^\circ + 10 \tan 35^\circ$ so

$$x = \frac{10 \tan 35^\circ}{\tan 50^\circ - \tan 35^\circ}$$

$$h = \frac{10 \tan 50^\circ \tan 35^\circ}{\tan 50^\circ - \tan 35^\circ} \approx 16.98$$

The pole is about 16.98 metres high.

52. See the following diagram. Since $\tan 40^\circ = h/a$, therefore $a = h/\tan 40^\circ$. Similarly, $b = h/\tan 70^\circ$.
 Since $a + b = 2$ km, therefore,

$$\frac{h}{\tan 40^\circ} + \frac{h}{\tan 70^\circ} = 2$$

$$h = \frac{2(\tan 40^\circ \tan 70^\circ)}{\tan 70^\circ + \tan 40^\circ} \approx 1.286 \text{ km.}$$




Fig. P.7.52

53. Area $\triangle ABC = \frac{1}{2}|BC|h = \frac{ah}{2} = \frac{ac \sin B}{2} = \frac{ab \sin C}{2}$
 By symmetry, area $\triangle ABC$ also $= \frac{1}{2}bc \sin A$




Fig. P.7.53

54. From Exercise 53, area = $\frac{1}{2}ac \sin B$. By Cosine Law,
 $\cos B = \frac{a^2 + c^2 - b^2}{2ac}$. Thus,

$$\begin{aligned}\sin B &= \sqrt{1 - \left(\frac{a^2 + c^2 - b^2}{2ac}\right)^2} \\ &= \frac{\sqrt{-a^4 - b^4 - c^4 + 2a^2b^2 + 2b^2c^2 + 2a^2c^2}}{2ac}.\end{aligned}$$

Hence, Area = $\frac{\sqrt{-a^4 - b^4 - c^4 + 2a^2b^2 + 2b^2c^2 + 2a^2c^2}}{4}$

square units. Since,

$$\begin{aligned}s(s-a)(s-b)(s-c) &= \frac{b+c+a}{2} \frac{b+c-a}{2} \frac{a-b+c}{2} \frac{a+b-c}{2} \\ &= \frac{1}{16} \left((b+c)^2 - a^2 \right) \left(a^2 - (b-c)^2 \right) \\ &= \frac{1}{16} \left(a^2 \left((b+c)^2 + (b-c)^2 \right) - a^4 - (b^2 - c^2)^2 \right) \\ &= \frac{1}{16} \left(2a^2b^2 + 2a^2c^2 - a^4 - b^4 - c^4 + 2b^2c^2 \right)\end{aligned}$$

Thus $\sqrt{s(s-a)(s-b)(s-c)}$ = Area of triangle.