

10728



Titu Andreescu

The American Mathematical Monthly, Vol. 106, No. 4. (Apr., 1999), p. 362.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9890%28199904%29106%3A4%3C362%3A1%3E2.0.CO%3B2-7>

The American Mathematical Monthly is currently published by Mathematical Association of America.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/maa.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

The JSTOR Archive is a trusted digital repository providing for long-term preservation and access to leading academic journals and scholarly literature from around the world. The Archive is supported by libraries, scholarly societies, publishers, and foundations. It is an initiative of JSTOR, a not-for-profit organization with a mission to help the scholarly community take advantage of advances in technology. For more information regarding JSTOR, please contact support@jstor.org.

PROBLEMS AND SOLUTIONS

Edited by **Gerald A. Edgar, Daniel H. Ullman, and Douglas B. West**

with the collaboration of Paul T. Bateman, Mario Benedicty, Paul Bracken, Duane M. Broline, Ezra A. Brown, Richard T. Bumby, Glenn G. Chappell, Randall Dougherty, Roger B. Eggleton, Ira M. Gessel, Bart Goddard, Jerrold R. Griggs, Douglas A. Hensley, Richard Holzsager, John R. Isbell, Robert Israel, Kiran S. Kedlaya, Murray S. Klamkin, Fred Kochman, Frederick W. Luttmann, Vania Mascioni, Frank B. Miles, Richard Pfeifer, Cecil C. Rousseau, Leonard Smiley, John Henry Steelman, Kenneth Stolarsky, Richard Stong, Charles Vanden Eynden, and William E. Watkins.

Proposed problems and solutions should be sent in duplicate to the MONTHLY problems address on the inside front cover. Submitted problems should include solutions and relevant references. Submitted solutions should arrive at that address before September 30, 1999; Additional information, such as generalizations and references, is welcome. The problem number and the solver's name and address should appear on each solution. An acknowledgement will be sent only if a mailing label is provided. An asterisk () after the number of a problem or a part of a problem indicates that no solution is currently available.*

PROBLEMS

10725. *Proposed by Vasile Mihai, Toronto, ON, Canada.* Fix a positive integer n . Given a permutation α of $\{1, 2, \dots, n\}$, let $f(\alpha) = \sum_{i=1}^n (\alpha(i) - \alpha(i+1))^2$, where $\alpha(n+1) = \alpha(1)$. Find the extreme values of $f(\alpha)$ as α ranges over all permutations of $\{1, 2, \dots, n\}$.

10726. *Proposed by Donald E. Knuth, Stanford University, Stanford, CA.* Start in state 0. For every nonnegative integer k , stay in state k for X_k units of time, then go to state $k+1$. What is the probability of being in state s after t units of time, assuming that X_k is distributed exponentially (a) with mean $1/(k+1)$? (b) with mean $1/2^k$?

10727. *Proposed by Jean Anglesio, Garches, France.* Let m be a fixed positive integer. For a positive integer n , let $s_m(n)$ be the sum of the m th powers of the decimal digits of n . For example, $s_3(172) = 1^3 + 7^3 + 2^3 = 352$. Starting with any positive integer n_0 , construct a sequence of positive integers by setting $n_k = s_m(n_{k-1})$ for every $k \geq 1$.

(a) Show that n_0, n_1, n_2, \dots is eventually periodic.

(b) Show that only finitely many periods are possible as n_0 varies.

10728. *Proposed by Titu Andreescu, American Mathematics Competitions, Lincoln, NE.* Determine all functions $f: \mathbb{Z} \rightarrow \mathbb{Z}$ satisfying

$$f(x^3 + y^3 + z^3) = (f(x))^3 + (f(y))^3 + (f(z))^3$$

for all integers x, y , and z .

10729. *Proposed by David P. Bellamy and Felix Lazebnik, University of Delaware, Newark, DE.* Let $I \subset \mathbb{R}$ be an open interval, and let n be a positive integer. Characterize the functions $f: I \rightarrow \mathbb{R}$ that have a continuous n th derivative and satisfy

$$f^{(n)} + p_1 f^{(n-1)} + \dots + p_{n-1} f' + p_n f = 0$$

for some continuous functions p_1, p_2, \dots, p_n on I .