

10756



Douglas Iannucci

The American Mathematical Monthly, Vol. 106, No. 8. (Oct., 1999), p. 778.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9890%28199910%29106%3A8%3C778%3A1%3E2.0.CO%3B2-C>

The American Mathematical Monthly is currently published by Mathematical Association of America.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/maa.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

The JSTOR Archive is a trusted digital repository providing for long-term preservation and access to leading academic journals and scholarly literature from around the world. The Archive is supported by libraries, scholarly societies, publishers, and foundations. It is an initiative of JSTOR, a not-for-profit organization with a mission to help the scholarly community take advantage of advances in technology. For more information regarding JSTOR, please contact support@jstor.org.

10756. Proposed by Douglas Iannucci, University of the Virgin Islands, St. Thomas, VI. Prove that

$$\cos \frac{\pi}{7} = \frac{1}{6} + \frac{\sqrt{7}}{6} \left(\cos \left(\frac{1}{3} \arccos \frac{1}{2\sqrt{7}} \right) + \sqrt{3} \sin \left(\frac{1}{3} \arccos \frac{1}{2\sqrt{7}} \right) \right).$$

10757. Proposed by Mark Kidwell, United States Naval Academy, Annapolis, MD. Given integers $a_0, a_1, a_2, \dots, a_n$ with $a_i \neq 0$ for $i \geq 1$, write $[a_0; a_1, a_2, \dots, a_n]$ for the continued fraction

$$a_0 + \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{\ddots + \frac{1}{a_n}}}}.$$

Every positive rational number has a unique representation as $[a_0; a_1, a_2, \dots, a_n]$ if we require that $a_0 \geq 0$, $a_i > 0$ for $1 \leq i \leq n-1$, and $a_n > 1$ (we call this the *standard representation*), but it can have other representations $[b_0; b_1, b_2, \dots, b_m]$ if we permit negative values for some of the b_i or if we permit $b_m = 1$. For example, $11/3 = [3; 1, 2] = [3; 1, 1, 1] = [4; -3]$. Prove or disprove: If r is a positive rational number, $r = [a_0; a_1, a_2, \dots, a_n]$ is the standard representation, and $r = [b_0; b_1, b_2, \dots, b_m]$ is another representation, then $a_0 + a_1 + \dots + a_n \leq |b_0| + |b_1| + \dots + |b_m|$, with strict inequality if any of the b_i are negative.

10758. Proposed by Mark Sapir, Vanderbilt University, Nashville, TN. Prove that the sum of the (decimal) digits of 9^n cannot equal 9 when $n > 2$.

10759. Proposed by Călin Popescu, Université Catholique de Louvain, Louvain-la-Neuve, Belgium. In triangle ABC , let h_a denote the altitude to the side BC and let r_a denote the exradius relative to side BC , i.e., the radius of the circle tangent to the extensions of sides AB and AC and to the side BC externally. Define h_b, h_c, r_b , and r_c correspondingly. Prove that $h_a^n r_a^n + h_b^n r_b^n + h_c^n r_c^n \leq r_a^n r_b^n + r_b^n r_c^n + r_c^n r_a^n$ for any integer n , and determine conditions for equality.

SOLUTIONS

Common Eigenvector of Commuting Matrices

10633 [1997, 975]. Proposed by Kiran S. Kedlaya, Princeton University, Princeton, NJ. Let S be a commuting family of n -by- n matrices over an arbitrary field. Suppose the matrices in S have a common eigenvector v , so that $Mv = \lambda_M v$ for all $M \in S$. Prove that the transposes of these matrices also have a common eigenvector with these eigenvalues, that is, a vector w satisfying $M^T w = \lambda_M w$ for all $M \in S$.

Solution by Alain Tissier, Montmermeil, France. Let K be the field. Set $\phi(M) = M - \lambda_M I$ and $\phi(S) = \{\phi(M) : M \in S\}$. Thus $\phi(S)$ is a commuting family of $n \times n$ matrices over K having a common nonzero vector v such that $\phi(M)v = 0$ for all $\phi(M) \in \phi(S)$. Since $\phi(M)^T = M^T - \lambda_M I$, we have to prove only that the transposes of the matrices in $\phi(S)$ have a common nonzero vector w satisfying $\phi(M)^T w = 0$ for $\phi(M) \in \phi(S)$. Thus we may suppose that $\lambda_M = 0$ for every M .

If all matrices in S are nilpotent, then the collection of transposes is also a commuting family of nilpotent matrices. In this case there is a nonzero vector w such that $M^T w = 0$ for all $M \in S$ (section 3.3 of J. E. Humphreys, *Introduction to Lie Algebras and Representation Theory*, Springer-Verlag, 1972). So we may assume that not all elements of S are nilpotent.

We proceed by induction on n . When $n = 1$ all the matrices are zero, so the conclusion is true. Take $n > 1$, and suppose the result is true for h -by- h matrices for each $h < n$. Let N