

10761



Fred Galvin

The American Mathematical Monthly, Vol. 106, No. 9. (Nov., 1999), p. 864.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9890%28199911%29106%3A9%3C864%3A1%3E2.0.CO%3B2-G>

The American Mathematical Monthly is currently published by Mathematical Association of America.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/maa.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

The JSTOR Archive is a trusted digital repository providing for long-term preservation and access to leading academic journals and scholarly literature from around the world. The Archive is supported by libraries, scholarly societies, publishers, and foundations. It is an initiative of JSTOR, a not-for-profit organization with a mission to help the scholarly community take advantage of advances in technology. For more information regarding JSTOR, please contact support@jstor.org.

PROBLEMS AND SOLUTIONS

Edited by **Gerald A. Edgar, Daniel H. Ullman, and Douglas B. West**

with the collaboration of Paul T. Bateman, Mario Benedicty, Paul Bracken, Duane M. Broline, Ezra A. Brown, Richard T. Bumby, Glenn G. Chappell, Randall Dougherty, Roger B. Eggleton, Ira M. Gessel, Bart Goddard, Jerrold R. Griggs, Douglas A. Hensley, John R. Isbell, Robert Israel, Kiran S. Kedlaya, Murray S. Klamkin, Fred Kochman, Frederick W. Luttmann, Vania Mascioni, Frank B. Miles, Richard Pfeifer, Cecil C. Rousseau, Leonard Smiley, John Henry Steelman, Kenneth Stolarsky, Richard Stong, Charles Vanden Eynden, and William E. Watkins.

Proposed problems and solutions should be sent in duplicate to the MONTHLY problems address on the inside front cover. Submitted problems should include solutions and relevant references. Submitted solutions should arrive at that address before April 30, 2000; Additional information, such as generalizations and references, is welcome. The problem number and the solver's name and address should appear on each solution. An acknowledgement will be sent only if a mailing label is provided. An asterisk () after the number of a problem or a part of a problem indicates that no solution is currently available.*

PROBLEMS

10760. *Proposed by Bruce Reznick, University of Illinois, Urbana, IL.* A function $f: \mathbb{N} \rightarrow \mathbb{C}$ is completely multiplicative if $f(1) = 1$ and $f(mn) = f(m)f(n)$ for all positive integers m and n . Find all completely multiplicative functions f with the property that the function $F(n) = \sum_{k=1}^n f(k)$ is also completely multiplicative.

10761. *Proposed by Fred Galvin, University of Kansas, Lawrence, KS.* Let G be a graph with n vertices. For each vertex v , let $f(v)$ be the maximum cardinality of an independent set of neighbors of v . Show that $\sum f(v) \leq n^2/2$, where the sum is taken over all vertices of G .

10762. *Proposed by Leroy Quet, Denver, CO.* Let $x_1 = 1$, and for $m \geq 1$ let $x_{m+1} = (m + 3/2)^{-1} \sum_{k=1}^m x_k x_{m+1-k}$. Evaluate $\lim_{m \rightarrow \infty} x_m/x_{m+1}$.

10763. *Proposed by Jean Anglesio, Garches, France.* Let ABC be a triangle; let O be its circumcenter, H its orthocenter, I its incenter, N its Nagel point, and X, Y, Z its excenters. Let S be defined so that O is the midpoint of HS , and let T denote the midpoint of SN . It is known that the orthocenter and the nine-point center of triangle XYZ are I and O , respectively. Prove that

(a) the circumcenter of triangle XYZ is T ; and

(b) the centroid of triangle XYZ is the centroid of SIN .

10764. *Proposed by Ray Redheffer, University of California, Los Angeles, CA.* Let $A = (a_{ij})$ be a real n -by- n matrix, and let x and y be real n -vectors satisfying $Ax = y$. Suppose that

$$\sum_{j \neq i} \max\{a_{ij}, 0\} < y_i \leq a_{ii} + \sum_{j \neq i} \min\{a_{ij}, 0\}$$

for all $i \in \{1, 2, \dots, n\}$. Show that $x_i > 0$ for all $i \in \{1, 2, \dots, n\}$.

10765. *Proposed by Peter J. Ferraro, Roselle Park, NJ.* Let f_n be the n th Fibonacci number, defined by $f_1 = f_2 = 1$ and $f_{n+2} = f_{n+1} + f_n$ for $n \geq 1$. Fix positive integers k and n with $n \geq 2k + 1$. Prove that $\lfloor \sqrt[k]{f_n} \rfloor - \lfloor \sqrt[k]{f_{n-k}} + \sqrt[k]{f_{n-2k}} \rfloor$ is 0 unless f_n is a k th power, when it is 1.