



## Logarithmic Convexity of Stirling's Ratio: 10680

Harold G. Diamond; Nathaniel Grossman

*The American Mathematical Monthly*, Vol. 106, No. 9. (Nov., 1999), p. 871.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9890%28199911%29106%3A9%3C871%3ALCOSR1%3E2.0.CO%3B2-R>

*The American Mathematical Monthly* is currently published by Mathematical Association of America.

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/maa.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.


---

The JSTOR Archive is a trusted digital repository providing for long-term preservation and access to leading academic journals and scholarly literature from around the world. The Archive is supported by libraries, scholarly societies, publishers, and foundations. It is an initiative of JSTOR, a not-for-profit organization with a mission to help the scholarly community take advantage of advances in technology. For more information regarding JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## Quadrilateral Center of Gravity

**10662** [1998, 464]. *Proposed by Joseph D. E. Konhauser and Stan Wagon, Macalester College, St. Paul, MN.* Find a construction for the center of gravity of the edges of a quadrilateral.

*Solution by the Con Amore Problems Group, Royal Danish School of Educational Studies, Copenhagen, Denmark.* If  $G$  is the center of gravity of the edges of the quadrilateral  $ABCD$  then  $G$  is also the center of gravity of particles with masses proportional to the lengths of the edges  $AB, BC, CD, DA$  placed at the midpoints  $P, Q, R, S$  of these edges. Construct these midpoints. The center of gravity for the particles at  $P$  and  $R$  is the point  $T$  on  $PR$  such that  $PT : TR = CD : AB$ , and the center of gravity for the particles at  $Q$  and  $S$  is the point  $U$  on  $QS$  with  $QU : US = DA : BC$ , so we construct the points  $T$  and  $U$ . If  $T$  and  $U$  coincide, we have  $G = T = U$ . If not, then  $G$  is the center of gravity of particles with masses proportional to the lengths of the line segments  $AB + CD$  and  $BC + DA$  placed at  $T$  and  $U$ , respectively. The sum of two line segments is constructed by placing them end to end. Thus  $G$  is the point on  $TU$  with  $TG : GU = (BC + DA) : (AB + CD)$ , and we construct this point.



Solved also by M. Benedicty, M. Boase (U. K.), G. D. Chakerian, R. J. Chapman (U. K.), S. S. Kim (Korea), J. H. Lindsey II, A. Nijenhuis, V. Pambuccian, C. R. Pranesachar (India), A. Sasane (The Netherlands), J. Schaer (Canada), Anchorage Math Solutions Group, GCHQ Problems Group, and the proposers.

## Logarithmic Convexity of Stirling's Ratio

**10680** [1998, 666]. *Proposed by Harold G. Diamond, University of Illinois, Urbana, IL.* For  $x > 0$  set  $g(x) = x \log \left( \Gamma(x+1) / (x^x e^{-x} \sqrt{2\pi x}) \right)$ . Show that  $g$  is concave down on  $(0, \infty)$ .

*Solution by Nathaniel Grossman, University of California, Los Angeles, CA.* It is enough to show that  $g''(x) < 0$  when  $x > 0$ . We begin with Binet's second expression for the gamma function, which we write in the form

$$\log \Gamma(x+1) = \left(x + \frac{1}{2}\right) \log x - x + \frac{1}{2} \log(2\pi) + 2k(x), \quad (*)$$

where  $k(x) = \int_0^\infty \arctan(t/x) (e^{2\pi t} - 1)^{-1} dt$  (M. Abramowitz and I. Stegun, *Handbook of Mathematical Functions*, Dover, New York, 1972, p. 258 (6.1.50)). From (\*) we find that  $g(x) = 2xk(x)$ , hence  $g''(x) = 2(xk''(x) + 2k'(x))$ . Easily justified differentiation under the integral sign leads to

$$xk''(x) + 2k'(x) = - \int_0^\infty \frac{2t^3}{(x^2 + t^2)^2 (e^{2\pi t} - 1)} dt,$$

in which the right hand side is clearly negative.

Solved also by J. Anglesio (France), P. Bracken (Canada), D. Bradley, E. Camouzis (Greece), R. J. Chapman (U. K.), R. A. Groeneveld, D. Krug, O. P. Lossers (The Netherlands), R. Martin (U. K.), A. McD. Mercer (Canada), P. Simeonov, A. Stadler (Switzerland), and NCCU Problems Group.