

10743



Calin Popescu

*The American Mathematical Monthly*, Vol. 106, No. 6. (Jun. - Jul., 1999), p. 586.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9890%28199906%2F07%29106%3A6%3C586%3A1%3E2.0.CO%3B2-9>

*The American Mathematical Monthly* is currently published by Mathematical Association of America.

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/maa.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

---

The JSTOR Archive is a trusted digital repository providing for long-term preservation and access to leading academic journals and scholarly literature from around the world. The Archive is supported by libraries, scholarly societies, publishers, and foundations. It is an initiative of JSTOR, a not-for-profit organization with a mission to help the scholarly community take advantage of advances in technology. For more information regarding JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# PROBLEMS AND SOLUTIONS

Edited by **Gerald A. Edgar**, **Daniel H. Ullman**, and **Douglas B. West**

with the collaboration of Paul T. Bateman, Mario Benedicty, Paul Bracken, Duane M. Broline, Ezra A. Brown, Richard T. Bumby, Glenn G. Chappell, Randall Dougherty, Roger B. Eggleton, Ira M. Gessel, Bart Goddard, Jerrold R. Griggs, Douglas A. Hensley, Richard Holzinger, John R. Isbell, Robert Israel, Kiran S. Kedlaya, Murray S. Klamkin, Fred Kochman, Frederick W. Luttmann, Vania Mascioni, Frank B. Miles, Richard Pfeifer, Cecil C. Rousseau, Leonard Smiley, John Henry Steelman, Kenneth Stolarsky, Richard Stong, Charles Vanden Eynden, and William E. Watkins.

*Proposed problems and solutions should be sent in duplicate to the MONTHLY problems address on the inside front cover. Submitted problems should include solutions and relevant references. Submitted solutions should arrive at that address before November 30, 1999; Additional information, such as generalizations and references, is welcome. The problem number and the solver's name and address should appear on each solution. An acknowledgement will be sent only if a mailing label is provided. An asterisk (\*) after the number of a problem or a part of a problem indicates that no solution is currently available.*

## PROBLEMS

**10739.** *Proposed by Oscar Ciaurri, Logroño, Spain.* Suppose that  $f: [0, 1] \rightarrow \mathbb{R}$  has a continuous second derivative with  $f''(x) > 0$  on  $(0, 1)$ , and suppose that  $f(0) = 0$ . Choose  $a \in (0, 1)$  such that  $f'(a) < f(1)$ . Show that there is a unique  $b \in (a, 1)$  such that  $f'(a) = f(b)/b$ .

**10740.** *Proposed by Charles Vanden Eynden, Illinois State University, Normal, IL.* A connected bipartite simple graph has a unique bipartition, meaning a partition of the vertices into two independent sets. Let  $\mathbf{G}$  be the set of such graphs that have no isomorphism that interchanges the two sets of the bipartition. Is there a criterion that for each  $G \in \mathbf{G}$  selects a well-defined set of the bipartition?

**10741.** *Proposed by Tim Keller, Fair Oaks, CA.* Is there an even base  $b$  for which there exist square integers of the form  $dddd_b$ ? By  $dddd_b$ , we mean the four-digit number in base  $b$  all of whose digits are  $d$ . For odd  $b$  we have the examples  $1111_7 = 20^2$  and  $4444_7 = 40^2$ .

**10742.** *Proposed by Emre Alkan, University of Wisconsin, Madison, WI.* Let us say that a finite group  $G$  has the *maximal property* if, for any prime  $p$  that divides  $|G|$ ,  $G$  has a maximal subgroup  $H$  such that  $p|H|$  divides  $|G|$ .

(a) Show that every finite solvable group has the maximal property.

(b) Show that there are infinitely many finite groups with the maximal property that are not solvable.

(c) Show that there are infinitely many finite groups without the maximal property that are not solvable.

**10743.** *Proposed by Călin Popescu, Université Catholique de Louvain, Louvain-La-Neuve, Belgium.* Let  $p \geq 5$  be prime, and let  $n$  be an integer such that  $(p+1)/2 \leq n \leq p-2$ . Let  $R = \sum (-1)^i \binom{n}{i}$ , where the sum is taken over the quadratic residues  $i$  modulo  $p$ , and let  $N = \sum (-1)^j \binom{n}{j}$ , where the sum is taken over the quadratic nonresidues  $j$  modulo  $p$ . Prove that exactly one of  $R$  and  $N$  is divisible by  $p$ .