

10744



Peter Lindqvist

The American Mathematical Monthly, Vol. 106, No. 6. (Jun. - Jul., 1999), p. 587.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9890%28199906%2F07%29106%3A6%3C587%3A1%3E2.0.CO%3B2-6>

The American Mathematical Monthly is currently published by Mathematical Association of America.

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/maa.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

The JSTOR Archive is a trusted digital repository providing for long-term preservation and access to leading academic journals and scholarly literature from around the world. The Archive is supported by libraries, scholarly societies, publishers, and foundations. It is an initiative of JSTOR, a not-for-profit organization with a mission to help the scholarly community take advantage of advances in technology. For more information regarding JSTOR, please contact support@jstor.org.

10744. Proposed by Peter Lindqvist, Norwegian University of Science and Technology, Trondheim, Norway, and Jaak Peetre, University of Lund, Lund, Sweden. Fix $p > 0$, and define functions $S(x)$, $C(x)$, and $T(x)$ for sufficiently small x by

$$x = \int_0^{S(x)} \frac{dt}{(1-t^p)^{(p-1)/p}}, \quad x = \int_{C(x)}^1 \frac{dt}{(1-t^p)^{(p-1)/p}}, \quad x = \int_0^{T(x)} \frac{dt}{(1+t^p)^{2/p}}.$$

Show that $S(x)^p + C(x)^p = 1$ and that $T(x) = S(x)/C(x)$. The case $p = 2$ yields the familiar trigonometric formulas.

10745. Proposed by M. J. Pelling, London, England. For $n \geq 1$, let $f(n)$ be the number of solutions (r, s, t) in positive integers to the Diophantine equation $rst = n(r + s + t)$.

(a) Prove that $f(n) = O(n^{1/2+\delta})$ for every $\delta > 0$.

(b)* Prove that $f(n) = O(n^\delta)$ for every $\delta > 0$.

SOLUTIONS

Using the Walls to Find the Center

10386 [1994, 474]. Proposed by Jordan Tabov, Bulgarian Academy of Sciences, Sofia, Bulgaria. Let a tetrahedron with vertices A_1, A_2, A_3, A_4 have altitudes that meet in a point H . For any point P , let P_1, P_2, P_3 , and P_4 be the feet of the perpendiculars from P to the faces $A_2A_3A_4, A_3A_4A_1, A_4A_1A_2$, and $A_1A_2A_3$, respectively. Prove that there exist constants a_1, a_2, a_3 , and a_4 such that one has

$$a_1 \overrightarrow{PP_1} + a_2 \overrightarrow{PP_2} + a_3 \overrightarrow{PP_3} + a_4 \overrightarrow{PP_4} = \overrightarrow{PH}$$

for every point P .

Solution by Murray S. Klamkin, University of Alberta, Edmonton, Alberta, Canada. More generally, let H and P be any two points in the space of the given tetrahedron and let P_1, P_2, P_3, P_4 be the feet of the lines through P parallel to HA_1, HA_2, HA_3, HA_4 in the faces of the tetrahedron opposite A_1, A_2, A_3, A_4 , respectively. Then there exist constants a_1, a_2, a_3, a_4 , independent of P , such that

$$a_1 \overrightarrow{PP_1} + a_2 \overrightarrow{PP_2} + a_3 \overrightarrow{PP_3} + a_4 \overrightarrow{PP_4} = \overrightarrow{PH}.$$

Let \mathbf{V} denote the vector from an origin outside the space of the given tetrahedron to any point V in the space of the tetrahedron. Then H and P have the representations (barycentric coordinates)

$$\mathbf{H} = x_1 \mathbf{A}_1 + x_2 \mathbf{A}_2 + x_3 \mathbf{A}_3 + x_4 \mathbf{A}_4 \quad (x_1 + x_2 + x_3 + x_4 = 1),$$

$$\mathbf{P} = u_1 \mathbf{A}_1 + u_2 \mathbf{A}_2 + u_3 \mathbf{A}_3 + u_4 \mathbf{A}_4 \quad (u_1 + u_2 + u_3 + u_4 = 1).$$

Since P_1 has the representation $\mathbf{P}_1 = r_2 \mathbf{A}_2 + r_3 \mathbf{A}_3 + r_4 \mathbf{A}_4$, where $r_2 + r_3 + r_4 = 1$, we must have

$$r_2 \mathbf{A}_2 + r_3 \mathbf{A}_3 + r_4 \mathbf{A}_4 - \mathbf{P} = \lambda_1 (\mathbf{H} - \mathbf{A}_1).$$

Since $\mathbf{A}_1, \mathbf{A}_2, \mathbf{A}_3, \mathbf{A}_4$ are independent vectors, we get $\lambda_1 = u_1/(1-x_1)$, so that $\overrightarrow{PP_1} = (\mathbf{P}_1 - \mathbf{P}) = (\mathbf{H} - \mathbf{A}_1)u_1/(1-x_1)$. Similarly,

$$(\mathbf{P}_i - \mathbf{P}) = (\mathbf{H} - \mathbf{A}_i) \frac{u_i}{1-x_i} \quad \text{for } i = 1, 2, 3, 4.$$

Choosing $a_i = 1 - x_i$, we obtain

$$\sum a_i (\mathbf{P}_i - \mathbf{P}) = \sum u_i (\mathbf{H} - \mathbf{A}_i) = \mathbf{H} - \mathbf{P} = \overrightarrow{PH}.$$

This proof generalizes to give an analogous result for n -dimensional simplices.

Solved also by J. Anglesio (France), R. J. Chapman (U. K.), M. Golomb, K. Hanes, N. Komanda, O. P. Lossers (The Netherlands), and the proposer.