

10743



Calin Popescu

*The American Mathematical Monthly*, Vol. 106, No. 7. (Aug. - Sep., 1999), p. 685.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9890%28199908%2F09%29106%3A7%3C685%3A1%3E2.0.CO%3B2-Z>

*The American Mathematical Monthly* is currently published by Mathematical Association of America.

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/maa.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

---

The JSTOR Archive is a trusted digital repository providing for long-term preservation and access to leading academic journals and scholarly literature from around the world. The Archive is supported by libraries, scholarly societies, publishers, and foundations. It is an initiative of JSTOR, a not-for-profit organization with a mission to help the scholarly community take advantage of advances in technology. For more information regarding JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# PROBLEMS AND SOLUTIONS

Edited by **Gerald A. Edgar, Daniel H. Ullman, and Douglas B. West**

with the collaboration of Paul T. Bateman, Mario Benedicty, Paul Bracken, Duane M. Broline, Ezra A. Brown, Richard T. Bumby, Glenn G. Chappell, Randall Dougherty, Roger B. Eggleton, Ira M. Gessel, Bart Goddard, Jerrold R. Griggs, Douglas A. Hensley, John R. Isbell, Robert Israel, Kiran S. Kedlaya, Murray S. Klamkin, Fred Kochman, Frederick W. Luttman, Vania Mascioni, Frank B. Miles, Richard Pfeifer, Cecil C. Rousseau, Leonard Smiley, John Henry Steelman, Kenneth Stolarsky, Richard Stong, Charles Vanden Eynden, and William E. Watkins.

*Proposed problems and solutions should be sent in duplicate to the MONTHLY problems address on the inside front cover. Submitted problems should include solutions and relevant references. Submitted solutions should arrive at that address before February 29, 2000; Additional information, such as generalizations and references, is welcome. The problem number and the solver's name and address should appear on each solution. An acknowledgement will be sent only if a mailing label is provided. An asterisk (\*) after the number of a problem or a part of a problem indicates that no solution is currently available.*

Problem **10743** [1999; 586] in the June–July 1999 issue was misstated. Here is the corrected version.

**10743.** *Proposed by Călin Popescu, Université Catholique de Louvain, Louvain-La-Neuve, Belgium.* Let  $R = \sum (-1)^i \binom{n}{i}$ , where the sum is taken over all  $i \in \{0, 1, \dots, n-1\}$  such that  $i+1$  is a quadratic residue modulo  $p$ , and let  $N = \sum (-1)^j \binom{n}{j}$ , where the sum is taken over all  $j \in \{0, 1, \dots, n-1\}$  such that  $j+1$  is a quadratic nonresidue modulo  $p$ . Prove that exactly one of  $R$  and  $N$  is divisible by  $p$ .

## PROBLEMS

**10746.** *Proposed by Stepan Tersian, University of Rousse, Rousse, Bulgaria.* Prove that

$$\int_0^{\infty} \left( e^{-y\sqrt{(s/x)^2+1}} - e^{-x\sqrt{(s/y)^2+1}} \right) \cos s \, ds = 0,$$

for all positive real numbers  $x$  and  $y$ .

**10747.** *Proposed by Athanasios Kalakos, Athens, Greece.* Find all differentiable functions  $f: \mathbb{R} \rightarrow \mathbb{R}$  that are twice differentiable on an open interval containing 0, have exactly one real root, satisfy  $f(1) = 1$ , and satisfy  $f'(f(t)) = 2f(t)$  for every  $t \in \mathbb{R}$ .

**10748.** *Proposed by Itshak Borosh, Douglas A. Hensley, and Joel Zinn, Texas A&M University, College Station, TX.* Let  $p$  and  $q$  be prime numbers, and let  $r$  be a positive integer such that  $q|(p-1)$ ,  $q \nmid r$ , and  $p > r^{q-1}$ . Show that for any integers  $a_1, a_2, \dots, a_r$ , if  $\sum_{j=1}^r a_j^{(p-1)/q} \equiv 0 \pmod{p}$ , then  $\prod_{j=1}^r a_j \equiv 0 \pmod{p}$ .

**10749.** *Proposed by Alain Grigis, Université Paris 13, Villetaneuse, France.* Let  $ABC$  be a triangle with a right angle at  $B$  and an angle of  $\pi/6$  at  $A$ . Consider a billiard path in the triangle that begins at  $A$ , reflects successively off side  $BC$  at  $P$ , off side  $AC$  at  $Q$ , off side  $AB$  at  $R$ , off side  $AC$  at  $S$ , and then ends at  $B$ .

(a) Show that  $AP$ ,  $QR$ , and  $SB$  are concurrent at a point  $X$ .

(b) Show that the angles formed at  $X$  measure  $\pi/3$ .

(c) Show that  $AX = XP + PQ + QX = XR + RS + SX = 2XB$ .