

10754



Paul Bracken

*The American Mathematical Monthly*, Vol. 106, No. 8. (Oct., 1999), p. 777.

Stable URL:

<http://links.jstor.org/sici?sici=0002-9890%28199910%29106%3A8%3C777%3A1%3E2.0.CO%3B2-F>

*The American Mathematical Monthly* is currently published by Mathematical Association of America.

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/about/terms.html>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/maa.html>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

---

The JSTOR Archive is a trusted digital repository providing for long-term preservation and access to leading academic journals and scholarly literature from around the world. The Archive is supported by libraries, scholarly societies, publishers, and foundations. It is an initiative of JSTOR, a not-for-profit organization with a mission to help the scholarly community take advantage of advances in technology. For more information regarding JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# PROBLEMS AND SOLUTIONS


Edited by **Gerald A. Edgar, Daniel H. Ullman, and Douglas B. West**

with the collaboration of Paul T. Bateman, Mario Benedicty, Paul Bracken, Duane M. Broline, Ezra A. Brown, Richard T. Bumby, Glenn G. Chappell, Randall Dougherty, Roger B. Eggleton, Ira M. Gessel, Bart Goddard, Jerrold R. Griggs, Douglas A. Hensley, John R. Isbell, Robert Israel, Kiran S. Kedlaya, Murray S. Klamkin, Fred Kochman, Frederick W. Luttmann, Vania Mascioni, Frank B. Miles, Richard Pfeifer, Cecil C. Rousseau, Leonard Smiley, John Henry Steelman, Kenneth Stolarsky, Richard Stong, Charles Vanden Eynden, and William E. Watkins.

*Proposed problems and solutions should be sent in duplicate to the MONTHLY problems address on the inside front cover. Submitted problems should include solutions and relevant references. Submitted solutions should arrive at that address before March 31, 2000; Additional information, such as generalizations and references, is welcome. The problem number and the solver's name and address should appear on each solution. An acknowledgement will be sent only if a mailing label is provided. An asterisk (\*) after the number of a problem or a part of a problem indicates that no solution is currently available.*

## PROBLEMS

**10753.** *Proposed by Louis Shapiro, Howard University, Washington, DC.* An ordered tree is a rooted tree in which the children of each node form a sequence as opposed to a set. The 5 ordered trees with 3 edges are



The number of ordered trees with  $n$  edges is the  $n$ th Catalan number  $\binom{2n}{n}/(n+1)$ . Therefore, if one draws each of the ordered trees with  $n$  edges, one draws a total of  $\binom{2n}{n}$  nodes. Prove that exactly half of these nodes are end-nodes (i.e., leaves with no children).

**10754.** *Proposed by Paul Bracken, Université de Montréal, Montréal, PQ, Canada.* Let  $\zeta(s) = \sum_{k=1}^{\infty} k^{-s}$ , and let  $\rho(s, n) = \sum_{k=n+1}^{\infty} k^{-s}$ . Show that for positive integers  $s \geq 2$ ,

$$\sum_{k=1}^{\infty} \frac{\rho(s, k)}{k} = \frac{s}{2} \zeta(s+1) - \frac{1}{2} \sum_{k=1}^{s-2} \zeta(s-k) \zeta(k+1).$$

**10755.** *Proposed by Jiro Fukuta, Motosu-gun, Gifu-ken, Japan.* An arbitrary circle  $O$  is drawn through vertices  $B$  and  $D$  of a convex quadrilateral  $ABCD$ . Let  $O_1$  be the circle tangent to lines  $AB$  and  $AD$  and tangent to  $O$  internally at a point of  $O$  on the opposite side of line  $BD$  from  $A$ . Let  $O_2$  be the circle tangent to lines  $CB$  and  $CD$  and tangent to  $O$  internally at a point of  $O$  on the opposite side of line  $BD$  from  $C$ . Let  $R_1$  and  $R_2$  be the radii of circles  $O_1$  and  $O_2$ , respectively, and let  $r_1$  and  $r_2$  be the radii of the incircles of triangles  $ABD$  and  $CBD$ , respectively. Prove that the quadrilateral  $ABCD$  is inscribable in a circle if and only if  $r_1/R_1 + r_2/R_2 = 1$ .