

June 21, 2004

CURVED FLATS, EXTERIOR DIFFERENTIAL SYSTEMS, AND CONSERVATION LAWS

CHUU-LIAN TERNG* AND ERXIAO WANG†

ABSTRACT. Let σ be an involution of a real semi-simple Lie group U , U_0 the subgroup fixed by σ , and U/U_0 the corresponding symmetric space. Ferus and Pedit called a submanifold M of a rank r symmetric space U/U_0 a *curved flat* if $T_p M$ is tangent to an r -dimensional flat of U/U_0 at p for each $p \in M$. They noted that the equation for curved flats is an integrable system. Bryant used the involution σ to construct an involutive exterior differential system \mathcal{I}_σ such that integral submanifolds of \mathcal{I}_σ are curved flats. Terng used r first flows in the U/U_0 -hierarchy of commuting soliton equations to construct the U/U_0 -system. She showed that the U/U_0 -system and the curved flat system are gauge equivalent, used the inverse scattering theory to solve the Cauchy problem globally with smooth rapidly decaying initial data, used loop group factorization to construct infinitely many families of explicit solutions, and noted that many these systems occur as the Gauss-Codazzi equations for submanifolds in space forms. The main goals of this paper are: (i) give a review of these known results, (ii) use techniques from soliton theory to construct infinitely many integral submanifolds and conservation laws for the exterior differential system \mathcal{I}_σ .

1. INTRODUCTION

Let G be a complex semi-simple Lie group, τ an involution of G such that its differential at the identity e is complex conjugate linear, and σ an involution of G such that the differential is complex linear. Assume that

$$\tau\sigma = \sigma\tau. \tag{1.1}$$

Let U be the fixed point set of τ , i.e., a real form of G . We will still use τ, σ to denote $d\tau_e$ and $d\sigma_e$ respectively. Let \mathcal{G}, \mathcal{U} denote the Lie algebras of G and U respectively. Since σ and τ commute, $\sigma(U) \subset U$. So $\sigma|_U$ is an involution of U . Let $\mathcal{U}_0, \mathcal{U}_1$ denote the $+1, -1$ eigenspaces of σ on \mathcal{U} . Then

$$[\mathcal{U}_0, \mathcal{U}_0] \subset \mathcal{U}_0, \quad [\mathcal{U}_0, \mathcal{U}_1] \subset \mathcal{U}_1, \quad [\mathcal{U}_1, \mathcal{U}_1] \subset \mathcal{U}_0.$$

The quotient space U/U_0 is a symmetric space, and the eigen-decomposition $\mathcal{U} = \mathcal{U}_0 + \mathcal{U}_1$ is called a *Cartan decomposition*.

Ferus and Pedit ([8]) called a submanifold M of a rank r symmetric space U/U_0 a *curved flat* if $T_p M$ is tangent to an r -dimensional flat of U/U_0 at p for each $p \in M$. They noted that the equation for curved flats is an integrable system. Bryant ([6]) used the involution σ to construct a natural

*Research supported in part by NSF Grant DMS-0306446.

†Research supported in part by Postdoctoral fellowship of MSRI.

involutive exterior differential system \mathcal{I}_σ such that integral submanifolds of \mathcal{I}_σ in U project down to curved flats in U/U_0 . Terng ([12]) used r first flows in the U/U_0 -hierarchy of commuting soliton equations to construct the U/U_0 -system. She showed that the U/U_0 -system and the curved flat system are gauge equivalent, used the inverse scattering theory to solve the Cauchy problem globally with smooth rapidly decaying initial data ([12]), used loop group factorization to construct infinitely many families of explicit solutions ([14]), and noted that many these systems occur as the Gauss-Codazzi equations for submanifolds in space forms ([12, 15]). The main goals of this paper are: (i) review some of these known results, (ii) use techniques from soliton theory to construct infinitely many integral submanifolds and conservation laws for the exterior differential system \mathcal{I}_σ .

We review the definitions of these systems next. An element $a \in \mathcal{U}_1$ is called *regular* if

- (i) $\mathcal{A} := \{y \in \mathcal{U}_1 \mid [a, y] = 0\}$ is a maximal abelian subspace in \mathcal{U}_1 ,
- (ii) $\text{Ad}(U_0)(\mathcal{A})$ is open in \mathcal{U}_1 .

Let (\cdot, \cdot) be an ad-invariant, non-degenerate bilinear form on \mathcal{U} . Given a linear subspace V of \mathcal{U} let

$$V^\perp = \{y \in \mathcal{U} \mid (y, V) = 0\}.$$

Assume that U/U_0 has rank r . Let \mathcal{A} be a maximal abelian subspace in \mathcal{U}_1 , and let $a_1, \dots, a_r \in \mathcal{A}$ be regular and form a basis of \mathcal{A} . The U/U_0 -system (cf. [12]) is the following PDE for $v : \mathbb{R}^r \rightarrow \mathcal{U}_1 \cap \mathcal{A}^\perp$:

$$[a_i, v_{x_j}] - [a_j, v_{x_i}] = [[a_i, v], [a_j, v]], \quad 1 \leq i \neq j \leq r. \quad (1.2)$$

These systems occur naturally in submanifold geometry. For example, the Gauss-Codazzi equations for isometric immersions of space forms in space forms ([9, 12, 2]), for isothermic surfaces in \mathbb{R}^n ([7, 2]), and for flat Lagrangian submanifolds in \mathbb{C}^n or in $\mathbb{C}P^n$ ([15]).

The U/U_0 -system also arises naturally from soliton theory (cf. [12]). In fact, given $1 \leq i \leq r$, $b \in \mathcal{A}$, and a positive integer j , the (b, j) -th soliton flow in the U/U_0 -hierarchy is a certain partial differential equation for $v : \mathbb{R}^2 \rightarrow \mathcal{U}_1 \cap \mathcal{A}^\perp$:

$$v_t = P_{b,j}(v).$$

For example, the second flow in the $SU(2)$ -hierarchy is the NLS (non-linear Schrödinger equation), the third flow in the $SU(2)/SO(2)$ -hierarchy is the modified KdV equation, and the first flow in the $SU(3)/SO(3)$ -hierarchy is the 3-wave equation. The U/U_0 -system (1.2) is given by the collection of the $(a_j, 1)$ -flows with $1 \leq j \leq r$ in the U/U_0 -hierarchy.

The *curved flat system associated to U/U_0* (cf. [9]) is the following first order system for $(A_1, \dots, A_r) : \mathbb{R}^r \rightarrow \prod_{i=1}^r \mathcal{U}_1$:

$$\begin{cases} (A_i)_{x_j} = (A_j)_{x_i}, & i \neq j, \\ [A_i, A_j] = 0, & i \neq j. \end{cases} \quad (1.3)$$

It is known that solutions of the curved flat system give rise to curved flats in U/U_0 ([8]).

Let $\alpha = g^{-1}dg$ be the Maurer-Cartan form on U . Write $\alpha = \alpha_0 + \alpha_1$ with respect to the Cartan decomposition $\mathcal{U} = \mathcal{U}_0 + \mathcal{U}_1$. Let \mathcal{I}_σ be the exterior differential ideal generated by α_0 . It was observed by Bryant ([6]) that (U, \mathcal{I}_σ) is involutive and the PDE for the exterior differential system \mathcal{I}_σ is the curved flat system (1.3). If $f : \mathcal{O} \rightarrow U$ is a maximal integral submanifold of the exterior differential system (EDS) (U, \mathcal{I}_σ) , then $f^*(\alpha_0) = 0$. The \mathcal{U}_0 component of the Maurer-Cartan equation $d\alpha + \frac{1}{2}[\alpha, \alpha] = 0$ gives

$$d\alpha_0 + \frac{1}{2}([\alpha_0, \alpha_0] + [\alpha_1, \alpha_1]) = 0.$$

So $f^*([\alpha_1, \alpha_1]) = 0$. This implies that $f^{-1}df$ is \mathcal{U}_1 -valued and the subspace $f^{-1}\text{Im}(df_p)$ is abelian for all $p \in \mathcal{O}$. This means that $(f^{-1}f_{y_1}, \dots, f^{-1}f_{y_r})$ is a solution of the curved flat system (1.3) with respect to any coordinate system y . Use the Cartan-Kähler Theorem we can see that the curved flat system should only depend on n functions of one variable, where $n = \dim(\mathcal{U}_1) - r$. But the curved flat system (1.3) is a system of $r(r-1)/2$ equations of nr functions. This indicates that the curved flat system has many redundant functions and we probably can use geometry to find a special coordinate system on integral submanifolds so that their PDE involves only n functions. This is indeed the case. We can find a special coordinate system x on an integral submanifold of (U, \mathcal{I}_σ) so that the corresponding PDE written in x coordinate is gauge equivalent to the U/U_0 -system.

Since the curved flat system is gauge equivalent to the U/U_0 -system, we can use techniques from soliton theory to construct infinitely many explicit integral submanifolds and conservations laws for the exterior differential system \mathcal{I}_σ .

This paper is organized as follows: We explain the gauge equivalence of the U/U_0 -system and the curved flat system in section 2, give a brief review of theory of exterior differential systems in section 3, and give Bryant's proof that the exterior system \mathcal{I}_σ on the Lie group U is involutive in section 4. Finally in section 5, we explain how to use the Birkhoff loop group factorization to construct infinitely many families of explicit solutions and commuting flows for the U/U_0 -system and conservation laws for \mathcal{I}_σ .

2. THE U/U_0 -SYSTEM

Let G, τ, σ, U, U_0 be as in section 1, U/U_0 the corresponding symmetric space, $\mathcal{U} = \mathcal{U}_0 + \mathcal{U}_1$ its Cartan decomposition, and (\cdot, \cdot) be an ad-invariant, non-degenerate bilinear form on \mathcal{U} .

Note that the U/U_0 -system (1.2) can be also defined invariantly as a system for maps $v : \mathcal{A} \rightarrow \mathcal{U}_1 \cap \mathcal{A}^\perp$ so that $[da, v]$ is flat, where da means the differential of the identity map $a(\xi) = \xi$ on \mathcal{A} . When we choose a basis

a_1, \dots, a_r of \mathcal{A} , the system becomes (1.2). Change basis of \mathcal{A} amounts to linear change of coordinates of \mathbb{R}^r .

The U/U_0 -system (1.2) and the curved flat system (1.3) are gauge equivalent. To explain this we first recall some known Propositions, which can be proved by direct computations.

Proposition 2.1. *The following statements are equivalent for smooth maps $u_i : \mathbb{R}^n \rightarrow \mathcal{G}$, $1 \leq i \leq n$:*

- (1) $\sum_{i=1}^n u_i dx_i$ is a flat \mathcal{G} -connection 1-form on \mathbb{R}^n ,
- (2) the first order system $E_{x_i} = Eu_i$, $1 \leq i \leq n$ is solvable,
- (3) there exists $g : \mathcal{O} \rightarrow G$ such that $g^{-1}dg = \sum_{i=1}^n u_i dx_i$ for some open subset \mathcal{O} of the origin in \mathbb{R}^n .

Proposition 2.2. ([12]) *The following statements are equivalent for $v : \mathbb{R}^n \rightarrow \mathcal{U}_1 \cap \mathcal{A}^\perp$:*

- (1) v is a solution of the U/U_0 -system (1.2),
- (2) $\sum_{i=1}^r [a_i, v] dx_i$ is a \mathcal{U} -valued flat connection on \mathbb{R}^r ,
- (3)

$$\theta_\lambda = \sum_{i=1}^r (a_i \lambda + [a_i, v]) dx_i \quad (2.1)$$

is a \mathcal{G} -valued flat connection on \mathbb{R}^r for all parameter $\lambda \in \mathbb{C}$,

- (4) there is an $s \in \mathbb{R}$ so that $\theta_s = \sum_{i=1}^r (a_i s + [a_i, v]) dx_i$ is a \mathcal{U} -valued flat connection on \mathbb{R}^r .

Proposition 2.3. *A smooth map $(A_1, \dots, A_r) : \mathbb{R}^r \rightarrow \prod_{i=1}^r \mathcal{U}_1$ is a solution of the curved flat system (1.3) associated to U/U_0 if and only if*

$$\omega_\lambda = \sum_{i=1}^r \lambda A_i dx_i$$

is a flat \mathcal{G} -valued connection 1-form on \mathbb{R}^r for all $\lambda \in \mathbb{C}$.

The flat connections θ_λ and ω_λ are called *Lax connections* of the U/U_0 -system and the U/U_0 curved flat system.

A map $\xi : \mathbb{C} \rightarrow \mathcal{G}$ is said to satisfy the *U/U_0 -reality condition* if

$$\tau(\xi(\bar{\lambda})) = \xi(\lambda), \quad \sigma(\xi(\lambda)) = \xi(-\lambda).$$

It follows from the definition that $\xi(\lambda) = \sum_j \xi_j \lambda^j$ satisfies the U/U_0 -reality condition if and only if $\xi_j \in \mathcal{U}_0$ if j is even and $\xi_j \in \mathcal{U}_1$ if j is odd. Note that both Lax connections θ_λ and ω_λ satisfy the U/U_0 -reality condition.

It follows from Proposition 2.1 that if v is a solution of the U/U_0 -system then there exists a unique $E(x, \lambda)$ so that

$$\begin{cases} E^{-1} E_{x_i} = a_i \lambda + [a_i, v], & 1 \leq i \leq r, \\ E(0, \lambda) = e. \end{cases}$$

Since θ_λ satisfies the U/U_0 -reality condition, E also satisfies the U/U_0 -reality condition:

$$\tau((E(x, \bar{\lambda})) = E(x, \lambda), \quad \sigma(E(x, \lambda)) = E(x, -\lambda).$$

We call such E the *parallel frame* of the Lax connection θ_λ associated to v .

The following Proposition says that solutions of the U/U_0 -system give rise to solutions of the curved flat system.

Proposition 2.4. ([15]) *Let $v : \mathbb{R}^r \rightarrow \mathcal{U}_1 \cap \mathcal{A}^\perp$ be a solution of the U/U_0 -system (1.2), and $E(x, \lambda)$ the parallel frame of the corresponding Lax connection θ_λ defined by (2.1). Let $g(x) = E(x, 0)$, and $A_i = ga_i g^{-1}$ for $1 \leq i \leq r$. Then*

(i) *the gauge transformation of θ_λ by g is*

$$g * \theta_\lambda = \sum_{i=1}^r \lambda ga_i g^{-1} dx_i,$$

(ii) (A_1, \dots, A_r) *is a solution of the curved flat system (1.3),*

Theorem 2.5. ([9]) *If (A_1, \dots, A_r) is a solution of the curved flat system (1.3) associated to U/U_0 , then there exists $f : \mathcal{O} \rightarrow U$ such that $f^{-1}f_{x_i} = A_i$ for all $1 \leq i \leq r$ and $\pi(f)$ is a curved flat in U/U_0 , where $\pi : U \rightarrow U/U_0$ is the natural projection. Conversely, every curved flat in U/U_0 can be lifted to a map f to U so that $(f^{-1}f_{x_1}, \dots, f^{-1}f_{x_r})$ is a solution of the curved flat system (1.3).*

A direct computation implies

Proposition 2.6. *If (A_1, \dots, A_r) is a solution of the curved flat system (1.3) associated to U/U_0 , then there exists a smooth map $f : \mathbb{R}^r \rightarrow U$ such that f satisfies the following conditions:*

$$\begin{cases} f^{-1}f_{x_i} \in \mathcal{U}_1, \\ [f^{-1}f_{x_i}, f^{-1}f_{x_j}] = 0, \quad \text{for all } i \neq j. \end{cases} \quad (2.2)$$

Conversely, if $f : \mathbb{R}^r \rightarrow U$ is an immersion satisfying (2.2), then

$$(f^{-1}f_{x_1}, \dots, f^{-1}f_{x_r})$$

is a solution of the curved flat system (1.3).

An immersed submanifold $f : \mathcal{O} \rightarrow \mathcal{U}_1$ is called *flat abelian* ([15]) if

- (1) $[f_{y_i}, f_{y_j}] = 0$ for all $1 \leq i \neq j \leq n$,
- (2) the induced metric on \mathcal{O} is flat.

The following Theorems give explicit algorithms to construct flat abelian submanifolds in \mathcal{U}_1 and curved flats in the symmetric space U/U_0 from solutions of the U/U_0 -system. The proofs can be found in [15].

Theorem 2.7. ([15]) *Let v and E be as in Proposition 2.4. Set $Y = \frac{\partial E}{\partial \lambda} E^{-1} \Big|_{\lambda=0}$. Then Y is an immersed flat abelian submanifold in \mathcal{U}_1 . Conversely, locally all flat abelian submanifolds in \mathcal{U}_1 can be constructed this way.*

For $g \in U$ and $x \in U$,

$$g * x = gx\sigma(g)^{-1}$$

defines an action of U on U (it is called the σ -action). The orbit at e is

$$M = \{g\sigma(g)^{-1} \mid g \in U\} \subset U.$$

Since the isotropy subgroup at e is U_0 , the orbit M is diffeomorphic to U/U_0 . In fact, M is a totally geodesic submanifold of U and is isometric to the symmetric space U/U_0 . This is the classical *Cartan embedding* of the symmetric space U/U_0 in U .

Theorem 2.8. ([15]) *With the same assumption as in Theorem 2.4, and set*

$$\psi(x) = E(x, 1)E(x, -1)^{-1}.$$

Then ψ is a curved flat in the symmetric space U/U_0 . Conversely, locally all curved flats in U/U_0 can be constructed this way.

Theorem 2.9. *Let \mathcal{O} be an open neighborhood of $0 \in \mathbb{R}^r$. If $f : \mathcal{O} \rightarrow U$ is an immersion satisfying (2.2), then there exists a local coordinate system x near 0 , a regular basis $\{a_1, \dots, a_r\}$ of the maximal abelian subspace $\mathcal{A} = \text{Im}(df_0)$, $g : \mathcal{O} \rightarrow U_0$, and a solution $v : \mathbb{R}^r \rightarrow \mathcal{U}_1 \cap \mathcal{A}^\perp$ of the U/U_0 -system (1.2) so that*

$$\begin{cases} f^{-1}f_{x_i} = ga_i g^{-1}, \\ g^{-1}g_{x_i} = [a_i, v]. \end{cases} \quad (2.3)$$

Conversely, if v is a solution of the U/U_0 -system, then $f(x) = E(x, 1)E(x, 0)^{-1}$ satisfies (2.2) and (2.3), where $E(x, \lambda)$ is the parallel frame for the Lax connection θ_λ corresponding to v .

Proof. Since generically all maximal abelian subalgebra are conjugate under elements of U_0 , there exist $g : \mathcal{O} \rightarrow U_0$ and $b_1, \dots, b_n : \mathcal{O} \rightarrow \mathcal{A}$ such that

$$f^{-1}f_{y_i} = gb_i g^{-1}$$

for $1 \leq i \leq n$. A direct computation implies

$$\begin{aligned} 0 &= d(df) = d\left(f \sum_{i=1}^n gb_i g^{-1} dy_i\right) \\ &= f \sum_{j \neq i} (g(b_i)_{y_j} g^{-1} + [g_{y_j} g^{-1}, gb_i g^{-1}]) dy_j \wedge dy_i \\ &= fg \left(\sum_{j \neq i} ((b_i)_{y_j} - [b_i, g^{-1}g_{y_j}]) dy_j \wedge dy_i \right) g^{-1}. \end{aligned}$$

This implies that

$$(b_i)_{y_j} - [b_i, g^{-1}g_{y_j}] = (b_j)_{y_i} - [b_j, g^{-1}g_{y_i}] \quad (2.4)$$

for all $i \neq j$. Let (\cdot, \cdot) denote a non-degenerate ad-invariant bilinear form on \mathcal{U} . Then $(\mathcal{A}, [\mathcal{A}, \mathcal{U}]) = 0$ and $[\mathcal{U}_{\mathcal{A}}, [\mathcal{A}, \mathcal{U}]] = 0$. So $[\mathcal{A}, \mathcal{U}] \subset \mathcal{U}_{\mathcal{A}}^{\perp}$ and $[\mathcal{A}, \mathcal{U}] \subset \mathcal{A}^{\perp}$. Also we have

$$\mathcal{U}_1 = \mathcal{A} \oplus (\mathcal{A}^{\perp} \cap \mathcal{U}_1).$$

Note $(b_i)_{y_j} \in \mathcal{A}$ and $[b_i, g^{-1}g_{y_j}] \in [\mathcal{A}, \mathcal{U}_0]$ is contained in \mathcal{A}^{\perp} . By (2.4) we get

$$(b_i)_{y_j} = (b_j)_{y_i}, \quad (2.5a)$$

$$[b_i, g^{-1}g_{y_i}] = [b_j, g^{-1}g_{y_i}], \quad (2.5b)$$

for all $1 \leq i \neq j \leq n$. Equation (2.5a) implies that $\sum_{i=1}^n b_i dy_i$ is closed. So there exist a local coordinate change $x = x(y)$ and constant a_1, \dots, a_n in \mathcal{A} such that $\sum_{i=1}^n b_i dy_i = \sum_{i=1}^n a_i dx_i$.

Let $\beta = \sum_{i=1}^n b_i dy_i$. Equation (2.5b) can be rewritten as $[\beta, g^{-1}dg] = 0$. Write β and $g^{-1}dg$ in x coordinate to get $\beta = \sum_{i=1}^n a_i dx_i$ and $g^{-1}dg = \sum_{i=1}^n g^{-1}g_{x_i} dx_i$. Then

$$0 = [\beta, g^{-1}dg] = \sum_{i \neq j} [a_i, g^{-1}g_{x_j}] dx_i \wedge dx_j.$$

So we have

$$[a_i, g^{-1}g_{x_j}] = [a_j, g^{-1}g_{x_i}], \quad \forall i \neq j.$$

Up to a linear change of coordinates of x , we may assume that a_i 's are regular. Note the kernel of $\text{ad}(a_i)$ on \mathcal{U}_1 is \mathcal{A} , and the tangent plane of the orbit $\text{Ad}(U_0)(a_i)$ at a_i is $[a_i, \mathcal{U}_0]$. By assumption $\text{Ad}(U_0)(\mathcal{A})$ is open in \mathcal{U}_1 . So the dimension of the tangent plane of the principal $\text{Ad}(U_0)$ -orbit at a_i is equal to $\dim(\mathcal{U}_1) - \dim(\mathcal{A})$. Thus $\text{ad}(a_i)$ maps $\mathcal{A}^{\perp} \cap \mathcal{U}_1$ isomorphically onto $\mathcal{U}_0 \cap (\mathcal{U}_0)_{\mathcal{A}}^{\perp}$, where $(\mathcal{U}_0)_{\mathcal{A}} = \{\xi \in \mathcal{U}_0 \mid [\xi, \mathcal{A}] = 0\}$. Then by (2.5b), there exists a $v : \mathcal{O} \rightarrow \mathcal{U}_1 \cap \mathcal{A}^{\perp}$ so that

$$g^{-1}g_{x_i} = [a_i, v], \quad 1 \leq i \leq n.$$

But $g^{-1}dg = \sum_i [a_i, v] dx_i$ is a flat connection. By Proposition 2.2, v is a solution of the U/U_0 -system (1.2).

To prove the converse, note that

$$E^{-1}dE = \theta_{\lambda} = \sum_{i=1}^r (a_i \lambda + [a_i, v]) dx_i.$$

Set $g(x) = E(x, 0)$ and $F(x, \lambda) = E(x, \lambda)E(x, 0)^{-1} = E(x, \lambda)g(x)^{-1}$. A direct computation implies that

$$F^{-1}dF = g\theta_{\lambda}g^{-1} - dgg^{-1} = \sum_{i=1}^r \lambda g a_i g^{-1} dx_i.$$

Note $f(x) = F(x, 1)$. So $f^{-1}df = \sum_{i=1}^r g a_i g^{-1} dx_i$. \square

Remark 2.10. The maps g and v in Theorem 2.9 are essentially unique. To see this, suppose we have g, \tilde{g} so that

$$ga_i g^{-1} = \tilde{g}a_i \tilde{g}^{-1} = f^{-1}f_{x_i},$$

$g^{-1}g_{x_i} = [a_i, v]$, and $\tilde{g}^{-1}\tilde{g}_{x_i} = [a_i, \tilde{v}]$. Since $g^{-1}\tilde{g}a_i = a_i$, there exists $(\mathcal{U}_0)_{\mathcal{A}}$ -valued map h such that $g^{-1}\tilde{g} = h$, i.e., $\tilde{g} = gh$. But

$$\begin{aligned} \tilde{g}^{-1}\tilde{g}_{x_i} &= [a_i, \tilde{v}] = h^{-1}[a_i, v]h + h^{-1}h_{x_i} \\ &= [a_i, h^{-1}vh] + h^{-1}h_{x_i} \in \mathcal{U}_{\mathcal{A}}^{\perp} + \mathcal{U}_{\mathcal{A}}. \end{aligned}$$

Thus

$$\begin{cases} h^{-1}h_{x_i} = 0, \\ [a_i, h^{-1}vh] = [a_i, \tilde{v}]. \end{cases}$$

The first equation implies h is a constant. Since $h^{-1}vh \in \mathcal{U}_1 \cap \mathcal{A}^{\perp}$ and $\text{ad}(a_i)$ is injective on $\mathcal{U}_1 \cap \mathcal{A}^{\perp}$, the second equation implies that $h^{-1}vh = \tilde{v}$. This proves that $\tilde{g} = gh$ and $\tilde{v} = h^{-1}vh$ for some constant $h \in (\mathcal{U}_0)_{\mathcal{A}}$.

3. BASICS OF EXTERIOR DIFFERENTIAL SYSTEMS

We give a brief account of Cartan-Kähler theory based on the lectures given by R. Bryant at MSRI in 1999 and 2003 (cf. [3] for details and references).

Let M be a smooth manifold, and $\Omega^*(M)$ the graded algebra of differential forms on M . An ideal \mathcal{I} of $\Omega^*(M)$ is called a *differential ideal* if \mathcal{I} satisfies the following conditions:

- (1) $\mathcal{I} = \bigoplus_j \mathcal{I}^j$, where $\mathcal{I}^j = \Omega^j(M) \cap \mathcal{I}$;
- (2) $d\mathcal{I} \subset \mathcal{I}$.

An *exterior differential system* (EDS) is a pair (M, \mathcal{I}) consisting of a smooth manifold M and a differential ideal $\mathcal{I} \subset \Omega^*(M)$.

A submanifold $N \subset M$ is called an *integral submanifold* for the EDS (M, \mathcal{I}) if $i^*\mathcal{I} = 0$, where $i : N \hookrightarrow M$ is the inclusion. In local coordinates this condition can be written as a system of PDE (or ODE).

A linear subspace $E \subset T_p M$ is said to be an *integral element* of \mathcal{I} if $\varphi|_E = 0$ for all $\varphi \in \mathcal{I}$. The set of all integral elements of \mathcal{I} of dimension n is denoted $v_n(\mathcal{I})$. A submanifold of M is an integral submanifold of \mathcal{I} if and only if each of its tangent space is an integral element of \mathcal{I} .

Note that $v_n(\mathcal{I}) \cap Gr_n(T_p M)$ is a real algebraic sub-variety of $Gr_n(T_p M)$, which may be very complicated. The set of *ordinary integral elements* $v_n^o(\mathcal{I}) \subset v_n(\mathcal{I})$ consists of those which are locally cut out 'cleanly' by finite number of n -forms in \mathcal{I} , so that the connected components of $v_n^o(\mathcal{I})$ are smooth embedded submanifolds of $Gr_n(TM)$. The rigorous definition can be found in [3].

Let $\{e_1, \dots, e_n\}$ be a basis of the linear subspace E of $T_p M$. The *polar space* of E is defined to be the vector space

$$H(E) = \{v \in T_p M \mid \varphi(v, e_1, \dots, e_n) = 0 \text{ for all } \varphi \in \mathcal{I}^{n+1}\}.$$

When $E \in v_n(\mathcal{I})$, a $(n+1)$ -plane E^+ containing E is an integral element of \mathcal{I} if and only if $E^+ \subset H(E)$. Define

$$r(E) = \dim H(E) - \dim E - 1.$$

This integer may jump up at certain points. An ordinary integral element E is called *regular* if r is locally constant in a neighborhood of E in $v_n^o(\mathcal{I})$. The set of regular integral elements is denoted $v_n^r(\mathcal{I})$ and is a dense open subset of $v_n^o(\mathcal{I})$. Thus $v_n^r(\mathcal{I}) \subset v_n^o(\mathcal{I}) \subset v_n(\mathcal{I}) \subset Gr_n(TM)$. An integral submanifold is called *regular* if all of its tangent spaces are regular integral elements.

We state the following two theorems that are given in [5]:

Theorem 3.1. (*Cartan-Kähler Theorem*) *Suppose (M, \mathcal{I}) is a real analytic EDS and that $N \subset M$ is a connected real analytic regular n -dimensional integral submanifold of \mathcal{I} with $r(N) \geq 0$. Let $R \subset M$ be a real analytic submanifold of codimension $r(N)$ containing N such that*

$$\dim(T_p R \cap H(T_p N)) = n + 1, \text{ for all } p \in N.$$

Then there exists a unique connected real analytic $(n+1)$ -dimensional integral submanifold \tilde{N} such that $N \subset \tilde{N} \subset R$.

A *regular flag* is a flag of integral elements

$$(0) = E_0 \subset E_1 \subset \dots \subset E_n = E \subset T_p M$$

where $E_j \in v_j^r(\mathcal{I})$ for $0 \leq j < n$ and $E_n \in v_n(\mathcal{I})$. Note that E_n may not be regular, but one can show that it must be ordinary. By applying Cartan-Kähler Theorem repeatedly to this flag, one can show that there is a real analytic n -dimensional integral manifold $N \subset M$ passing through p and satisfying $T_p N = E$. Set

$$c(E_j) = \dim(T_p M) - \dim H(E_j).$$

Theorem 3.2. (*Cartan's Test*) *Let (M, \mathcal{I}) be an EDS, and $F = (E_0, \dots, E_n)$ an integral flag of \mathcal{I} . Then $v_n(\mathcal{I})$ has codimension at least*

$$c(F) = c(E_0) + \dots + c(E_{n-1})$$

in $Gr_n(TM)$ at E_n . Moreover, F is a regular flag if and only if $v_n(\mathcal{I})$ is a smooth submanifold of $Gr_n(TM)$ in a neighborhood E_n and has codimension exactly $c(F)$.

The *Cartan characters* of the flag F are the numbers

$$s_j(F) = \dim H(E_{j-1}) - \dim H(E_j), \quad 0 \leq j \leq n,$$

with the convention $c(E_{-1}) = 0$ or $H(E_{-1}) = T_p M$. These numbers exhibit the generality of integral submanifolds. Roughly speaking, the integral manifolds near N will depend on s_0 constants, s_1 functions of one variable, \dots , s_n functions of n variables.

A connected open subset Z of $v_n^o(\mathcal{I})$ is called *involutive* if every $E \in Z$ is the terminus of a regular flag. When Z is clear from the context, we simply say that our EDS (M, \mathcal{I}) is involutive.

Suppose (M, \mathcal{I}) is an EDS with n -dimensional integral submanifold. A *conservation law* for (M, \mathcal{I}) is an $(n-1)$ -form $\phi \in \Omega^{n-1}(M)$ such that $d(f^*\phi) = 0$ for every integral submanifold $f : N^n \hookrightarrow M$ of \mathcal{I} . Actually, one only considers as conservation laws those ϕ such that $d\phi \in \mathcal{I}$. Two “trivial” type of conservation laws are $\phi \in \mathcal{I}^{n-1}$ or ϕ being exact on M . Factoring out these cases, *the space of conservation laws* is defined to be $\mathcal{C} = H^{n-1}(\Omega^*(M)/\mathcal{I})$. It also makes sense factor out those conservation laws represented by closed $(n-1)$ -forms on M (then the quotient space is called the space of *proper* conservation laws). One can study the symmetries of the EDS and then apply Noether’s Theorem to compute the corresponding conservation laws (cf. [4] for details).

4. INVOLUTIVITY OF THE EDS (U, \mathcal{I}_σ)

Let $G, \tau, \sigma, U, \mathcal{U}_0$ and \mathcal{U}_1 be as in section 1. Let α be the canonical left invariant 1-form $g^{-1}dg$ on U . Write

$$\alpha = \alpha_0 + \alpha_1$$

with respect to the Cartan decomposition $\mathcal{U} = \mathcal{U}_0 + \mathcal{U}_1$. The \mathcal{U}_j -component of the Maurer-Cartan equation $d\alpha + \frac{1}{2}[\alpha, \alpha] = 0$ gives

$$\begin{cases} d\alpha_0 + \frac{1}{2}([\alpha_0, \alpha_0] + [\alpha_1, \alpha_1]) = 0, \\ d\alpha_1 + [\alpha_0, \alpha_1] = 0. \end{cases}$$

Let $\mathcal{I}_\sigma \subset \Omega^*(U)$ be the differential ideal generated by the components of α_0 . It follows from the Maurer-Cartan equation that

$$\begin{aligned} \mathcal{I}_\sigma &= \langle \alpha_0, d\alpha_0 \rangle \\ &= \langle \alpha_0, [\alpha_1, \alpha_1] \rangle. \end{aligned}$$

Here $\langle \cdot, \cdot \rangle$ denotes the algebraic ideal generated by the enclosed forms.

The following Proposition was proved by R. Bryant.

Proposition 4.1. ([6]) *The EDS (U, \mathcal{I}_σ) is involutive.*

Proof. Since everything is homogeneous, we only need to look at the integral elements $E \subset T_e U = \mathcal{U}$. Note that $E \subset \mathcal{U}_1 = \bigcap_{j \neq 1} \ker(\alpha_j)$. For $E = (0) \in v_0(\mathcal{I}_\sigma)$, we have $H(E) = \mathcal{U}_1$ and $v_0(\mathcal{I}_\sigma) \cong U$. Thus $v_0(\mathcal{I}_\sigma) = v_0^o(\mathcal{I}_\sigma) = v_0^r(\mathcal{I}_\sigma)$. Now consider $E = \mathbb{R}x \in v_1(\mathcal{I}_\sigma)$ for some $x \in \mathcal{U}_1 - \{0\}$. Its polar space is

$$H(E) = \{y \in \mathcal{U}_1 \mid [x, y] = 0\},$$

since $[\alpha_1, \alpha_1]_e(x, y) = [x, y]$. For generic such x , $H(\mathbb{R}x)$ will be a maximal abelian subalgebra of \mathcal{U}_1 , and set $\dim H(\mathbb{R}x) = \dim \mathcal{A} = r$. Therefore

$$v_1(\mathcal{I}_\sigma) = v_1^o(\mathcal{I}_\sigma) \not\subseteq v_1^r(\mathcal{I}_\sigma).$$

Furthermore, when $\mathbb{R}x \in v_1^r(\mathcal{I}_\sigma)$, every subspace E of $H(\mathbb{R}x)$ containing $\mathbb{R}x$ is also regular and has $H(E) = H(\mathbb{R}x)$. Thus generic $E \in v_1^o(\mathcal{I}_\sigma)$ is the terminus of a regular flag, and our EDS is involutive. In fact, every regular integral curve of \mathcal{I}_σ lies in a unique r -dimensional integral submanifold of \mathcal{I}_σ , or locally the integral submanifolds depend on $s_0 = \dim \mathcal{U} - \dim \mathcal{U}_1$ constants and $s_1 = \dim(\mathcal{U}_1) - r$ functions of one variable (since $s_2 = \dots = s_r = 0$). \square

Corollary 4.2. *Let \mathcal{O} be an open subset of \mathbb{R}^r , and $f : \mathcal{O} \rightarrow U$ an immersion. Then the following statements are equivalent:*

- (1) f is a k -dimensional integral submanifold of (U, \mathcal{I}_σ) ,
- (2) f satisfies (2.2),
- (3) $(f^{-1}f_{x_1}, \dots, f^{-1}f_{x_r})$ is a solution of the curved flat system (1.3).

Hence the PDE for the EDS (U, \mathcal{I}_σ) is the curved flat system (1.3) associated to U/U_0 .

As a consequence of the Cartan Kähler Theorem 3.1, Proposition 4.1 and Corollary 4.2 that real analytic curved flats in U/U_0 or real analytic solutions of the curved flat systems (1.3) depend only on $\dim(\mathcal{U}_1 \cap \mathcal{A}^\perp)$ functions of one variable along a non-characteristic line.

By Theorem 2.9, there is a special coordinate system x on \mathbb{R}^r so that the curved flat system (1.3) written in x coordinate system is gauge equivalent to the U/U_0 -system (1.2). The Cartan-Kähler theory implies that the Cauchy problem of the U/U_0 -system has a unique local solution for any given local real analytic initial data on the x_1 -axis. But it was also proved in [12] using the inverse scattering theory of Beals and Coifman ([1]) that given any smooth rapidly decaying function $v_0 : \mathbb{R} \rightarrow \mathcal{U}_1 \cap \mathcal{A}^\perp$, there exists a unique smooth solution $v : \mathbb{R}^r \rightarrow \mathcal{U}_1 \cap \mathcal{A}^\perp$ so that

$$v(x_1, \dots, x_r) = v_0(x_1, 0, \dots, 0).$$

Although the theory exterior differential system seems to give a weaker result concerning the Cauchy problem, it may prove to be a very good tool to detect “integrability”.

Remark 4.3. Let G, τ, U be as above, and ρ an order k automorphism of G so that $d\rho_e$ is complex linear. Assume that

$$\tau\rho = \rho^{-1}\tau^{-1}.$$

Let \mathcal{G}_j denote the eigenspace of $d\rho_e$ with eigenvalue $e^{\frac{2\pi ij}{k}}$, and $\mathcal{K}_j = \mathcal{U} \cap \mathcal{G}_j$. Then

$$\mathcal{U} = \mathcal{K}_0 + \dots + \mathcal{K}_{k-1}. \quad (4.1)$$

Let $\alpha = g^{-1}dg$, and

$$\alpha = \alpha_0 + \dots + \alpha_{k-1}$$

the decomposition of α with respect to (4.1). Let \mathcal{I}_ρ denote the differential ideal on U generated by $\alpha_0, \alpha_2, \dots, \alpha_{k-1}$. Then

$$\begin{aligned}\mathcal{I}_\rho &= \langle \alpha_0, \alpha_2, \dots, \alpha_{k-1}, d\alpha_0, d\alpha_2, \dots, d\alpha_{k-1} \rangle \\ &= \langle \alpha_0, \alpha_2, \dots, \alpha_{k-1}, [\alpha_1, \alpha_1] \rangle.\end{aligned}$$

We define regular elements in \mathcal{K}_1 the same way as before, namely, $a \in \mathcal{K}_1$ is regular if it is contained in a maximal abelian subspace \mathcal{A} in \mathcal{K}_1 and $\text{Ad}(U_0)(\mathcal{A})$ is open in \mathcal{K}_1 . If \mathcal{K}_1 admits regular elements, then the proof of Proposition 4.1 works for (U, \mathcal{I}_ρ) . In fact, in this case, we have:

- (1) (U, \mathcal{I}_ρ) is involutive.
- (2) If $\dim(\mathcal{A}) = r$, then any r -dimensional integral submanifold depend on $\dim(\mathcal{K}_1) - \dim(\mathcal{A})$ number of functions of one variable.
- (3) every regular integral curve is contained in a unique r -dimensional integral submanifold of (U, \mathcal{I}_ρ) .
- (4) The curved flat system associated to U/K is the system (1.3) for $(A_1, \dots, A_r) : \mathbb{R}^r \rightarrow \mathcal{K}_1$, and the U/K -system is the system (1.2) for $v : \mathbb{R}^r \rightarrow \mathcal{K}_{k-1}$. Modulo a change of coordinate system of \mathbb{R}^r , these two system are gauge equivalent.
- (5) Given an immersion $f : \mathbb{R}^r \rightarrow U$, the following statements are equivalent:
 - (a) f is an integral submanifold of (U, \mathcal{I}_ρ) ,
 - (b) $f^{-1}f_{x_i} \in \mathcal{K}_1$ and $[f^{-1}f_{x_i}, f^{-1}f_{x_j}] = 0$ for all i, j ,
 - (c) $(f^{-1}f_{x_1}, \dots, f^{-1}f_{x_r})$ is a solution of the curved flat system associated to U/K .
- (6) The PDE for the EDS (U, \mathcal{I}_ρ) is the curved flat system associated to U/K .

5. CONSERVATION LAWS AND COMMUTING FLOWS

We construct infinitely many conservation laws and commuting flows for the U/U_0 -system, and indicate how to construct infinitely many explicit solutions of the U/U_0 -system.

First we review the Birkhoff Factorization Theorem (for detail see [10]). Let $\epsilon > 0$ be a small number, and $\mathcal{O}_\epsilon = \{\lambda \in \mathbb{C} \mid \frac{1}{\epsilon} < |\lambda| \leq \infty\}$ the open neighborhood at ∞ in $S^2 = \mathbb{C} \cup \{\infty\}$. $L(G)$ denote the group of holomorphic maps $g : \mathcal{O}_\epsilon \setminus \{\infty\} \rightarrow G$, $L_+(G)$ the subgroup of $g \in L(G)$ such that g can be extended to a holomorphic map in \mathbb{C} , and $L_-(G)$ the subgroup of $g \in L(G)$ that can be extended to a holomorphic map in \mathcal{O}_ϵ and is equal to the identity e at ∞ .

Theorem 5.1. (*Birkhoff Factorization Theorem*) *The multiplication map*

$$\mu : L_+(G) \times L_-(G) \rightarrow L(G), \quad (g_+, g_-) \mapsto g_+g_-$$

is one to one, and the image is an open dense subset of $L(G)$.

In other words, for generic $g \in L(G)$, we can factor $g = g_+g_-$ uniquely with $g_{\pm} \in L_{\pm}(G)$. Let \hat{e} denote the constant map from $\mathcal{O}_e \setminus \{\infty\}$ to G with constant e . Since \hat{e} lies in the image of the multiplication map μ , there is an open subset of \hat{e} so that all elements in this open subset can be factored uniquely.

Let $\tilde{\tau}$ and $\hat{\sigma}$ denote the map on $L(G)$ defined by

$$(\tilde{\tau}(g))(\lambda) = \tau(g(\bar{\lambda})), \quad (\hat{\sigma}(g))(\lambda) = \sigma(g(-\lambda)).$$

It is easy to check that

- (1) $\tilde{\tau}$ and $\hat{\sigma}$ are conjugate linear and complex linear involutions of $L(G)$,
- (2) $g \in L(G)$ is a fixed point of both $\tilde{\tau}$ and $\hat{\sigma}$ if and only if g satisfies the U/U_0 -reality condition: $\tau(g(\bar{\lambda})) = g(\lambda)$, $\sigma(g(\lambda)) = \sigma(-\lambda)$.
- (3) both $\tilde{\tau}$ and $\hat{\sigma}$ leave $L_{\pm}(G)$ invariant.

Let $L^{\tau,\sigma}(G)$ and $L_{\pm}^{\tau,\sigma}(G)$ denote the subgroup of fixed points of $\tilde{\tau}$ and $\hat{\sigma}$ of $L(G)$ and $L_{\pm}(G)$ respectively. Then we have

Corollary 5.2. *The multiplication map*

$$L_+^{\tau,\sigma}(G) \times L_-^{\tau,\sigma}(G) \rightarrow L^{\tau,\sigma}(G)$$

is one to one and the image is open and dense in $L^{\tau,\sigma}(G)$.

We want to use this factorization to construct infinitely many solutions and commuting flows for the U/U_0 -system. Given $b \in \mathcal{A}$ and $j > 1$ an odd integer, $x \in \mathbb{R}^r$, and $t \in \mathbb{R}$, let $e^A(x, t) \in L_+^{\tau,\sigma}(G)$ be defined by

$$e^A(x, t)(\lambda) = \exp((a_1x_1 + \cdots + a_r x_r)\lambda + b\lambda^j t).$$

Given $f \in L_-^{\tau,\sigma}(G)$, since $e^A(0, 0) = \hat{e}$ is the identity in $L^{\tau,\sigma}(G)$ and e^A is smooth from $\mathbb{R}^r \times \mathbb{R}$ to $L^{\tau,\sigma}(G)$, by Corollary 5.2 there is an open subset of $(0, 0)$ in $\mathbb{R}^r \times \mathbb{R}$ so that we can factor $f^{-1}e^A(x, t)$ uniquely as

$$f^{-1}e^A(x, t) = E(x, t)m(x, t)^{-1}, \quad (5.1)$$

where $E(x, t) \in L_+^{\tau,\sigma}(G)$ and $m(x, t) \in L_-^{\tau,\sigma}(G)$.

Given $c \in \mathcal{A}$, let

$$m^{-1}cm = Q_{c,0} + Q_{c,1}\lambda^{-1} + Q_{c,2}\lambda^{-2} + \cdots \quad (5.2)$$

denote the Taylor series of $(m(x, t)^{-1}cm(x, t))(\lambda)$ at $\lambda = \infty$. Since $m(x, t)(\lambda) = e$ at $\lambda = \infty$,

$$Q_{c,0} = c. \quad (5.3)$$

We want to explain how to compute $Q_{c,n}$, $m^{-1}dm$ and $E^{-1}dE$ next. To do this, we take ∂_{x_i} of (5.1) to get

$$f^{-1}e^A(x, t)a_i\lambda = E_{x_i}m^{-1} - Em^{-1}m_{x_i}m^{-1}.$$

Multiply E^{-1} on the left and m on the right of the above equation and use (5.1) to get

$$m^{-1}a_i\lambda m = E^{-1}E_{x_i} - m^{-1}m_{x_i}. \quad (5.4)$$

Take ∂_t of (5.1) and use a similar calculation to get

$$m^{-1}b\lambda^j m = E^{-1}E_t - m^{-1}m_t. \quad (5.5)$$

Note that $E^{-1}E_{x_i}$ and $E^{-1}E_t$ lie in the Lie algebra $\mathcal{L}_+^{\tau,\sigma}(\mathcal{G})$ of $L_+^{\tau,\sigma}(G)$, and $m^{-1}m_{x_i}$ and $m^{-1}m_t$ lie in the Lie algebra $\mathcal{L}_-^{\tau,\sigma}(\mathcal{G})$ of $L_-^{\tau,\sigma}(G)$. But it follows from the factorization theorem that

$$\mathcal{L}(\mathcal{G}) = \mathcal{L}_+^{\tau,\sigma}(\mathcal{G}) \oplus \mathcal{L}_-^{\tau,\sigma}(\mathcal{G})$$

as direct sum of vector spaces. Let ξ_{\pm} denote the $\mathcal{L}_{\pm}^{\tau,\sigma}(\mathcal{G})$ component of $\xi \in \mathcal{L}^{\tau,\sigma}(\mathcal{G})$. Then (5.4) and (5.5) imply that

$$E^{-1}E_{x_i} = (m^{-1}a_i m \lambda)_+, \quad (5.6a)$$

$$E^{-1}E_t = (m^{-1}b m \lambda^j)_+ \quad (5.6b)$$

$$m^{-1}m_{x_i} = -(m^{-1}a_i m \lambda)_-, \quad (5.6c)$$

$$m^{-1}m_t = -(m^{-1}b m \lambda^j)_-. \quad (5.6d)$$

Use (5.2) to see that

$$\begin{aligned} (m^{-1}a_i m \lambda)_+ &= Q_{a_i,0}\lambda + Q_{a_i,1}, \\ (m^{-1}b m \lambda^j)_+ &= Q_{b,0}\lambda^j + Q_{b,1}\lambda^{j-1} + \cdots + Q_{b,j}. \end{aligned}$$

So we get

$$\begin{cases} E^{-1}E_{x_i} = Q_{a_i,0}\lambda + Q_{a_i,1}, \\ E^{-1}E_t = Q_{b,0}\lambda^j + Q_{b,1}\lambda^{j-1} + \cdots + Q_{b,j}. \end{cases} \quad (5.7)$$

Lemma 5.3. *If $c_1, c_2 \in \mathcal{A}$, then*

$$[m^{-1}c_1 m, m^{-1}c_2 m] = 0, \quad (5.8a)$$

$$[m^{-1}c_1 m, -(m^{-1}c_2 \lambda^n m)_-] = [m^{-1}c_1 m, (m^{-1}c_2 \lambda^n m)_+]. \quad (5.8b)$$

Theorem 5.4. ([12]) *There exists $v : \mathbb{R}^r \times \mathbb{R} \rightarrow \mathcal{U}_1 \cap \mathcal{A}^{\perp}$ so that*

$$Q_{a_i,1} = [a_i, v].$$

Moreover, for each $t \in \mathbb{R}$, $v(\cdots, t)$ is a solution of the U/U_0 -system.

Proof. By (5.8a),

$$[m^{-1}a_i m, m^{-1}a_j m] = 0, \quad 1 \leq i \neq j \leq r.$$

So the coefficient of λ^{-1} of the left hand side has to be zero, i.e.,

$$[a_i, Q_{a_j,1}] + [Q_{a_i,1}, a_j] = 0, \quad 1 \leq i \neq j \leq r.$$

But this implies that there exists $v : \mathbb{R}^r \times \mathbb{R} \rightarrow \mathcal{U}_1 \cap \mathcal{A}^{\perp}$ so that

$$Q_{a_i,1} = [a_i, v].$$

By (5.7) and Proposition 2.1 we see that $\sum_i (a_i \lambda + [a_i, v]) dx_i$ is a flat \mathcal{G} -valued connection on \mathbb{R}^r for all $\lambda \in \mathbb{C}$. Hence for each fixed t , $v(\cdots, t)$ is a solution of the U/U_0 -system. \square

The following is well-known (cf. [11, 13]):

Theorem 5.5. (1) $Q_{b,j}(x, t)$ is a polynomial in $u, \partial_x v, \dots, \partial_x^{j-1} v$,
 (2) $Q_{b,j}$ satisfies the following recursive formula

$$(Q_{b,j})_{x_i} + [[a_i, v], Q_{b,j}] = [Q_{b,j+1}, a_i], \quad (5.9)$$

(3) $Q_{b,0} = b, Q_{b,1} = [b, v]$.

Proof. A direct computation gives

$$\begin{aligned} (m^{-1}bm)_{x_i} &= [m^{-1}bm, m^{-1}m_{x_i}], \quad \text{by (5.6a)} \\ &= [m^{-1}bm, -(m^{-1}a_i\lambda m)_-], \quad \text{by (5.8b)} \\ &= [m^{-1}bm, (m^{-1}a_i\lambda m)_+]. \end{aligned}$$

Substitute (5.2) to the above equation to get

$$(m^{-1}bm)_{x_i} = [m^{-1}bm, a_i\lambda + u]. \quad (5.10)$$

Compare coefficient of λ^{-j} of λ^{-j} of (5.10) to get (5.9).

It was proved in [11, 13] that $Q_{b,j}$ is a polynomial in $u_i, \partial_{x_i} u_i, \dots, \partial_{x_i}^{j-1} u_i$, where $u_i = [a_i, v]$. Since $\text{ad}(a_i)$ is a linear isomorphism between $\mathcal{U}_1 \cap \mathcal{A}^\perp$ and $\mathcal{U}_0 \cap \mathcal{U}_A^\perp$, (1) follows. Since $m(\dots, \infty) = I$, $Q_{b,0} = b$. Use (5.9) to prove $Q_{b,1} = [b, v]$. \square

Use (5.7) and Proposition 2.1 to see that

$$\Theta_\lambda^{b,j} = \sum_{i=1}^r (a_i\lambda + [a_i, v]) dx_i + (b\lambda^j + Q_{b,1}\lambda^{j-1} + \dots + Q_{b,j}) dt$$

is a flat connection on $\mathbb{R}^r \times \mathbb{R}$ for all $\lambda \in \mathbb{C}$. It follows from the recursive formula (5.9) and the flat equation

$$d\Theta_\lambda^{b,j} + \Theta_\lambda^{b,j} \wedge \Theta_\lambda^{b,j} = 0$$

that we have

$$\begin{cases} [a_i, v_{x_j}] = [a_j, v_{x_i}] + [[a_i, v], [a_j, v]], & i \neq j, \\ [a_i, v_t] = (Q_{b,j})_x + [[a_i, v], Q_{b,j}], & 1 \leq i \leq r. \end{cases} \quad (5.11)$$

The first set of equations just means $v(\dots, t)$ is a solution of the U/U_0 -system for each t , and the second set of equations give the flow on the space of solutions of the U/U_0 -system.

Let $A_{\mathbb{C}}$ denote the subgroup of G whose Lie subalgebra is $\mathcal{A} \otimes \mathbb{C}$, and $L_+^{\tau,\sigma}(A_{\mathbb{C}})$ is the subgroup of $f \in L_+^{\tau,\sigma}(G)$ such that $g(\lambda) \in A_{\mathbb{C}}$ for all $\lambda \in \mathbb{C}$. Given $b \in \mathcal{A}$ and j a positive integer, then $\xi_{b,j}$ lies in the Lie algebra $\mathcal{L}_+^{\tau,\sigma}(\mathcal{A} \otimes \mathbb{C})$, where $\xi_{b,j}(\lambda) = b\lambda^j$. Let $e_{b,j}(t)$ be the one-parameter subgroup in $L_+^{\tau,\sigma}(G)$ generated by $\xi_{b,j}$, i.e.,

$$e_{b,j}(t)(\lambda) = e^{b\lambda^j t}.$$

It was proved in [13] that if $v(x, t)$ is a solution of (5.11) then

$$e_{b,j}(t) \cdot v(\dots, 0) := v(\dots, t)$$

is the dressing action of $e_{b,j}(t) \in L_+^{\tau,\sigma}(A_{\mathbb{C}})$ on the space of solutions of the U/U_0 -systems. The second set of equations of (5.11) is the vector field on the space of solutions of the U/U_0 -system corresponding to the one-parameter subgroup generated by $\xi_{b,j}$. Since the group $L_+^{\tau,\sigma}(A_{\mathbb{C}})$ is abelian, the flows generated by these $\xi_{b,j}$ are commuting. So we have

Theorem 5.6. ([12], [13]) *Given $b \in \mathcal{A}$ and a positive integer j , the flow*

$$[a_i, v_t] = (Q_{b,j})_{x_i} + [[a_i, v], Q_{b,j}], \quad 1 \leq i \leq r, \quad (5.12)$$

leaves the space of solutions of the U/U_0 -system (1.2) invariant. Moreover, all these flows commute.

We sketch the method of constructing solutions of the U/U_0 -system below. Let $e^{A_0}(x) \in L_+^{\tau,\sigma}(G)$ be defined by

$$e^{A_0}(x) = \exp\left(\sum_{i=1}^r a_i x_i \lambda\right).$$

Theorem 5.7. ([14]) *Given $f \in L_-^{\tau,\sigma}(G)$, factor*

$$f^{-1}e^{A_0}(x) = E(x)m^{-1}(x) \quad (5.13)$$

with $E(x) \in L_+^{\tau,\sigma}(G)$ and $m(x) \in L_-^{\tau,\sigma}(G)$. Expand $m(x)(\lambda)$ at $\lambda = \infty$:

$$m(x)(\lambda) = e + m_{-1}(x)\lambda^{-1} + m_2(x)\lambda^{-2} + \dots.$$

Then

- (1) $m_{-1}(x) \in \mathcal{U}_1$,
- (2) $v = (m_{-1})^\perp$ is a solution of the U/U_0 -system, where $(m_{-1})^\perp$ is the projection of m_{-1} onto $\mathcal{U}_1 \cap \mathcal{A}^\perp$ with respect to $\mathcal{U}_1 = \mathcal{A} \oplus (\mathcal{U}_1 \cap \mathcal{A}^\perp)$.

Proof. Use the same computation as for the proof of (5.6a) to conclude that

$$E^{-1}E_{x_i} = (m^{-1}a_i m)_+ = a_i \lambda + Q_{a_i,1}.$$

Expand $m(x)(\lambda)$ at $\lambda = \infty$:

$$m(x)(\lambda) = e + m_{-1}(x)\lambda^{-1} + m_{-2}(x)\lambda^{-2} + \dots.$$

A direct computation implies that

$$m^{-1}a_i m = a_i + [a_i, m_{-1}]\lambda^{-1} + \dots.$$

Therefore $Q_{a_i,1} = [a_i, m_{-1}]$. Since $m \in L^{\tau,\sigma}(G)$, $m_{-1}(x) \in \mathcal{U}_1$. So

$$[a_i, v] = [a_i, m_{-1}^\perp] = [a_i, m_{-1}] = Q_{a_i,1}.$$

Hence we have shown that

$$E^{-1}E_{x_i} = a_i \lambda + [a_i, v], \quad 1 \leq i \leq r.$$

By Proposition 2.2, v is a solution of the U/U_0 -system. □

Remark 5.8. It was proved in [14] that if each entry of $f \in L_-^{\tau,\sigma}(G)$ is a meromorphic function on $S^2 = \mathbb{C} \cup \{\infty\}$, then the factorization (5.13) can be carried out explicitly using residue calculus. In particular, $m(x)(\lambda)$ and $E(x, \lambda) = E(x)(\lambda)$ can be given by explicit formulas. Therefore, we get explicit solutions $v = (m_{-1})^\perp$ for the U/U_0 -system. Since the parallel frame $E(x, \lambda)$ for the solution v is also given explicitly, it follows from Corollary 4.2 and Theorem 2.9 that $F(x) = E(x, 1)E(x, 0)^{-1}$ is an explicit integral submanifold of the EDS (U, \mathcal{I}_σ) .

Next we derive conservation laws of the flows for the U/U_0 -system.

Theorem 5.9. *Let $c, b \in \mathcal{A}$, and n a positive integer. Then*

$$(Q_{c,n}, a_i)_{x_j} = (Q_{c,n}, a_j)_{x_i}, \quad 1 \leq i \neq j \leq r. \quad (5.14)$$

In particular,

$$\phi_{c,n} := \sum_{i=1}^r (Q_{c,n}, a_i) dx_i \quad (5.15)$$

is a closed 1-form on \mathbb{R}^r

Proof. Compute directly to get

$$\begin{aligned} (m^{-1}cm, a_i)_{x_j} &= ([m^{-1}cm, m^{-1}m_{x_j}], a_i), \quad \text{by (5.6c),} \\ &= ([m^{-1}cm, -(m^{-1}a_j\lambda m)_-], a_i), \quad \text{by (5.8b),} \\ &= ([m^{-1}cm, (m^{-1}a_j\lambda m)_+], a_i) = ([m^{-1}cm, a_j\lambda + Q_{a_j,1}], a_i). \end{aligned}$$

Use (5.2) and compare coefficient of λ^{-n} of the above equation to get

$$\begin{aligned} (Q_{c,n}, a_i)_{x_j} &= ([Q_{c,n}, Q_{a_j,1}], a_i) + ([Q_{c,n+1}, a_j], a_i) \\ &= (Q_{c,n}, [Q_{a_j,1}, a_i]) + (Q_{c,n+1}, [a_j, a_i]) = (Q_{c,n}, [[a_j, v], a_i]) + 0 \\ &= (Q_{c,n}, [[a_i, v], a_j]) = ([Q_{c,n}, [a_i, v]], a_j), \quad \text{by (5.9),} \\ &= ((Q_{c,n})_{x_i} - [Q_{n+1, a_i}], a_j) = ((Q_{c,n})_{x_i}, a_j). \end{aligned}$$

□

If $f : \mathcal{O} \rightarrow U$ is a r -dimensional integral submanifold of the EDS (U, \mathcal{I}_σ) , then by Theorem 2.9 and Corollary 4.2 there exist a special local coordinate system x of \mathcal{O} , $g : \mathcal{O} \rightarrow U_0$ and a solution v of the U/U_0 -system (1.2) such that $f^{-1}f_{x_i} = ga_i g^{-1}$ and $g^{-1}g_{x_i} = [a_i, v]$ for all $1 \leq i \leq r$. Let $*$ denote the Hodge star operator for the Euclidean space \mathbb{R}^r . Given $1 \leq i \neq j \leq r$, let

$$\psi_{c,n}^{ij} = \phi_{c,n} \wedge *(dx_i \wedge dx_j) = \left(\sum_{\ell=1}^r (Q_{c,n}, a_\ell) dx_\ell \right) \wedge *(dx_i \wedge dx_j).$$

Then $\psi_{c,n}^{ij}$ is a closed $(r-1)$ -form on the integrable submanifold. In other words, $\psi_{c,n}^{ij}$ is a conservation law for the EDS (U, \mathcal{I}_σ) for all $1 \leq i < j \leq r$, $c \in \mathcal{A}$, and positive integer n .

Next we derive the conservation laws for the flow (5.12) on the space of solutions of the U/U_0 -system (1.2). Given $a, c \in \mathcal{A}$, compute

$$\begin{aligned} (m^{-1}cm, a)_t &= ([m^{-1}cm, m^{-1}m_t], a), \quad \text{by (5.6d)} \\ &= ([m^{-1}cm, -(m^{-1}b\lambda^j m)_-], a), \quad \text{by (5.8b)} \\ &= ([m^{-1}cm, (m^{-1}b\lambda^j m)_+], a). \end{aligned}$$

Substitute (5.2) to the above equation and compare coefficient of λ^{-n} to get

$$((Q_{c,n})_t, a) = \sum_{i=0}^{j-1} ([Q_{c,n+i}, Q_{b,j-i}], a). \quad (5.16)$$

Here we have used

$$([Q_{c,n}, Q_{b,0}], a) = ([Q_{c,n}, b], a) = (Q_{c,n}, [b, a]) = 0.$$

We claim that

$$([Q_{c,n}, Q_{b,j}], a_i) = \sum_{i=1}^j (Q_{c,n+i-1}, Q_{b,j-i})_{x_i} \quad (5.17)$$

We prove this claim by induction on j . For $j = 1$, we have

$$\begin{aligned} (Q_{c,n}, Q_{b,1}, a_i) &= (Q_{c,n}, [Q_{b,1}, a_i]) = (Q_{c,n}, [[b, v], a_i]), \\ &= (Q_{c,n}, [[a_i, v], b]) = -([[a_i, v], Q_{c,n}], b) \\ &= -([Q_{c,n+1}, a_i] - (Q_{c,n})_{x_i}, b) = ((Q_{c,n})_{x_i}, b). \end{aligned}$$

(We used the Jacobi identity for the first line of the computation above). This proves (5.17) for $j = 1$. Now assume (5.17) is true for j and we want to prove the identity for $j + 1$. We compute

$$\begin{aligned} ([Q_{c,n}, Q_{b,j+1}], a_i) &= (Q_{c,n}, [Q_{b,j+1}, a_i]), \quad \text{by (5.9)} \\ &= (Q_{c,n}, (Q_{b,j})_{x_i} + [u_i, Q_{b,j}]) \\ &= (Q_{c,n}, Q_{b,j})_{x_i} - ((Q_{c,n})_{x_i}, Q_{b,j}) + (Q_{c,n}, [u_i, Q_{b,j}]) \\ &= (Q_{c,n}, Q_{b,j})_{x_i} - ((Q_{c,n})_{x_i}, Q_{b,j}) - ([u_i, Q_{c,n}], Q_{b,j}) \\ &= (Q_{c,n}, Q_{b,j})_{x_i} + ([a, Q_{c,n+1}], Q_{b,j}), \quad \text{by (5.9),} \\ &= (Q_{c,n}, Q_{b,j})_{x_i} + (a, [Q_{c,n+1}, Q_{b,j}]) \end{aligned}$$

Then the induction hypothesis implies (5.17) is true for $j + 1$.

It follows from (5.16) and (5.17) that we have

Theorem 5.10. *Let $v : \mathbb{R}^r \times \mathbb{R} \rightarrow \mathcal{U}_1 \cap \mathcal{A}^\perp$ be a solution of (5.11), $c \in \mathcal{A}$, and n a positive integer. Then*

$$(Q_{c,n}, a_i)_t = \sum_{\ell=0}^{j-1} \sum_{s=1}^{j-\ell} (Q_{c,n+\ell+s-1}, Q_{b,j-\ell-s})_{x_i}. \quad (5.18)$$

As a consequence, we see that

$$\int_{\mathbb{R}^r} (Q_{c,n}, a_i) dx_1 \wedge \cdots \wedge dx_r$$

is a conserved quantity for the flow (5.12) on the space of rapidly decaying solutions of the U/U_0 -system.

REFERENCES

- [1] Beals, R., Coifman, R.R., *Scattering and inverse scattering for first order systems*, Commun. Pure Appl. Math. **37** (1984), 39–90
- [2] Brück, M., Du, X., Park, J., and Terng, C.L., *Submanifold geometry of real Grassmannian systems*, The Memoirs, vol 155, AMS, **735** (2002), 1–95
- [3] Bryant, R., Chern, S.S., Gardner, R., Goldschmidt, H., and Griffiths, P., *Exterior Differential Systems*, Springer-Verlag, New York (1991)
- [4] Bryant, R., Griffiths, P., and Grossman, D., *Exterior Differential Systems and Euler-Lagrange Partial Differential Equations*, Chicago Lectures in Mathematics, the University of Chicago Press, 2003.
- [5] Bryant, R., *Lectures given at MSRI summer school, 1999*, unpublished notes.
- [6] Bryant, R., *Lectures given at MSRI “Integrable system seminar”, 2003*, unpublished notes.
- [7] Burstall, F., *Isothermic surfaces: conformal geometry, Clifford algebras and integrable systems*, to appear in “Integrable systems, Geometry, and Topology”, International Press, math-dg/0003096
- [8] Ferus, D., Pedit, F., *Curved flats in symmetric spaces*, Manuscripta Math., **91** (1996), 445–454
- [9] Ferus, D., Pedit, F., *Isometric immersions of space forms and soliton theory*, Math. Ann., **305** (1996), 329–342
- [10] Pressley, A., Segal, G. B., *Loop Groups*, Oxford Science Publ., Clarendon Press, Oxford, (1986)
- [11] Sattinger, D.H., *Hamiltonian hierarchies on semi-simple Lie algebras*, Stud. Appl. Math., **72** (1984), 65–86
- [12] Terng, C.L., *Soliton equations and differential geometry*, J. Differential Geometry, **45** (1997), 407–445
- [13] Terng, C.L., Uhlenbeck, K., *Poisson actions and scattering theory for integrable systems*, Surveys in Differential Geometry: Integrable systems (A supplement to J. Differential Geometry), **4** (1998), 315–402
- [14] Terng, C.L., Uhlenbeck, K., *Bäcklund transformations and loop group actions*, Comm. Pure. Appl. Math., **53** (2000), 1–75
- [15] Terng, C.L., *Geometries and symmetries of soliton equations and integrable elliptic systems*, to appear in Surveys on Geometry and Integrable Systems, Advanced Studies in Pure Mathematics, Mathematical Society of Japan

NORTHEASTERN UNIVERSITY AND UC IRVINE

E-mail address: terng@neu.edu

MSRI, BERKELEY, CA 94720

E-mail address: ewang@mrsi.org