



A Quick Proof of the Classification of Simple Real Lie Algebras

Author(s): A. W. Knap

Source: *Proceedings of the American Mathematical Society*, Vol. 124, No. 10 (Oct., 1996), pp. 3257-3259

Published by: American Mathematical Society

Stable URL: <http://www.jstor.org/stable/2161875>

Accessed: 10/06/2009 03:25

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=ams>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We work with the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact support@jstor.org.



American Mathematical Society is collaborating with JSTOR to digitize, preserve and extend access to *Proceedings of the American Mathematical Society*.

<http://www.jstor.org>

A QUICK PROOF OF THE CLASSIFICATION OF SIMPLE REAL LIE ALGEBRAS

A. W. KNAPP

(Communicated by Roe W. Goodman)

ABSTRACT. Élie Cartan's classification of the simple Lie algebras over \mathbb{R} is derived quickly from some structure theory over \mathbb{R} and the classification over \mathbb{C} .

Élie Cartan classified the simple Lie algebras over \mathbb{R} for the first time in 1914. There have been a number of simplifications in the proof since then, and these are described in [3, p. 537]. All proofs assume the classification over \mathbb{C} and a certain amount of structure theory over \mathbb{R} . Recent proofs tend to run to 25 pages. Here is a shorter argument.

Theorem. *Up to isomorphism, the only simple Lie algebras over \mathbb{R} that are neither complex nor compact are those in Cartan's list as organized in [3, p. 518].*

We use terminology as in [3]. Let $\mathfrak{g}_0 = \mathfrak{k}_0 \oplus \mathfrak{p}_0$ be a Cartan decomposition of a noncomplex simple Lie algebra over \mathbb{R} , and let θ be the Cartan involution. Choose a maximal abelian subspace \mathfrak{t}_0 of \mathfrak{k}_0 and extend to a maximally compact Cartan subalgebra $\mathfrak{h}_0 = \mathfrak{t}_0 \oplus \mathfrak{a}_0$ of \mathfrak{g}_0 . Removal of subscripts 0 will indicate complexifications. Let $\Delta = \Delta(\mathfrak{g}, \mathfrak{h})$ be the root system. Roots are imaginary on \mathfrak{t}_0 and real on \mathfrak{a}_0 . All roots are imaginary-valued or complex on \mathfrak{h}_0 ; there are no real-valued roots. Introduce a positive system Δ^+ that takes $i\mathfrak{t}_0$ before \mathfrak{a}_0 . The map θ carries roots to roots and permutes the simple roots. The complex simple roots move in two-element orbits, while the imaginary simple roots are fixed. By the **Diagram** of $(\mathfrak{g}_0, \mathfrak{h}_0, \Delta^+)$, we mean the Dynkin diagram of Δ with the two-element orbits under θ so labeled and with the imaginary roots shaded or not, according as the simple root is noncompact (root vector in \mathfrak{p}) or compact (root vector in \mathfrak{k}).

Lemma 1. *If $(\mathfrak{g}_0, \mathfrak{h}_0, \Delta^+)$ and $(\mathfrak{g}'_0, \mathfrak{h}'_0, (\Delta')^+)$ have the same Diagram, then \mathfrak{g}_0 and \mathfrak{g}'_0 are isomorphic.*

Proof. We may assume that the complexifications $(\mathfrak{g}, \mathfrak{h}, \Delta^+)$ are the same and that the associated compact forms are the same: $\mathfrak{u}_0 = \mathfrak{k}_0 \oplus i\mathfrak{p}_0 = \mathfrak{k}'_0 \oplus i\mathfrak{p}'_0$. Using the conjugacy of compact forms, the conjugacy of maximal abelian subspaces within them, and the standard construction of a compact form from \mathfrak{h} , we see that we can normalize root vectors X_α , $\alpha \in \Delta$, as in Theorem 5.5 of [3, p. 176] and obtain \mathfrak{u}_0 from $\{X_\alpha\}$ as in Theorem 6.3 of [3, p. 181].

Received by the editors April 12, 1995.

1991 *Mathematics Subject Classification.* Primary 17B20, 22E15.

First suppose $\mathfrak{a}_0 = 0$, so that all roots are imaginary. For α simple we have $\theta X_\alpha = \pm X_\alpha$, the sign being $+$ if α is compact and $-$ if α is noncompact. The same formula holds for θ' . Since \mathfrak{h} and the X_α 's for α simple generate \mathfrak{g} , it follows that $\theta = \theta'$, hence that $\mathfrak{k} = \mathfrak{k}'$ and $\mathfrak{p} = \mathfrak{p}'$. Then $\mathfrak{g}_0 = \mathfrak{g}'_0$ is recovered as $(\mathfrak{u}_0 \cap \mathfrak{k}) \oplus i(\mathfrak{u}_0 \cap \mathfrak{p})$.

If $\mathfrak{a}_0 \neq 0$, we may not have $\theta = \theta'$. For $\alpha \in \Delta$, write $\theta X_\alpha = a_\alpha X_{\theta\alpha}$. Then $a_\alpha a_{-\alpha} = 1$ and $a_\alpha a_{\theta\alpha} = 1$. Since θ maps $\mathfrak{u}_0 \cap \text{span}\{X_\alpha, X_{-\alpha}\}$ to $\mathfrak{u}_0 \cap \text{span}\{X_{\theta\alpha}, X_{-\theta\alpha}\}$, we see that $\bar{a}_\alpha = a_{-\alpha}$. Therefore $|a_\alpha| = 1$. For each pair of complex simple roots α and $\theta\alpha$, choose square roots $a_\alpha^{1/2}$ and $a_{\theta\alpha}^{1/2}$ whose product is 1. Similarly write $\theta' X_\alpha = b_\alpha X_{\theta'\alpha}$ with $|b_\alpha| = 1$, and define $b_\alpha^{1/2}$ and $b_{\theta'\alpha}^{1/2}$ for α and $\theta'\alpha$ simple. Define H and H' in $\mathfrak{h} \cap \mathfrak{u}_0$ by the conditions that $\alpha(H) = \alpha(H') = 0$ for α simple imaginary and that $\exp(\frac{1}{2}\alpha(H)) = a_\alpha^{1/2}$, $\exp(\frac{1}{2}\theta\alpha(H)) = a_{\theta\alpha}^{1/2}$, $\exp(\frac{1}{2}\alpha(H')) = b_\alpha^{1/2}$, and $\exp(\frac{1}{2}\theta'\alpha(H')) = b_{\theta'\alpha}^{1/2}$ if α and $\theta\alpha$ are complex simple. A little computation shows that $\theta' \circ \text{Ad}(\exp \frac{1}{2}(H - H')) = \text{Ad}(\exp \frac{1}{2}(H - H')) \circ \theta$, from which it follows that $\mathfrak{k}' = \text{Ad}(\exp \frac{1}{2}(H - H'))\mathfrak{k}$, $\mathfrak{p}' = \text{Ad}(\exp \frac{1}{2}(H - H'))\mathfrak{p}$, and $\mathfrak{g}'_0 = \text{Ad}(\exp \frac{1}{2}(H - H'))\mathfrak{g}_0$. □

The next step is to identify some pairs of distinct Diagrams that correspond merely to changes of Δ^+ . The argument is inspired by [2]. First let us assume that $\mathfrak{a}_0 = 0$, i.e., that the automorphism of Δ given by θ is the identity. Let Λ be the subset of $i\mathfrak{t}_0$ where all roots take integer values and where all noncompact roots take odd-integer values. If $\{\omega_j\}$ is the basis dual to the simple roots, then the sum of those ω_j corresponding to the noncompact simple roots is a member of Λ . The set Λ is discrete, and we let H_0 be a member of Λ as close to 0 as possible.

Lemma 2. *If $(\Delta^+)'$ is a positive system that makes H_0 dominant, then there is at most one noncompact simple root, say α_i . If the basis dual to the simple roots of $(\Delta^+)'$ is $\{\omega_j\}$, then there cannot exist i' such that $\langle \omega_i - \omega_{i'}, \omega_{i'} \rangle > 0$.*

Proof. Since H_0 is in Λ and is dominant, $H_0 = \sum n_j \omega_j$ with all n_j integers ≥ 0 . If $n_i > 0$, then $H_0 - \omega_i$ is dominant and thus has $\langle H_0 - \omega_i, \omega_i \rangle \geq 0$ with equality if and only if $H_0 = \omega_i$. Then $|H_0 - 2\omega_i|^2 \leq |H_0|^2$ with equality only if $H_0 = \omega_i$, and minimality forces $H_0 = \omega_i$. Now let $H_0 = \omega_i$. If $\langle \omega_i - \omega_{i'}, \omega_{i'} \rangle > 0$, then $|H_0 - 2\omega_{i'}|^2 < |H_0|^2$, in contradiction to minimality. □

When $\mathfrak{a}_0 \neq 0$, Lemma 2 is to be applied to the part of $i\mathfrak{t}_0$ corresponding to the span of the imaginary simple roots. The result is that we can associate to any \mathfrak{g}_0 at least one Diagram in which at most one imaginary root is shaded.

Now we can read off the possibilities. First suppose that the automorphism of Δ is the identity. If all roots are unshaded, then \mathfrak{g}_0 is the compact form. Otherwise exactly one simple root is shaded. For the classical Dynkin diagrams, let the double line or triple point be at the right end, and let the i^{th} root be shaded. In A_n , we are led to $\mathfrak{su}(i, n + 1 - i)$. In B_n , we are led to $\mathfrak{so}(2i, 2n + 1 - 2i)$. In C_n , we are led to $\mathfrak{sp}(i, n - i)$ if $i < n$ and to $\mathfrak{sp}(n, \mathbb{R})$ if $i = n$. In D_n , we are led to $\mathfrak{so}(2i, 2n - 2i)$ if $i \leq n - 2$ and to $\mathfrak{so}^*(2n)$ otherwise.

For the exceptional Dynkin diagrams, a little checking that compares the second conclusion of Lemma 2 with the fundamental weights (see [1, pp. 260-275]) shows that α_i in Lemma 2 has to be a node (endpoint vertex) of the Dynkin diagram. Moreover, in G_2 , α_i has to be the long simple root, while in E_8 , it cannot be the node on the short branch. In E_6 two nodes are equivalent by outer automorphism. Thus we obtain at most three Lie algebras for E_7 ; at most two for E_6 , E_8 , F_4 ; and

at most one for G_2 . These are E II, E III for E_6 , E V, E VI, E VII for E_7 ; E VIII, E IX for E_8 ; F I, F II for F_4 ; and G for G_2 .

When the automorphism of Δ is not the identity, the Dynkin diagram is A_n , D_n , or E_6 . For A_n , there is no imaginary simple root if n is even, and there is one if n is odd. For n even we are led to $\mathfrak{sl}(n+1, \mathbb{R})$, while for n odd we are led to $\mathfrak{sl}(n+1, \mathbb{R})$ if the root is shaded and to $\mathfrak{su}^*(n+1)$ if the root is unshaded. For D_n , the first $n-2$ simple roots are imaginary. If all are unshaded, we are led to $\mathfrak{so}(1, 2n-1)$. If the i^{th} simple root is shaded, $i \leq n-2$, we are led to $\mathfrak{so}(2i+1, 2n-2i-1)$. For E_6 , the triple point and the node on the short branch are imaginary. If neither is shaded, we are led to E IV, while if either one is shaded, we are led to E I.

Note added in proof. David Vogan has pointed out that any Dynkin diagram marked with an involution and having a subset of its one-element orbits shaded is a Diagram for some \mathfrak{g}_0 . The proof is in the spirit of Lemma 1. Existence of the exceptional simple real Lie algebras follows.

REFERENCES

1. N. Bourbaki, *Groupes et Algèbres de Lie, Chapitres 4, 5, et 6*, Hermann, Paris, 1968. MR **39**:1590
2. A. Borel and J. De Siebenthal, *Les sous-groupes fermés de rang maximum des groupes de Lie clos*, Comment. Math. Helvetici **23** (1949), 200–221. MR **11**:326d
3. S. Helgason, *Differential Geometry, Lie Groups, and Symmetric Spaces*, Academic Press, New York, 1978. MR **80k**:53081

DEPARTMENT OF MATHEMATICS, STATE UNIVERSITY OF NEW YORK, STONY BROOK, NEW YORK 11794

E-mail address: aknapp@ccmail.sunysb.edu