

I: LIE SYMMETRIES OF PARTIAL DIFFERENTIAL EQUATIONS AND CR GEOMETRY

JOËL MERKER

Table of contents

1. Completely integrable systems of partial differential equations	1.
2. Submanifold of solutions	6.
3. Classification problems	10.
4. Punctual and infinitesimal Lie symmetries	12.
5. Examples	15.
6. Transfer of Lie symmetries to the parameter space	18.
7. Equivalence problems and normal forms	25.
8. Study of two specific examples	27.
9. Dual system of partial differential equations	45.
10. Fundamental pair of foliations and covering property	46.
11. Formal and smooth equivalences	50.

This memoir is divided in three parts¹. Part I endeavours a general, new theory (inspired by modern CR geometry) of Lie symmetries of completely integrable PDE systems, viewed from their associated submanifold of solutions. Part II builds general combinatorial formulas for the prolongations of vector fields to jet spaces. Part III characterizes explicitly flatness of some systems of second order. The results presented here are original and did not appear in print elsewhere; most formulas of Parts II and III were checked by means of Maple Release 7.

§1. COMPLETELY INTEGRABLE SYSTEMS OF PARTIAL DIFFERENTIAL EQUATIONS

1.1. General systems. Let $\mathbb{K} = \mathbb{R}$ or \mathbb{C} . Let $n \in \mathbb{N}$ with $n \geq 1$ and let $x = (x^1, \dots, x^n) \in \mathbb{K}^n$. Also, let $m \in \mathbb{N}$ with $m \geq 1$ and let $y = (y^1, \dots, y^m) \in \mathbb{K}^m$. For $\alpha \in \mathbb{N}^n$, we denote by a subscript y_{x^α} the partial derivative $\partial^{|\alpha|} y / \partial x^\alpha$ of a local map $\mathbb{K}^n \ni x \mapsto y(x) \in \mathbb{K}^m$.

Let $\kappa \in \mathbb{N}$ with $\kappa \geq 1$, let $p \in \mathbb{N}$ with $p \geq 1$, choose a collection of p multiindices $\beta(1), \dots, \beta(p) \in \mathbb{N}^n$ with $|\beta(q)| \geq 1$ for $q = 1, \dots, p$ and $\max_{1 \leq q \leq p} |\beta(q)| = \kappa$, and choose integers $j(1), \dots, j(p)$ with $1 \leq j(q) \leq m$ for $q = 1, \dots, p$. In the present Part I, we study the Lie symmetries of a general system of analytic partial differential equations of the form:

$$(\mathcal{E}) \quad y_{x^\alpha}^j(x) = F_\alpha^j \left(x, y(x), (y_{x^{\beta(q)}}^{j(q)}(x))_{1 \leq q \leq p} \right),$$

where j with $1 \leq j \leq m$ and $\alpha \in \mathbb{N}^n$ satisfy

$$(1.2) \quad (j, \alpha) \neq (j, 0) \quad \text{and} \quad (j, \alpha) \neq (j(q), \beta(q)).$$

In particular, all $(\kappa+1)$ -th partial derivatives of the unknown $y = y(x)$ depend on a certain precise set of derivatives of order $\leq \kappa$: the system is *complete*. In addition, all the other partial derivatives of order $\leq \kappa$ do also depend on the same precise set of derivatives.

Here, we assume that $u = 0$ is a local solution of the system (\mathcal{E}) and that the functions F_α^j are \mathbb{K} -algebraic (in the sense of Nash) or \mathbb{K} -analytic, in a neighborhood of the origin in \mathbb{K}^{n+m+p} . Even if our concern will be local throughout, we will not introduce any special notation to speak of open subsets and simply refer to various \mathbb{K}^μ . We will study five concrete instances, the first three ones being classical.

Date: 2008-2-2.

¹ Part II of [Me2005a] already appeared as [Me2005b].

Example 1.3. With $n = m = \kappa = 1$, a second order ordinary differential equation

$$(\mathcal{E}_1) \quad y_{xx} = F(x, y, y_x),$$

and more generally $y_{x^{\kappa+1}} = F(x, y, y_x, \dots, y_x^\kappa)$, where $x, y \in \mathbb{K}$, see [Lie1883, EL1890, Tr1896, Ca1924, Se1931, Ca1932a, Ol1986, Ar1988, BK1989, GTW1989, HK1989, Ib1992, Ol1995, N2003].

Example 1.4. With $n \geq 2$, $m = 1$ and $\kappa = 1$, a complete system of second order equations

$$(\mathcal{E}_2) \quad y_{x^{i_1}x^{i_2}} = F_{i_1, i_2}(x^{i_1}, y, y_x^{i_2}), \quad 1 \leq i_1, i_2 \leq n,$$

see [Ha1937, Ch1975, Su2001] and Part III below.

Example 1.5. Dually, with $n = 1$, $m \geq 2$ and $\kappa = 1$, an ordinary system of second order

$$(\mathcal{E}_3) \quad y_{xx}^j = F^j(x, y^{j_1}, y_x^{j_1}), \quad j = 1, \dots, m,$$

see [Fe1995, Me2004] and the references therein.

Example 1.6. With $n = 1$, $m = 2$ and $\kappa = 1$, a system of the form

$$(\mathcal{E}_4) \quad \begin{cases} y_x^2 = F(x, y^1, y^2, y_x^1) \\ y_{xx}^1 = G(x, y^1, y^2, y_x^1). \end{cases}$$

Differentiating the first equation with respect to x and substituting, we get the missing equation:

$$(1.7) \quad \begin{aligned} y_{xx}^2 &= F_x + y_x^1 F_{y^1} + y_x^2 F_{y^2} + y_{xx}^1 F_{y_x^1} \\ &= F_x + y_x^1 F_{y^1} + y_x^2 F_{y^2} + G F_{y_x^1} \\ &=: H(x, y^1, y^2, y_x^1). \end{aligned}$$

Example 1.8. With $n = 2$, $m = 1$ and $\kappa = 2$, a system of the form

$$(\mathcal{E}_5) \quad \begin{cases} y_{x^2} = F(x^1, x^2, y, y_{x^1}, y_{x^1 x^1}) \\ y_{x^1 x^1 x^1} = G(x^1, x^2, y, y_{x^1}, y_{x^1 x^1}). \end{cases}$$

Here, five equations are missing. Differentiating the first equation with respect to x^1 and substituting:

$$(1.9) \quad \begin{aligned} y_{x^1 x^2} &= F_{x^1} + y_{x^1} F_y + y_{x^1 x^1} F_{y_{x^1}} + y_{x^1 x^1 x^1} F_{y_{x^1 x^1}} \\ &= F_{x^1} + y_{x^1} F_y + y_{x^1 x^1} F_{y_{x^1}} + G F_{y_{x^1 x^1}} \\ &=: H(x^1, x^2, y, y_{x^1}, y_{x^1 x^1}), \end{aligned}$$

and then similarly for $y_{x^2 x^2}$, $y_{x^1 x^1 x^2}$, $y_{x^1 x^2 x^2}$, $y_{x^2 x^2 x^2}$.

1.10. Finitely nondegenerate generic submanifolds of \mathbb{C}^{n+m} . Examples 1.3, 1.4, 1.6 and 1.8 (but *not* 1.5) are intrinsically linked to real submanifolds of complex submanifolds.

Let M be a real algebraic or analytic local generic CR^2 submanifold of \mathbb{C}^{n+m} of codimension $m \geq 1$ and of CR dimension $n \geq 1$, and let $p \in M$. Classically, there exists local holomorphic coordinates $t = (z, w) \in \mathbb{C}^n \times \mathbb{C}^m$ centered at p in which M is represented by

$$(1.11) \quad w^j = \overline{\Theta}^j(z, \bar{z}, \bar{w}), \quad j = 1, \dots, m,$$

² Fundamentals about Cauchy-Riemann geometry may be found in [Bo1991, BER1999, Me2005a, Me2005b, MP2005].

for some local \mathbb{C} -analytic map $\Theta = (\Theta^1, \dots, \Theta^m)$ satisfying the identity

$$(1.12) \quad w \equiv \bar{\Theta}(z, \bar{z}, \Theta(\bar{z}, z, w)),$$

reflecting the fact that M is real.

Definition 1.13. ([BER1999, Me2005a, Me2005b, MP2005]) M is *finitely nondegenerate* if there exists an integer $\kappa \geq 1$ such that the local holomorphic map

$$(1.14) \quad (\bar{z}, \bar{w}) \longmapsto (\bar{\Theta}_{z\beta}^j(0, \bar{z}, \bar{w}))_{\substack{1 \leq j \leq m \\ |\beta| \leq \kappa}}$$

is of rank $n + m$ at $(\bar{z}, \bar{w}) = (0, 0)$.

From (1.12), the map $\bar{w} \mapsto \bar{\Theta}(0, 0, \bar{w})$ is already of rank m at $\bar{w} = 0$. One then verifies ([BER1999, Me2005a, Me2005b, MP2005]) that there exist multiindices $\beta(1), \dots, \beta(n) \in \mathbb{N}^n$ with $|\beta(k)| \geq 1$ for $k = 1, \dots, n$ and $\max_{1 \leq k \leq n} |\beta(k)| = \kappa$ together with integers $j(1), \dots, j(n)$ with $1 \leq j(k) \leq m$ such that the local holomorphic map

$$(1.15) \quad \mathbb{C}^{n+m} \ni (\bar{z}, \bar{w}) \longmapsto \left((\bar{\Theta}^j(0, \bar{z}, \bar{w}))_{1 \leq j \leq m}, (\bar{\Theta}_{z\beta(k)}^{j(k)}(0, \bar{z}, \bar{w}))_{1 \leq k \leq n} \right) \in \mathbb{C}^{m+n}$$

is of rank $n + m$ at $(\bar{z}, \bar{w}) = (0, 0)$.

1.16. Associated system of partial differential equations. Generalizing an idea which goes back to B. Segre in [Se1931, Se1932] ($n = m = 1$), applied by É. Cartan in [Ca1932a] and studied more recently in [Su2001, GM2003a], we may associate to M a system of partial differential equations of the form (\mathcal{E}) as follows. Complexifying the variables \bar{z} and \bar{w} , we introduce new independent variables $\zeta \in \mathbb{C}^n$ and $\xi \in \mathbb{C}^m$ together with the complex algebraic or analytic m -codimensional submanifold \mathcal{M} of $\mathbb{C}^{2(n+m)}$ defined by

$$(1.17) \quad w^j = \bar{\Theta}^j(z, \zeta, \xi), \quad j = 1, \dots, m.$$

We then consider the “dependent variables” w^j as algebraic or analytic functions of the “independent variables” z^k , with additional dependence on the extra “parameters” (ζ, ξ) . Then by applying the differentiation $\partial^{|\alpha|}/\partial z^\alpha$ to (1.17), we get $w_{z^\alpha}^j(z) = \bar{\Theta}_{z^\alpha}^j(z, \zeta, \xi)$. Assuming finite nondegeneracy and writing these equations for $(j, \alpha) = (j(k), \beta(k))$, we obtain a system of $m + n$ equations:

$$(1.18) \quad \begin{cases} w^j(z) = \bar{\Theta}^j(z, \zeta, \xi), & j = 1, \dots, m, \\ w_{z^{\beta(k)}}^{j(k)}(z) = \bar{\Theta}_{z^{\beta(k)}}^{j(k)}(z, \zeta, \xi), & k = 1, \dots, n. \end{cases}$$

By means of the implicit function theorem we can solve:

$$(1.19) \quad (\zeta, \xi) = R(z^k, w^j(z), w_{z^{\beta(k)}}^{j(k)}(z)).$$

Finally, for every pair (j, α) different from $(j, 0)$ and from $(j(k), \beta(k))$, we may replace (ζ, ξ) by R in the differentiated expression $w_{z^\alpha}^j(z) = \bar{\Theta}_{z^\alpha}^j(z, \zeta, \xi)$, which yields

$$(1.20) \quad \begin{aligned} w_{z^\alpha}^j(z) &= \bar{\Theta}_{z^\alpha}^j \left(z, R(z^k, w^j(z), w_{z^{\beta(k)}}^{j(k)}(z)) \right) \\ &=: F_\alpha^j \left(z^k, w^j(z), w_{z^{\beta(k)}}^{j(k)}(z) \right). \end{aligned}$$

This is the *system of partial differential equations associated to M* .

Example 1.21. (Continued) With $n = m = 1$, i.e. $M \subset \mathbb{C}^2$ and $\kappa = 1$, i.e. M is Levi nondegenerate of equation

$$(1.22) \quad w = \bar{w} + i z \bar{z} + O_3,$$

where z, \bar{z} are assigned weight 1 and w, \bar{w} weight 2, B. Segre [Se1931] obtained $w_{zz} = F(z, w, w_z)$. J. Faran [Fa1980] found some examples of such equations that *cannot* come from a $M \subset \mathbb{C}^2$. But the following was left unsolved.

Open problem 1.23. Characterize equations $y_{xx} = F(x, y, y_x)$ associated to a real analytic, Levi nondegenerate (i.e. $\kappa = 1$) hypersurface $M \subset \mathbb{C}^2$. Can one read the reality condition (1.12) on F ? In case of success, generalize to arbitrary $M \subset \mathbb{C}^{n+m}$.

Example 1.24. (Continued) Similarly, the system (\mathcal{E}_2) comes from a Levi nondegenerate hypersurface $M \subset \mathbb{C}^{n+1}$ ([Ha1937, CM1974, Ch1975, Su2001]). Exercise: why (\mathcal{E}_3) cannot come from any $M \subset \mathbb{C}^\nu$?

Example 1.25. (Continued) With $n = 1, m = 2$ and $\kappa = 1$, the system (\mathcal{E}_4) comes from a $M \subset \mathbb{C}^3$ which is Levi nondegenerate and satisfies

$$(1.26) \quad T^c M + [T^c M, T^c M] + [T^c M, [T^c M, T^c M]] = TM$$

at the origin, namely which has equations of the following form, after some elementary transformations ([Be1997, BES2005]):

$$(1.27) \quad \begin{aligned} w^1 &= \bar{w}^1 + i z \bar{z} + O_4, \\ w^2 &= \bar{w}^2 + i z \bar{z}(z + \bar{z}) + O_4, \end{aligned}$$

where z, \bar{z} are assigned weight 1 and $w^1, w^2, \bar{w}^1, \bar{w}^2$ weight 2.

Example 1.28. (Continued) With $n = 2, m = 1$ and $\kappa = 2$, the system (\mathcal{E}_5) comes from a hypersurface $M \subset \mathbb{C}^3$ of equation ([Eb1998, GM2003b, FK2005a, FK2005b, Eb2006, GM2006]):

$$(1.29) \quad w = \bar{w} + i \frac{2 z^1 \bar{z}^1 + z^1 z^1 \bar{z}^2 + \bar{z}^1 \bar{z}^1 z^2}{1 - z^2 \bar{z}^2} + O_4,$$

where $z_1, \bar{z}_1, z_2, \bar{z}_2$ are assigned weight 1 and w, \bar{w} weight 2, with the assumption that the Levi form has rank exactly one at every point, and with the assumption that M is 2-nondegenerate at 0.

1.30. Jet spaces, contact forms and Frobenius integrability. Throughout the present Part I, we assume that the system (\mathcal{E}) is *completely integrable*, namely that the Pfaffian system naturally associated to (\mathcal{E}) in the appropriate jet space is involutive in the sense of Frobenius. This holds automatically in case (\mathcal{E}) comes from a generic submanifold $M \subset \mathbb{C}^{n+m}$. In general, we will construct a *submanifold of solutions* associated to (\mathcal{E}) . So, we must explain complete integrability.

We denote by $\mathcal{J}_{n,m}^\kappa$ the space of κ -th jets of maps $\mathbb{K}^n \ni x \mapsto y(x) \in \mathbb{K}^m$. Let

$$(1.31) \quad (x^i, y^j, y_{i_1}^j, y_{i_1, i_2}^j, \dots, y_{i_1, i_2, \dots, i_\kappa}^j) \in \mathbb{K}^{n+m+mn+mn^2+\dots+mn^\kappa},$$

denote the natural coordinates on $\mathcal{J}_{n,m}^\kappa \simeq \mathbb{K}^{n+m(1+n+\dots+n^\kappa)}$. For instance, $(x, y, y_1) \in \mathcal{J}_{1,1}^1$. We shall sometimes write them shortly:

$$(1.32) \quad (x^i, y^j, y_\beta^j) \in \mathbb{K}^{n+m+m(n+\dots+n^\kappa)},$$

where $\beta \in \mathbb{N}^n$ varies and satisfies $|\beta| \leq \kappa$. Sometimes also, we consider these jet coordinates only up to their symmetries $y_{i_1, i_2, \dots, i_\lambda}^j = y_{i_{\sigma(1)}, i_{\sigma(2)}, \dots, i_{\sigma(\lambda)}}^j$, where σ is a permutation of $\{1, 2, \dots, \lambda\}$, so that $\mathcal{J}_{n,m}^\kappa \simeq \mathbb{K}^{n+m} C_{n+\kappa}^\kappa$, with $C_{n+\kappa}^\kappa := \frac{(n+\kappa)!}{\kappa! n!}$.

Having these notations at hand, we may develop the canonical system of contact forms on $\mathcal{J}_{n,m}^\kappa$ ([OI1995], [Stk2000]):

$$(1.33) \quad \left\{ \begin{array}{l} \theta^j := dy^j - \sum_{k=1}^n y_k^j dx^k, \\ \theta_{i_1}^j := dy_{i_1}^j - \sum_{k=1}^n y_{i_1,k}^j dx^k, \\ \dots\dots\dots \\ \theta_{i_1,\dots,i_{\kappa-1}}^j := dy_{i_1,\dots,i_{\kappa-1}}^j - \sum_{k=1}^n y_{i_1,\dots,i_{\kappa-1},k}^j dx^k. \end{array} \right.$$

For instance, with $n = m = 1$ and $\kappa = 2$, we have $\theta^1 = dy - y_1 dx$ and $\theta_1^1 = dy_1 - y_2 dx$. These (linearly independent) one-forms generate a subspace $\mathcal{CT}_{n,m}^\kappa$ of the cotangent $T^*\mathcal{J}_{n,m}^\kappa$ whose dimension equals $m C_{n+\kappa-1}^{\kappa-1}$. For the duality between forms and vectors, the orthogonal $(\mathcal{CT}_{n,m}^\kappa)^\perp$ in $T\mathcal{J}_{n,m}^\kappa$ is spanned by the $n + m C_{n+\kappa-1}^\kappa$ vector fields:

$$(1.34) \quad \left\{ \begin{array}{l} D_i := \frac{\partial}{\partial x^i} + \sum_{j_1=1}^m y_{i_1}^{j_1} \frac{\partial}{\partial y^{j_1}} + \dots + \sum_{j_1=1}^m \sum_{k_1,\dots,k_{\kappa-1}=1}^n y_{i_1,k_1,\dots,k_{\kappa-1}}^{j_1} \frac{\partial}{\partial y_{k_1,\dots,k_{\kappa-1}}^{j_1}}, \\ T_{i_1,\dots,i_\kappa}^{j_1} := \frac{\partial}{\partial y_{i_1,\dots,i_\kappa}^{j_1}}, \end{array} \right.$$

the first n ones being the total differentiation operators, considered in Part II. For $n = m = 1, \kappa = 2$, we get $\frac{\partial}{\partial x} + y_1 \frac{\partial}{\partial y} + y_2 \frac{\partial}{\partial y_1}$ and $\frac{\partial}{\partial y_2}$.

Classically ([OI1986, BK1989, OI1995]), one associates to (\mathcal{E}) its *skeleton* $\Delta_{\mathcal{E}}$, namely the $(n + m + p)$ -dimensional submanifold of $\mathcal{J}_{n,m}^{\kappa+1}$ simply defined by the graphed equations:

$$(1.35) \quad y_\alpha^j = F_\alpha^j(x, y, (y_{\beta(q)}^{j(q)})_{1 \leq q \leq p}),$$

for $(j, \alpha) \neq (j, 0)$ and $\neq (j(q), \beta(q))$ with $|\alpha| \leq \kappa + 1$. Clearly, the natural coordinates on $\Delta_{\mathcal{E}}$ are:

$$(1.36) \quad (x, y, (y_{\beta(q)}^{j(q)})_{1 \leq q \leq p}) \equiv (x, y, (y_{l_1(q), \dots, l_{\lambda_q}(q)}^{j(q)})_{1 \leq q \leq p}) \in \mathbb{K}^n \times \mathbb{K}^m \times \mathbb{K}^p,$$

where $\lambda_q := |\beta(q)|$ and $(l_1(q), \dots, l_{\lambda_q}(q)) := \beta(q)$.

Next, in view of the form (1.34) of the generators of $(\mathcal{CT}_{n,m}^{\kappa+1})^\perp$ and in view of the equations of $\Delta_{\mathcal{E}}$, the intersection

$$(1.37) \quad (\mathcal{CT}_{n,m}^{\kappa+1})^\perp \cap T\Delta_{\mathcal{E}}$$

is a vector subbundle of $T\Delta_{\mathcal{E}}$ that is generated by n linearly independent vector fields obtained by restricting the D_i to $\Delta_{\mathcal{E}}$, which yields:

$$(1.38) \quad \left\{ \begin{array}{l} D_i = \frac{\partial}{\partial x^i} + \sum_{j=1}^m \mathbf{A}_i^j(x^{i_1}, y^{j_1}, y_{\beta(q_1)}^{j(q_1)}) \frac{\partial}{\partial y^j} + \\ + \sum_{q=1}^p \mathbf{B}_i^q(x^{i_1}, y^{j_1}, y_{\beta(q_1)}^{j(q_1)}) \frac{\partial}{\partial y_{\beta(q)}^{j(q)}}, \end{array} \right.$$

$i = 1, \dots, n$, where the coefficients \mathbf{A}_i^j and \mathbf{B}_i^q are given by:

$$(1.39) \quad \begin{aligned} \mathbf{A}_i^j &:= \begin{cases} y_i^j & \text{if the variable } y_i^j \text{ appears among the } p \text{ variables } y_{\beta^{(q_1)}}^{j(q_1)}; \\ F_i^j & \text{otherwise;} \end{cases} \\ \mathbf{B}_i^q &:= \begin{cases} y_{i, l_1(q), \dots, l_{\lambda_q}(q)}^{j(q)} & \text{if } y_{l_1(q), \dots, l_{\lambda_q}(q)}^{j(q)} \text{ appears among the } p \text{ variables } y_{\beta^{(q_1)}}^{j(q_1)}; \\ F_{i, l_1(q), \dots, l_{\lambda_q}(q)}^{j(q)} & \text{otherwise.} \end{cases} \end{aligned}$$

Example 1.40. For (\mathcal{E}_1) , we get $D = \frac{\partial}{\partial x} + y_1 \frac{\partial}{\partial x} + F(x, y, y_1) \frac{\partial}{\partial y_2}$; exercise: treat (\mathcal{E}_2) and (\mathcal{E}_3) . For (\mathcal{E}_4) , we get $D = \frac{\partial}{\partial x} + y_1^1 \frac{\partial}{\partial y^1} + F \frac{\partial}{\partial y^2} + G \frac{\partial}{\partial y_1^1}$. For (\mathcal{E}_5) , whose skeleton is written $y_2 = F$, $y_{1,1,1} = G$, $y_{1,2} = H$, $y_{1,1,2} = K$, with F, G, H, K being functions of $(x^1, x^2, y, y_1, y_{1,1})$, we get


$$(1.41) \quad \begin{aligned} D_1 &= \frac{\partial}{\partial x^1} + y_1 \frac{\partial}{\partial y} + y_{1,1} \frac{\partial}{\partial y_1} + G \frac{\partial}{\partial y_{1,1}}, \\ D_2 &= \frac{\partial}{\partial x^2} + F \frac{\partial}{\partial y} + H \frac{\partial}{\partial y_1} + K \frac{\partial}{\partial y_{1,1}}. \end{aligned}$$

Definition 1.42. The system (\mathcal{E}) is *completely integrable* if the n vector fields (1.38) satisfy the Frobenius integrability condition, namely every Lie bracket $[D_{i_1}, D_{i_2}]$, $1 \leq i_1, i_2 \leq n$, is a linear combination of the vector fields D_1, \dots, D_n .

Because of their specific form (1.38), we must then have in fact $[D_{i_1}, D_{i_2}] = 0$. For $n = 1$, the condition is of course void.

§2. SUBMANIFOLD OF SOLUTIONS

2.1. Fundamental foliation of the skeleton. As the vector fields D_i commute, they equip the skeleton $\Delta_{\mathcal{E}} \simeq \mathbb{K}^{n+m+p}$ with a foliation $F_{\Delta_{\mathcal{E}}}$ by n -dimensional integral manifolds which are (approximately) directed along the x -axis. We draw a diagram (see only the left side).



The (abstract, not numerical) integration of (\mathcal{E}) is thus straightforwardly completed: the set of solutions coincides with the set of leaves of $F_{\Delta_{\mathcal{E}}}$. This is the true geometric content, viewed in the appropriate jet space, of the assumption of complete integrability.

2.2. General solution and submanifold of solutions. To construct the submanifold of solutions $\mathcal{M}_{(\mathcal{E})}$ associated to (\mathcal{E}) (sketched in the right hand side), we execute some elementary analytico-geometric constructions.

At first, we *duplicate* the coordinates $(y_{\beta(q)}^{j(q)}, y^j) \in \mathbb{K}^p \times \mathbb{K}^m$ by introducing a new subspace of coordinates $(a, b) \in \mathbb{K}^p \times \mathbb{K}^m$; thus, on the left diagram, we draw a vertical plane together with a - and b -axes. The leaves of the foliation $F_{\Delta_{\mathcal{E}}}$ are uniquely determined by their intersections with this plane, consisting of points of coordinates $(0, a, b) \in \mathbb{K}^p \times \mathbb{K}^m$.

Such points $(0, a, b)$ correspond to the *initial conditions* $(y_{x\beta(q)}^{j(q)}(0), y(0))$ for the general solution of (\mathcal{E}) . In fact, the (concatenated, multiple) flow of $\{D_1, \dots, D_n\}$ is given by

$$(2.3) \quad \exp(x^n D_n (\dots (\exp(x^1 D_1(0, a, b))) \dots)) = (x, \Pi(x, a, b), \Omega(x, a, b)) \in \mathbb{K}^n \times \mathbb{K}^m \times \mathbb{K}^p,$$

for some two local analytic maps $\Pi = (\Pi^1, \dots, \Pi^m)$ and $\Omega = (\Omega^1, \dots, \Omega^p)$ and the next lemma is straightforward.

Lemma 2.4. *The general solution of (\mathcal{E}) is*

$$(2.5) \quad y(x) := \Pi(x, a, b),$$

where (a, b) varies in $\mathbb{K}^p \times \mathbb{K}^m$. Furthermore, for $q = 1, \dots, p$:

$$(2.6) \quad \Omega^q(x, a, b) \equiv \Pi_{x\beta(q)}^{j(q)}(x, a, b).$$

This leads to introducing a fundamental geometric object.

Definition 2.7. *The submanifold of solutions $\mathcal{V}_S(\mathcal{E})$ associated to (\mathcal{E}) is the analytic submanifold of $\mathbb{K}_x^n \times \mathbb{K}_a^p \times \mathbb{K}_b^m$ defined by the Cartesian equations:*

$$(2.8) \quad 0 = -y^j + \Pi^j(x, a, b), \quad j = 1, \dots, m.$$

There is a strong interplay between the study of (\mathcal{E}) and the geometry of $\mathcal{V}_S(\mathcal{E})$. By construction, the diffeomorphism:

$$(2.9) \quad \begin{cases} A : \mathbb{K}^{n+p+m} [\text{coordinates } (x^i, a^q, b^j)] \longrightarrow \mathbb{K}^{n+m+p} [\text{coordinates } (x^i, y^j, y_{\beta(q)}^{j(q)})] \\ A(x^i, a^q, b^j) := (x^i, \Pi^j(x, a, b), \Pi_{x\beta(q)}^{j(q)}(x, a, b)), \end{cases}$$

sends the *foliation* F_v by the variables x whose leaves are $\{a = \text{cst.}, b = \text{cst.}\}$ (see the diagram), to the previous foliation $F_{\Delta_{\mathcal{E}}}$.

2.10. PDE system associated to a submanifold. Inversely, let \mathcal{M} be a submanifold of $\mathbb{K}_x^n \times \mathbb{K}_y^m \times \mathbb{K}_a^p \times \mathbb{K}_b^m$ of the form

$$(2.11) \quad y^j = \Pi^j(x, a, b), \quad j = 1, \dots, m.$$

A necessary condition for it to be the complexification of a generic $M \subset \mathbb{C}^{n+m}$ is that $p = n$ (answer to an exercise above).

Definition 2.12. \mathcal{M} is *solvable with respect to the parameters* if $b \mapsto \Pi(0, 0, b)$ of rank m at $b = 0$ and if there exist $\kappa \geq 1$, multiindices $\beta(1), \dots, \beta(p) \in \mathbb{N}^n$ with $|\beta(q)| \geq 1$ for $q = 1, \dots, p$ and $\max_{1 \leq q \leq p} |\beta(q)| = \kappa$, together with integers $j(1), \dots, j(p)$ with $1 \leq j(q) \leq m$ such that the local \mathbb{K} -analytic map

$$(2.13) \quad \mathbb{K}^{m+p} \ni (a, b) \longmapsto \left((\Pi^j(0, a, b))^{1 \leq j \leq m}, \left(\Pi_{x\beta(q)}^{j(q)}(0, a, b) \right)_{1 \leq q \leq p} \right) \in \mathbb{K}^{m+p}$$

is of rank equal to $m + p$ at $(a, b) = (0, 0)$

When \mathcal{M} is the submanifold of solutions of a system (\mathcal{E}) , it is automatically solvable with respect to the variables, the pairs $(j(q), \beta(q))$ being the same as in the arguments of the right hand sides F_α^j in (\mathcal{E}) . Proceeding as in §1.16, we may associate to \mathcal{M} a system of the form (\mathcal{E}) . Since we need introduce some new notation, let us repeat the argument.

Considering $y = y(x) = \Pi(x, a, b)$ as a function of x with extra parameters (a, b) and applying $\partial^{|\alpha|}/\partial x^\alpha$, we get $y_{x^\alpha}^j(x) = \Pi_{x^\alpha}^j(x, a, b)$. Writing only the relevant $(m + p)$ equations:

$$(2.14) \quad \begin{cases} y^j(x) = \Pi^j(x, a, b), \\ y_{x^{\beta(q)}}^{j(q)} = \Pi_{x^{\beta(q)}}^{j(q)}(x, a, b), \end{cases}$$

the assumption of solvability with respect to parameters enables to get

$$(2.15) \quad \begin{cases} a^q = A^q(x^i, y^j, y_{\beta(q_1)}^{j(q_1)}), \\ b^j = B^j(x^i, y^{j_1}, y_{\beta(q)}^{j(q)}). \end{cases}$$

For every $(j, \alpha) \neq (j, 0)$ and $\neq (j(q), \beta(q))$, we then replace (a, b) in $y_{x^\alpha}^j = \Pi_{x^\alpha}^j$:

$$(2.16) \quad \begin{aligned} y_{x^\alpha}^j(x) &= \Pi_{x^\alpha}^j\left(x, A(x^i, y^{j_1}(x), y_{\beta(q)}^{j(q)}(x)), B(x^i, y^{j_1}(x), y_{\beta(q)}^{j(q)}(x))\right) \\ &=: F_\alpha^j(x^i, y^{j_1}(x), y_{\beta(q)}^{j(q)}(x)). \end{aligned}$$

Proposition 2.17. *There is a one-to-one correspondence*

$$(2.18) \quad (\mathcal{E}_{\mathcal{M}}) = (\mathcal{E}) \longleftrightarrow \mathcal{M} = \mathcal{M}_{(\mathcal{E})},$$

between completely integrable systems of partial differential equations of the general form (\mathcal{E}) and submanifolds (of solutions) \mathcal{M} of the form (2.11) which are solvable with respect to the parameters. Of course

$$(2.19) \quad (\mathcal{E}_{\mathcal{M}(\mathcal{E})}) = (\mathcal{E}) \quad \text{and} \quad \mathcal{M}_{(\mathcal{E}_{\mathcal{M}})} = \mathcal{M}.$$

2.20. Transfer of total differentiations. We notice that the auxiliary functions A^q and B^j enable to express the inverse of \mathbf{A} :

$$(2.21) \quad \mathbf{A}^{-1} : (x^{i_1}, y^{j_1}, y_{\beta(q_1)}^{j(q_1)}) \longmapsto \left(x^i, A^q(x^{i_1}, y^{j_1}, y_{\beta(q_1)}^{j(q_1)}), B^j(x^{i_1}, y^{j_1}, y_{\beta(q_1)}^{j(q_1)})\right).$$

More importantly, the total differentiation operator considerably simplifies when viewed on \mathcal{M} . This observation is useful for translating differential invariants of (\mathcal{E}) as differential invariants of \mathcal{M} .

Lemma 2.22. *Through \mathbf{A} , for $i = 1, \dots, n$, the pull-back of the total differentiation operator D_i is simply $\frac{\partial}{\partial x^i}$, or equivalently:*

$$(2.23) \quad \mathbf{A}_* \left(\frac{\partial}{\partial x^i} \right) = D_i.$$

Proof. Let $\ell = \ell(x^i, y^j, y_{\beta(q)}^{j(q)})$ be any function defined on $\Delta_{\mathcal{E}}$. Composing with \mathbf{A} yields the function $\Lambda := \ell \circ \mathbf{A}$, i.e.

$$(2.24) \quad \Lambda(x, a, b) \equiv \ell\left(x^i, \Pi^j(x, a, b), \Pi_{x^{\beta(q)}}^{j(q)}(x, a, b)\right).$$

Differentiating with respect to x^i , we get, dropping the arguments:

$$(2.25) \quad \frac{\partial \Lambda}{\partial x^i} = \frac{\partial \ell}{\partial x^i} + \sum_{j=1}^m \Pi_{x^i}^j \frac{\partial \ell}{\partial y^j} + \sum_{q=1}^p \Pi_{x^i x^{\beta(q)}}^{j(q)} \frac{\partial \ell}{\partial y_{x^{\beta(q)}}^{j(q)}}.$$

Replacing the appearing $\Pi_{x^\alpha}^j$ for which $(j, \alpha) \neq (j, 0)$ and $\neq (j(q), \beta(q))$ by F_α^j , we recover D_i as defined by (1.38), whence $\frac{\partial \Lambda}{\partial x^i} = D_i \ell$. \square

2.26. Transfer of algebrico-differential expressions. The diffeomorphism A may be used to translate algebrico-differential expressions from \mathcal{M} to (\mathcal{E}) and vice-versa:

$$(2.27) \quad \mathfrak{l}_{\mathcal{M}}(J_{x,a,b}^{\lambda+\kappa+1} \Pi) \longleftrightarrow \mathfrak{l}_{(\mathcal{E})}(J_{x,y,y_1}^\lambda F).$$

Here, $\lambda \in \mathbb{N}$, the letter J is used to denote jets, and $\mathfrak{l} = \mathfrak{l}_{\mathcal{M}}$ or $= \mathfrak{l}_{(\mathcal{E})}$ is a polynomial or more generally, a quotient of polynomials with respect to its jet arguments. Notice the shift by $\kappa + 1$ of the jet orders.

Example 2.28. Suppose $n = m = 1$ and $\kappa = 1$. Then $F = \Pi_{xx}$. As an exercise, let us compute F_x, F_y, F_{y_1} in terms of $J_{x,a,b}^3 \Pi$. We start with the identity

$$(2.29) \quad F(x, y, y_1) \equiv \Pi_{xx}(x, A(x, y, y_1), B(x, y, y_1)),$$

that we differentiate with respect to x , to y and to y_1 :

$$(2.30) \quad \begin{aligned} F_x &= \Pi_{xxx} + \Pi_{xxa} A_x + \Pi_{xxb} B_x, \\ F_y &= \Pi_{xxa} A_y + \Pi_{xxb} B_y, \\ F_{y_1} &= \Pi_{xxa} A_{y_1} + \Pi_{xxb} B_{y_1}. \end{aligned}$$

Thus, we need to compute $A_x, A_y, A_{y_1}, B_x, B_y, B_{y_1}$. This is easy: it suffices to differentiate the two identities that define A and B as implicit functions, namely:

$$(2.31) \quad \begin{aligned} y &\equiv \Pi(x, A(x, y, y_1), B(x, y, y_1)) \quad \text{and} \\ y_1 &\equiv \Pi_x(x, A(x, y, y_1), B(x, y, y_1)) \end{aligned}$$

with respect to x , to y and to y_1 , which gives six new identities:

$$(2.32) \quad \begin{aligned} 0 &= \Pi_x + \Pi_a A_x + \Pi_b B_x, & 0 &= \Pi_{xx} + \Pi_{xa} A_x + \Pi_{xb} B_x, \\ 1 &= \Pi_a A_y + \Pi_b B_y, & 0 &= \Pi_{xa} A_y + \Pi_{xb} B_y \\ 0 &= \Pi_a A_{y_1} + \Pi_b B_{y_1}, & 1 &= \Pi_{xa} A_{y_1} + \Pi_{xb} B_{y_1}, \end{aligned}$$

and to solve each of the three linear systems of two equations located in a line, noticing that their common determinant $\Pi_b \Pi_{xa} - \Pi_a \Pi_{xb}$ does not vanish at the origin, since $\Pi = b + xa + O_3$. By elementary Cramer formulas, we get:

$$(2.33) \quad \left\{ \begin{aligned} A_x &= \frac{-\Pi_b \Pi_{xx} + \Pi_x \Pi_{xb}}{\Pi_b \Pi_{xa} - \Pi_a \Pi_{xb}}, & B_x &= \frac{-\Pi_x \Pi_{xa} + \Pi_a \Pi_{xx}}{\Pi_b \Pi_{xa} - \Pi_a \Pi_{xb}}, \\ A_y &= \frac{-\Pi_{xb}}{\Pi_b \Pi_{xa} - \Pi_a \Pi_{xb}}, & B_y &= \frac{\Pi_{xa}}{\Pi_b \Pi_{xa} - \Pi_a \Pi_{xb}}, \\ A_{y_1} &= \frac{\Pi_b}{\Pi_b \Pi_{xa} - \Pi_a \Pi_{xb}}, & B_{y_1} &= \frac{-\Pi_a}{\Pi_b \Pi_{xa} - \Pi_a \Pi_{xb}}. \end{aligned} \right.$$

Replacing in (2.30), no simplification occurs and we get what we wanted:

$$(2.34) \quad \left\{ \begin{aligned} F_x &= \Pi_{xxx} + \frac{\Pi_{xxa} [-\Pi_b \Pi_{xx} + \Pi_x \Pi_{xb}] + \Pi_{xxb} [-\Pi_x \Pi_{xa} + \Pi_a \Pi_{xx}]}{\Pi_b \Pi_{xa} - \Pi_a \Pi_{xb}}, \\ F_y &= \frac{-\Pi_{xxa} \Pi_{xb} + \Pi_{xxb} \Pi_{xa}}{\Pi_b \Pi_{xa} - \Pi_a \Pi_{xb}}, \\ F_{y_1} &= \frac{\Pi_{xxa} \Pi_b - \Pi_{xxb} \Pi_a}{\Pi_b \Pi_{xa} - \Pi_a \Pi_{xb}}, \end{aligned} \right.$$

One sees $D F = F_x + \Pi_x F_y + \Pi_{xx} F_{y_1} = \Pi_{xxx}$ simply, as predicted by Lemma 2.22.

Second order derivatives $F_{xx}, F_{xy}, F_{xy_1}, F_{yy}, F_{yy_1}, F_{y_1y_1}$ have still reasonable complexity, when expressed in terms of $J_{x,a,b}^4 \Pi$. Beyond, the computations explode.

Open question 2.35. *A second order ordinary differential equation $y_{xx} = F(x, y, y_x)$ has two fundamental differential invariants, namely ([Tr1896, Ca1924, GTW1989, Ol1995]):*

$$(2.36) \quad \begin{aligned} I_{(\mathcal{E}_1)}^1 &:= \frac{\partial^4 F}{\partial y_1^4} \quad \text{and} \\ I_{(\mathcal{E}_1)}^2 &:= DD(F_{y_1y_1}) - F_{y_1} D(F_{y_1y_1}) - 4D(F_{yy_1}) + 6F_{yy} - 3F_y F_{y_1y_1} + 4F_{y_1} F_{yy_1}. \end{aligned}$$

Compute $I_{\mathcal{M}_1}^1$ and $I_{\mathcal{M}_1}^2$.

Although the notion of diffeomorphism is clear and apparently obvious from the intuitive, geometric and conceptual viewpoints, in concrete applications and in explicit computations, it almost never straightforward to transfer algebrico-differential objects.

Open problem 2.37. *For general (\mathcal{E}) and \mathcal{M} , build closed combinatorial formulas executing the double translation (2.27).*

2.38. Plan for the sequel. We will endeavour a general theory showing that the study of systems (\mathcal{E}) and the study of submanifolds of solutions \mathcal{M} gives complementary views on the same object. In fact, Lie symmetries, equivalence problems, Cartan connections, normal forms and classification lists may be endeavoured on both sides, yielding essentially equivalent results, though the translation is seldom straightforward. In Section 3, 4 and 5, we review some features from the side (\mathcal{E}) , before studying some aspects from the side of \mathcal{M} . A more systematic and complete approach shall appear as a monography.

§3. CLASSIFICATION PROBLEMS

3.1. Transformations of PDE systems. Through a local \mathbb{K} -analytic change of variables close to the identity $(x, y) \mapsto \varphi(x, y) =: (x', y')$, the system (\mathcal{E}) transforms to a similar system, with primes:

$$(\mathcal{E}') \quad y'^j_{x'^\alpha}(x') = F'^j_\alpha(x', y'(x'), (y'^{j(q)}_{x'^{\beta(q)}}(x'))_{1 \leq q \leq p}).$$

Example 3.2. Coming back temporarily to the notations of §1.12(II), with $n = m = \kappa = 1$, assume that $y_{xx} = f(x, y, y_x)$ transforms to $Y_{XX} = F(X, Y, Y_X)$ through a local diffeomorphism $(x, y) \mapsto (X, Y) = (X(x, y), Y(x, y))$. How F is related to f ? By symmetry, it suffices to compute f in terms of F, X, Y . The prolongation to $\mathcal{J}_{1,1}^2$ of the diffeomorphism has components ([BK1989, Me2004]):

$$(3.3) \quad Y_X = \frac{Y_x + y_x Y_y}{X_x + y_x X_y},$$

and

$$(3.4) \quad \begin{aligned} Y_{XX} = \frac{1}{[X_x + y_x X_y]^3} & \left(y_{xx} \cdot \left| \begin{array}{cc} X_x & X_y \\ Y_x & Y_y \end{array} \right| + \left| \begin{array}{cc} X_x & X_{xx} \\ Y_x & Y_{xx} \end{array} \right| + \right. \\ & + y_x \cdot \left\{ 2 \left| \begin{array}{cc} X_x & X_{xy} \\ Y_x & Y_{xy} \end{array} \right| - \left| \begin{array}{cc} X_{xx} & X_y \\ Y_{xx} & Y_y \end{array} \right| \right\} + \\ & + y_x y_x \cdot \left\{ \left| \begin{array}{cc} X_x & X_{yy} \\ Y_x & Y_{yy} \end{array} \right| - 2 \left| \begin{array}{cc} X_{xy} & X_y \\ Y_{xy} & Y_y \end{array} \right| \right\} + \\ & \left. + y_x y_x y_x \cdot \left\{ - \left| \begin{array}{cc} X_{yy} & X_y \\ Y_{yy} & Y_y \end{array} \right| \right\} \right). \end{aligned}$$

It then suffices to replace Y_{XX} above by $F(X, Y, Y_X)$ and to solve y_{xx} :

$$\begin{aligned}
 (3.5) \quad y_{xx} &= \frac{1}{\begin{vmatrix} X_x & X_y \\ Y_x & Y_y \end{vmatrix}} \left([X_x + y_x X_y]^3 F \left(X, Y, \frac{Y_x + y_x Y_y}{X_x + y_x X_y} \right) - \begin{vmatrix} X_x & X_{xx} \\ Y_x & Y_{xx} \end{vmatrix} + \right. \\
 &\quad \left. + y_x \cdot \left\{ -2 \begin{vmatrix} X_x & X_{xy} \\ Y_x & Y_{xy} \end{vmatrix} + \begin{vmatrix} X_{xx} & X_y \\ Y_{xx} & Y_y \end{vmatrix} \right\} + \right. \\
 &\quad \left. + y_x y_x \cdot \left\{ - \begin{vmatrix} X_x & X_{yy} \\ Y_x & Y_{yy} \end{vmatrix} + 2 \begin{vmatrix} X_{xy} & X_y \\ Y_{xy} & Y_y \end{vmatrix} \right\} + \right. \\
 &\quad \left. + y_x y_x y_x \cdot \left\{ \begin{vmatrix} X_{yy} & X_y \\ Y_{yy} & Y_y \end{vmatrix} \right\} \right) \\
 &=: f(x, y, y_x).
 \end{aligned}$$

Open problem 3.6. Find general formulas expressing the F_α^j in terms of F_α^{ij} , x^i , y^j .

Conversely, given two such systems (\mathcal{E}) and (\mathcal{E}') , when do they transform to each other? Let $\pi'_{\kappa,p}$ denote the projection from $\mathcal{J}'_{n,m}{}^{\kappa+1}$ to $\Delta_{\mathcal{E}'}$ defined by

$$(3.7) \quad \pi'_{\kappa,p}(x'^i, y'^j, y'^j_{i_1}, \dots, y'^j_{i_1, \dots, i_{\kappa+1}}) := (x'^i, y'^j, y'^j_{\beta(q)})$$

Let $\varphi^{(\kappa+1)}$ be the $(\kappa+1)$ -th prolongation of φ (Section 1(II)).

Lemma 3.8. ([O11986, BK1989, O11995]) *The following three conditions are equivalent:*

- (1) φ transforms (\mathcal{E}) to (\mathcal{E}') ;
- (2) its $(\kappa+1)$ -th prolongation $\varphi^{(\kappa+1)} : \mathcal{J}_{n,m}^{\kappa+1} \rightarrow \mathcal{J}'_{n,m}{}^{\kappa+1}$ maps $\Delta_{\mathcal{E}}$ to $\Delta_{\mathcal{E}'}$;
- (3) $\varphi^{(\kappa+1)} : \mathcal{J}_{n,m}^{\kappa+1} \rightarrow \mathcal{J}'_{n,m}{}^{\kappa+1}$ maps $\Delta_{\mathcal{E}}$ to $\Delta_{\mathcal{E}'}$ and the associated map

$$(3.9) \quad \Phi_{\mathcal{E},\mathcal{E}'} := \pi'_{\kappa,p} \circ (\varphi^{(\kappa+1)}|_{\Delta_{\mathcal{E}}})$$

sends every leaf of $F_{\Delta_{\mathcal{E}}}$ to some leaf of $F_{\Delta_{\mathcal{E}'}}$.

Equivalence problem 3.10. Find an algorithm to decide whether two given (\mathcal{E}) and (\mathcal{E}') are equivalent.

Élie Cartan's widely applicable method (not reviewed here; [Ca1937, Ste1983, G1989, HK1989, Fe1995, O11995]) provides an answer "in principle" to this question by reducing to an $\{e\}$ -structure an initial G -structure associated to (\mathcal{E}) . Due to the incredible size-length-complexity of the underlying computations, this approach almost never abutes: it is forced to incompleteness. But in fact, the main question is to classify.

Classification problem 3.11. Classify systems (\mathcal{E}) , namely provide complete lists of all possible such equations written in simplified "normal", easily recognizable forms.

Both problems are deeply linked to the classification of Lie algebras of local vector fields. For $n = 1$, $m = 1$ and $\kappa = 1$, namely (\mathcal{E}_1) : $y_{xx} = F(x, y, y_x)$, Lie and Tresse solved the two problems³. Table 7 of [O11986], below reproduced, describes the results.

³ The author knows no complete confirmation of the Lie-Tresse classification by means of É. Cartan's method of equivalence.

	Symmetry group	Dimension	Invariant equation
(1)		0	$y_{xx} = F(x, y, y_x)$
(2)	∂_y	1	$y_{xx} = F(x, y_x)$
(3)	∂_x, ∂_y	2	$y_{xx} = F(y_x)$
(4)	$\partial_x, e^x \partial_y$	2	$y_{xx} - y_x = F(y_x - y)$
(5)	$\partial_x, \partial_x - y \partial_y, x^2 \partial_x - 2xy \partial_y$	3	$y_{xx} = \frac{3y_x^2}{2y} + cy^3$
(6)	$\partial_x, x \partial_x - y \partial_y,$ $x^2 \partial_x - (2xy + 1) \partial_y$	3	$y_{xx} = 6yy_x - 4y^3 +$ $+c(y_x - y^2)^{3/2}$
(7)	$\partial_x, \partial_y, x \partial_x + \alpha y \partial_y,$ $\alpha \neq 0, \frac{1}{2}, 1, 2$	3	$y_{xx} = c(y_x)^{\frac{\alpha-2}{\alpha-1}}$
(8)	$\partial_x, \partial_y, x \partial_x + (x + y) \partial_y$	3	$y_{xx} = ce^{-y_x}$
(9)	$\partial_x, \partial_y, y \partial_x, x \partial_y, y \partial_y,$ $x^2 \partial_x + xy \partial_y, xy \partial_x + y^2 \partial_y$	8	$y_{xx} = 0$

Table 1.

However, the author knows no modern reference offering a complete proof of this classification, with precise insight on the assumptions (some normal forms hold true only at a generic point). In addition, the above Lie-Tresse list is still slightly incomplete in the sense that it does not precise which are the conditions satisfied by F (Table 7 in [O11986]) insuring in the first four lines that $\mathfrak{S}\mathfrak{Y}\mathfrak{M}(\mathcal{E}_1)$ is indeed of small dimension 0, 1 or 2.

Open question 3.12. *Specify some precise nondegeneracy conditions upon F in the first four lines of Table 1.*

§4. PUNCTUAL AND INFINITESIMAL LIE SYMMETRIES

4.1. Lie symmetries of (\mathcal{E}) . Let $\varphi = (\phi, \psi)$ be a diffeomorphism of $\mathbb{K}_x^n \times \mathbb{K}_y^n$ as in (1.7)(II).

Definition 4.2. ([O11986, O11995, BK1989]) φ is a (local) Lie symmetry of (\mathcal{E}) if it transforms the graph of every solution of (\mathcal{E}) into the graph of another solution.

To explain, we must pass to jet spaces. Denote the components of the $(\kappa + 1)$ -th prolongation $\varphi^{(\kappa+1)} : \mathcal{J}_{n,m}^{\kappa+1} \rightarrow \mathcal{J}_{n,m}^{\kappa+1}$ by

$$(4.3) \quad \varphi^{(\kappa+1)} = \left(\phi^{i_1}, \psi^{j_1}, \Phi_{i_1}^j, \Phi_{i_1, i_2}^j, \dots, \Phi_{i_1, i_2, \dots, i_{\kappa+1}}^j \right).$$

The restriction $\varphi^{(\kappa+1)}|_{\Delta_{\mathcal{E}}}$ is obtained by replacing each jet variable y_{α}^j by F_{α}^j , whenever $(j, \alpha) \neq (j, 0)$ and $\neq (j(q), \beta(q))$, and wherever it appears⁴ in the $\Phi_{i_1, \dots, i_{\lambda}}^j$.

Let $\pi_{\kappa,p}$ denote the projection from $\mathcal{J}_{n,m}^{\kappa+1}$ to $\Delta_{\mathcal{E}} \simeq \mathbb{K}^{m+n+p}$ defined by

$$(4.4) \quad \pi_{\kappa,p}(x^i, y^j, y_{i_1}^j, \dots, y_{i_1, \dots, i_{\kappa+1}}^j) := \left(x^i, y^j, y_{\beta(q)}^j \right),$$

and introduce the map

$$(4.5) \quad \varphi_{\Delta_{\mathcal{E}}} := \pi_{\kappa,p} \circ \left(\varphi^{(\kappa+1)}|_{\Delta_{\mathcal{E}}} \right) \equiv \left(\varphi(x^i, y^j), \Phi_{\beta(q)}^{j(q)}(x^i, y^j, y_{\beta(q_1)}^j) \right).$$

Lemma 4.6. ([O11986, O11995, BK1989], [*]) *The following three conditions are equivalent:*

⁴ Remind from Section 1(II) that we have not (open problem) provided a complete explicit expression of $\Phi_{i_1, \dots, i_{\lambda}}^j$ for general $n \geq 1, m \geq 1$ and $\lambda \geq 1$.

- (1) the diffeomorphism φ is a Lie symmetry of (\mathcal{E}) ;
- (2) $\varphi^{(\kappa+1)}|_{\Delta_{\mathcal{E}}}$ sends $\Delta_{\mathcal{E}}$ to $\Delta_{\mathcal{E}}$;
- (3) $\varphi^{(\kappa+1)}|_{\Delta_{\mathcal{E}}}$ sends $\Delta_{\mathcal{E}}$ to $\Delta_{\mathcal{E}}$ and $\varphi_{\Delta_{\mathcal{E}}} = \pi_{\kappa,p}(\varphi^{(\kappa+1)}|_{\Delta_{\mathcal{E}}})$ is a symmetry of the foliation $F_{\Delta_{\mathcal{E}}}$, namely it sends every leaf to some other leaf.

Then the set of Lie symmetries of (\mathcal{E}) constitutes a local Lie (pseudo)group.

4.7. Infinitesimal Lie symmetries of (\mathcal{E}) . Let

$$(4.8) \quad \mathcal{L} = \sum_{i=1}^n \mathcal{X}^i(x, y) \frac{\partial}{\partial x^i} + \sum_{j=1}^m \mathcal{Y}^j(x, y) \frac{\partial}{\partial y^j},$$

be a (local) vector field on \mathbb{K}^{n+m} having analytic coefficients. Denote its flow by $\varphi_t(x, y) := \exp(t\mathcal{L})(x, y)$, $t \in \mathbb{K}$. As in Section 1(II), by differentiating the prolongation $(\varphi_t)^{(\kappa+1)}$ with respect to t at $t = 0$, we get the prolonged vector field $\mathcal{L}^{(\kappa+1)}$ on $\mathcal{J}_{n,m}^{\kappa+1}$, having the general form (Part II):

$$(4.9) \quad \mathcal{L}^{(\kappa+1)} = \mathcal{L} + \sum_{j=1}^m \sum_{i_1=1}^n \mathbf{Y}_{i_1}^j \frac{\partial}{\partial y_{i_1}^j} + \cdots + \sum_{j=1}^m \sum_{i_1, \dots, i_{\kappa+1}=1}^n \mathbf{Y}_{i_1, \dots, i_{\kappa+1}}^j \frac{\partial}{\partial y_{i_1, \dots, i_{\kappa+1}}^j},$$

with known explicit expressions for the $\mathbf{Y}_{i_1, \dots, i_{\lambda}}^j$.

Definition 4.10. \mathcal{L} is an *infinitesimal symmetry* of (\mathcal{E}) if for every small t , its time- t flow map φ_t is a Lie symmetry of (\mathcal{E}) .

The restriction $\mathcal{L}^{(\kappa+1)}|_{\Delta_{\mathcal{E}}}$ is obtained by replacing every y_{α}^j by F_{α}^j in all coefficients $\mathbf{Y}_{i_1}^j, \dots, \mathbf{Y}_{i_1, \dots, i_{\kappa+1}}^j$. Then the coefficients become functions of $(x^{i_1}, y^{j_1}, y_{\beta(q_1)}^{j(q_1)})$ only.

Lemma 4.11. ([O1986, O1995, BK1989], [*]) *The following three conditions are equivalent:*

- (1) the vector field \mathcal{L} is an infinitesimal Lie symmetry of (\mathcal{E}) ;
- (2) its $(\kappa + 1)$ -th prolongation $\mathcal{L}^{(\kappa+1)}$ is tangent to the skeleton $\Delta_{\mathcal{E}}$;
- (3) $\mathcal{L}^{(\kappa+1)}$ is tangent to $\Delta_{\mathcal{E}}$ and the push-forward

$$(4.12) \quad \mathcal{L}_{\Delta_{\mathcal{E}}} := (\pi_{\kappa,p})_*(\mathcal{L}^{(\kappa+1)}|_{\Delta_{\mathcal{E}}})$$

is an infinitesimal symmetry of the foliation $F_{\Delta_{\mathcal{E}}}$, namely for every $i = 1, \dots, n$, the Lie bracket $[\mathcal{L}_{\Delta_{\mathcal{E}}}, D_i]$ is a linear combination of $\{D_1, \dots, D_n\}$.

According to [O1986, BK1989, O1995], the set of infinitesimal Lie symmetries constitutes a Lie algebra, with the property $[\mathcal{L}^{(\kappa+1)}, \mathcal{L}'^{(\kappa+1)}] = [\mathcal{L}, \mathcal{L}']^{(\kappa+1)}$. We summarize by a diagram.

$$\begin{array}{ccc}
 \mathcal{J}_{n,m}^{\kappa+1} & \xrightarrow{\varphi^{(\kappa+1)}} & \mathcal{J}_{n,m}^{\kappa+1} & & \mathcal{L}^{(\kappa+1)} \\
 \downarrow \pi_{\kappa,p} & & \downarrow \pi_{\kappa,p} & & \downarrow \\
 \Delta_{\mathcal{E}} & \xrightarrow{\varphi_{\Delta_{\mathcal{E}}}} & \Delta_{\mathcal{E}} & & \mathcal{L}_{\Delta_{\mathcal{E}}} \\
 \downarrow \pi_{\kappa} & & \downarrow \pi_p & & \downarrow \\
 \mathbb{K}_x^n \times \mathbb{K}_y^m & \xrightarrow{\varphi} & \mathbb{K}_x^n \times \mathbb{K}_y^m & & \mathcal{L}
 \end{array}$$

4.13. Sophus Lie's algorithm. We describe the general process. Its complexity will be exemplified in Section 5 (to be read simultaneously).

The tangency of $\mathcal{L}^{(\kappa+1)}$ to $\Delta_{\mathcal{E}}$ is expressed by applying $\mathcal{L}^{(\kappa+1)}$ to the equations $0 = -y_{\alpha}^j + F_{\alpha}^j$, which yields:

$$(4.14) \quad 0 = -\mathbf{Y}_{\alpha}^j + \sum_{i=1}^n \mathcal{X}^i \frac{\partial F_{\alpha}^j}{\partial x^i} + \sum_{l=1}^n \mathcal{Y}^l \frac{\partial F_{\alpha}^j}{\partial y^l} + \sum_{q=1}^p \mathbf{Y}_{\beta(q)}^{j(q)} \frac{\partial F_{\alpha}^j}{\partial y_{\beta(q)}^j},$$

for $(j, \alpha) \neq (j, 0)$ and $\neq (j(q), \beta(q))$. Restricting a coefficient $\mathbf{Y}_{i_1, \dots, i_{\lambda}}^j$ to $\Delta_{\mathcal{E}}$, namely replacing everywhere in it each y_{α}^j by F_{α}^j , provides a specialized coefficient

$$(4.15) \quad \widehat{\mathbf{Y}}_{i_1, \dots, i_{\lambda}}^j = \widehat{\mathbf{Y}}_{i_1, \dots, i_{\lambda}}^j(x^{i_1}, y^{j_1}, y_{\beta(q_1)}^{j(q_1)}, J_{x,y}^{\lambda} \mathcal{X}^{i_1}, J_{x,y}^{\lambda} \mathcal{Y}^{j_1}),$$

that depends **linearly** on the λ -th jet of the coefficients of \mathcal{L} , as confirmed by an inspection of Part II's formulas. Here, we use the jet notation $J_{x,y}^{\lambda} \mathcal{Z} := (\partial_x^{\alpha_1} \partial_y^{\beta_1} \mathcal{Z})_{|\alpha_1|+|\beta_1| \leq \lambda}$. We thus get equations

$$(4.16) \quad 0 \equiv -\widehat{\mathbf{Y}}_{\alpha}^j + \sum_{i=1}^n \mathcal{X}^i \frac{\partial F_{\alpha}^j}{\partial x^i} + \sum_{l=1}^n \mathcal{Y}^l \frac{\partial F_{\alpha}^j}{\partial y^l} + \sum_{q=1}^p \widehat{\mathbf{Y}}_{\beta(q)}^{j(q)} \frac{\partial F_{\alpha}^j}{\partial y_{\beta(q)}^j},$$

involving only the variables $(x^{i_1}, y^{j_1}, y_{\beta(q_1)}^{j(q_1)})$.

Next, we develop every such equation with respect to the powers of $y_{\beta(q_1)}^{j(q_1)}$:

$$(4.17) \quad 0 \equiv \sum_{\mu_1, \dots, \mu_p \geq 0} (y_{\beta(1)}^{j(1)})^{\mu_1} \dots (y_{\beta(p)}^{j(p)})^{\mu_p} \Psi_{\alpha, \mu_1, \dots, \mu_p}^j(x^{i_1}, y^{j_1}, J_{x,y}^{\kappa+1} \mathcal{X}^{i_1}, J_{x,y}^{\kappa+1} \mathcal{Y}^{j_1}).$$

The $\Psi_{\alpha, \mu_1, \dots, \mu_p}^j$ are **linear** with respect to $(J_{x,y}^{\kappa+1} \mathcal{X}^{i_1}, J_{x,y}^{\kappa+1} \mathcal{Y}^{j_1})$, with certain coefficients analytic with respect to (x, y) , which depend intrinsically (but in a complex manner) on the right hand sides F_{α}^j .

Proposition 4.18. *The vector field \mathcal{L} is an infinitesimal Lie symmetry of (\mathcal{E}) if and only if its coefficients $\mathcal{X}^{i_1}, \mathcal{Y}^{j_1}$ satisfy the linear PDE system:*

$$(4.19) \quad 0 = \Psi_{\alpha, \mu_1, \dots, \mu_p}^j(x^{i_1}, y^{j_1}, J_{x,y}^{\kappa+1} \mathcal{X}^{i_1}, J_{x,y}^{\kappa+1} \mathcal{Y}^{j_1})$$

for all $(j, \alpha) \neq (j, 0)$ and $\neq (j(q), \beta(q))$ and for all $(\mu_1, \dots, \mu_p) \in \mathbb{N}^p$.

In all known instances, a finite number of these equations suffices.

Example 4.20. With $n = m = \kappa = 1$, a second prolongation $\mathcal{L}^{(2)} = \mathcal{X} \frac{\partial}{\partial x} + \mathcal{Y} \frac{\partial}{\partial y} + \mathbf{Y}_1 \frac{\partial}{\partial y_1} + \mathbf{Y}_2 \frac{\partial}{\partial y_2}$ is tangent to the skeleton $0 = -y_2 + F(x, y, y_1)$ of (\mathcal{E}_1) if and only if $0 = -\mathbf{Y}_2 + \mathcal{X} F_x + \mathcal{Y} F_y + \mathbf{Y}_1 F_{y_1}$, or, developing:

$$(4.21) \quad \left\{ \begin{array}{l} 0 = -\mathcal{Y}_{xx} + [-2\mathcal{Y}_{xy} + \mathcal{X}_{xx}] y_1 + [-\mathcal{Y}_{yy} + 2\mathcal{X}_{xy}] (y_1)^2 + [\mathcal{X}_{yy}] (y_1)^3 + \\ \quad + [-\mathcal{Y}_y + 2\mathcal{X}_x] F + [3\mathcal{X}_y] y_1 F + [\mathcal{X}] F_x + [\mathcal{Y}] F_y + \\ \quad + [\mathcal{Y}_x] F_{y_1} + [\mathcal{Y}_y - \mathcal{X}_x] y_1 F_{y_1} + [-\mathcal{X}_y] (y_1)^2 F_{y_1}. \end{array} \right.$$

Developing $F = \sum_{k \geq 0} (y_1)^k F_k(x, y)$, we may obtain equations (4.19).

§5. EXAMPLES

5.1. Second order ordinary differential equation. Pursuing the study of (\mathcal{E}_1) , according to Section 7 below, we may assume that $F = O(y_x)$, or equivalently $F(x, y, 0) \equiv 0$.

Convention 5.2. The letters R will denote various functions of (x, y, y_1) , changing with the context. Similarly, $r = r(x, y)$, excluding the pure jet variable y_1 . Hence, symbolically:

$$(5.3) \quad R = r + y_1 r + (y_1)^2 r + (y_1)^3 r + \dots$$

So the skeleton is

$$(5.4) \quad y_2 = F(x, y, y_1) = y_1 R = y_1 r + (y_1)^2 r + (y_1)^3 r + \dots$$

Applying $\mathcal{L}^{(2)}$, see (2.3)(II) for its expression, we get:

$$(5.5) \quad 0 = -Y_2 + \mathcal{X} F_x + \mathcal{Y} F_y + Y_1 F_{y_1}.$$

Observe that $F_x = (y_1 R)_x = r y_1 + r (y_1)^2 + \dots$ and similarly for F_y , but that $(y_1 R)_{y_1} = r + r y_1 + r (y_1)^2 + \dots$. Inserting above Y_1, Y_2 given by (2.6)(II), replacing y_2 by $y_1 R$ and computing mod $(y_1)^4$, we get:

$$(5.6) \quad \begin{aligned} 0 \equiv & -\mathcal{Y}_{xx} + [-2\mathcal{Y}_{xy} + \mathcal{X}_{xx}] y_1 + [-\mathcal{Y}_{yy} + 2\mathcal{X}_{xy}] (y_1)^2 + [\mathcal{X}_{yy}] (y_1)^3 + \\ & + [-\mathcal{Y}_y + 2\mathcal{X}_x] (y_1 r + (y_1)^2 r + (y_1)^3 r) + [3\mathcal{X}_y] ((y_1)^2 r + (y_1)^3 r) + \\ & + [\mathcal{X}] (y_1 r + (y_1)^2 r + (y_1)^3 r) + [\mathcal{Y}] (y_1 r + (y_1)^2 r + (y_1)^3 r) + \\ & + [\mathcal{Y}_x] (r + y_1 r + (y_1)^2 r + (y_1)^3 r) + \\ & + [\mathcal{Y}_y - \mathcal{X}_x] (y_1 r + (y_1)^2 r + (y_1)^3 r) + [-\mathcal{X}_y] ((y_1)^2 r + (y_1)^3 r). \end{aligned}$$

We gather the powers $\text{cst.}, y_1, (y_1)^2$ and $(y_1)^3$, equating their coefficients to 0:

$$(5.7) \quad \begin{aligned} 0 &= -\mathcal{Y}_{xx} + P(\mathcal{Y}_x), \\ 0 &= -2\mathcal{Y}_{xy} + \mathcal{X}_{xx} + P(\mathcal{Y}_y, \mathcal{X}_x, \mathcal{X}, \mathcal{Y}, \mathcal{Y}_x), \\ 0 &= -\mathcal{Y}_{yy} + 2\mathcal{X}_{xy} + P(\mathcal{Y}_y, \mathcal{X}_x, \mathcal{X}_y, \mathcal{X}, \mathcal{Y}, \mathcal{Y}_x), \\ 0 &= \mathcal{X}_{yy} + P(\mathcal{Y}_y, \mathcal{X}_x, \mathcal{X}_y, \mathcal{X}, \mathcal{Y}, \mathcal{Y}_x) \end{aligned}$$

Convention 5.8. The letter P will denote various *linear* combinations of some precise partial derivatives of \mathcal{X}, \mathcal{Y} which have analytic coefficients in (x, y) .

By cross-differentiations and substitutions in the above system, all third, fourth, fifth, etc. order derivatives of \mathcal{X}, \mathcal{Y} may be expressed as $P(\mathcal{X}, \mathcal{Y}, \mathcal{X}_x, \mathcal{X}_y, \mathcal{Y}_x, \mathcal{Y}_y, \mathcal{Y}_{xy}, \mathcal{Y}_{yy})$.

Proposition 5.9. *An infinitesimal Lie symmetry $\mathcal{X} \frac{\partial}{\partial x} + \mathcal{Y} \frac{\partial}{\partial y}$ of (\mathcal{E}_1) is uniquely determined by the eight initial Taylor coefficients:*

$$(5.10) \quad \mathcal{X}(0), \mathcal{Y}(0), \mathcal{X}_x(0), \mathcal{X}_y(0), \mathcal{Y}_x(0), \mathcal{Y}_y(0), \mathcal{Y}_{xy}(0), \mathcal{Y}_{yy}(0).$$

The bound $\dim \mathfrak{Sym}(\mathcal{E}_1) \leq 8$ is attained with $F = 0$, whence all $P = 0$ and

$$(5.11) \quad \begin{cases} A := \partial_y, & E := y \partial_y, \\ B := \partial_x, & F := y \partial_x, \\ C := x \partial_y, & G := xx \partial_x + xy \partial_y, \\ D := x \partial_x, & H := xy \partial_x + yy \partial_y. \end{cases}$$

are infinitesimal generators of the group $\mathrm{PGL}_3(\mathbb{K}) = \mathrm{Aut}(P_2(\mathbb{K}))$ of projective transformations

$$(5.12) \quad (x, y) \mapsto \left(\frac{\alpha x + \beta y + \gamma}{\lambda x + \mu y + \nu}, \frac{\delta x + \eta y + \epsilon}{\lambda x + \mu y + \nu} \right)$$

stabilizing the collections of all affine lines of \mathbb{K}^2 , namely the solutions of the model equation $y_{xx} = 0$. The model Lie algebra $\mathfrak{pgl}_3(\mathbb{K}) \simeq \mathfrak{sl}_3(\mathbb{K})$ is simple.

Theorem 5.13. *The bound $\dim \mathfrak{G}\mathfrak{M}(\mathcal{E}_1) \leq 8$ is attained if and only if (\mathcal{E}_1) is equivalent, through a diffeomorphism $(x, y) \mapsto (X, Y)$, to $Y_{XX} = 0$.*

Proof. The statement is well known ([Lie1883, EL1890, Tr1896, Se1931, Ca1932a, Ol1986, HK1989, Ib1992, Ol1995, Sh1997, Su2001, N2003, Me2004]). We provide a (new?) proof which has the advantage to enjoy direct generalizations to all PDE systems whose model Lie algebras are semisimple, for instance (\mathcal{E}_2) , (\mathcal{E}_3) and (\mathcal{E}_5) .

The Lie brackets between the eight generators (5.11) are:

	A	B	C	D	E	F	G	H
A	0	0	0	0	A	B	C	$D + 2E$
B	0	0	A	B	0	0	$E + 2D$	F
C	0	$-A$	0	$-C$	C	$D - E$	0	G
D	0	$-B$	C	0	0	$-F$	G	0
E	$-A$	0	$-C$	0	0	F	0	H
F	$-B$	0	$-D + E$	F	$-F$	0	H	0
G	$-C$	$-E - 2D$	0	$-G$	0	H	0	0
H	$-D - 2E$	$-F$	$-G$	0	$-H$	0	0	0

Table 2.

Assuming that $\dim \mathfrak{G}\mathfrak{M}(\mathcal{E}_1) = 8$, taking account of (5.7), after making some linear combinations, there must exist eight generators of the form

$$(5.14) \quad \begin{cases} A' := \partial_y + O(1), & E' := y \partial_y + O(2), \\ B' := \partial_x + O(1), & F' := y \partial_x + O(2), \\ C' := x \partial_y + O(2), & G' := xx \partial_x + xy \partial_y + O(3), \\ D' := x \partial_x + O(2), & H' := xy \partial_x + yy \partial_y + O(3). \end{cases}$$

To insure that the Lie brackets between these vector fields are small perturbations of the model ones, we can in advance replace (x, y) by $(\varepsilon x, \varepsilon y)$, so that $y_{xx} = \varepsilon F(\varepsilon x, \varepsilon y, y_x)$ is an $O(\varepsilon)$, hence all the remainders $O(1)$, $O(2)$ and $O(3)$ above are also $O(\varepsilon)$. It follows that the structure constants for A', \dots, H' are ε -close to those of Table 2.

Theorem 5.15. ([OV1994]) *Every semisimple Lie algebra over \mathbb{R} or \mathbb{C} is rigid: small deformations of the structure constants just give isomorphic Lie algebras.*

Consequently, there exists a change of basis close to the identity leading to new generators $A'', B'', \dots, G'', H''$ having exactly the same structure constants as in Table 2. Then $A''(0)$ and $B''(0)$ are still linearly independent. Since $[A'', B''] = [A, B] = 0$, there exist local coordinates (X, Y) centered at 0 in which $A'' = \partial_X$ and $B'' = \partial_Y$. Since $[A'', C''] = [A, C] = 0$ and $[B'', C''] = [B, C] = A$, it follows that $C'' = X \partial_Y$. The tangency to $0 = -Y_2 + F(X, Y, Y_1)$ (with $F(0) = 0$) of $(\partial_X)^{(2)} = \partial_X$, of $(\partial_Y)^{(2)} = \partial_Y$ and of $(X \partial_Y)^{(2)} = X \partial_Y + \partial_{Y_1}$ yields $F = 0$. \square

Open question 5.16. Does this proof generalize to $y_{x^{\kappa+1}} = F(x, y, y_x, \dots, y_{x^\kappa})$?

5.17. Complete system of second order. We now summarize a generalization to (\mathcal{E}_2) . According to Section 7 below, one may assume that the submanifold of solutions is $y = b + \sum_{i=1}^n a^i [x^i + O(|x|^2) + O(a) + O(b)]$, whence $y_{x^{i_1 x^{i_2}}} = F_{i_1, i_2}(x^i, y, y_{x^k})$ with $F(x, y, 0) \equiv 0$. Applying to the skeleton $0 = -y_{i_1, i_2} + F_{i_1, i_2}(x^i, y, y_k)$ a second prolongation $\mathcal{L}^{(2)}$ having coefficients \mathbf{Y}_{i_1} given by (3.9)(II) and \mathbf{Y}_{i_1, i_2} given by (3.20)(II), we get

$$(5.18) \quad 0 = -\mathbf{Y}_{i_1, i_2} + \sum_{k=1}^n [\mathcal{X}^k] \frac{\partial F_{i_1, i_2}}{\partial x^k} + [\mathcal{Y}] \frac{\partial F_{i_1, i_2}}{\partial y} + \sum_{k=1}^n [\mathbf{Y}_k] \frac{\partial F_{i_1, i_2}}{\partial y_k}.$$

Replacing y_{i_1, i_2} everywhere by $F_{i_1, i_2} = y_1 R + \dots + y_n R$, developping in powers of the pure jet variables y_l and picking the coefficients of cst. , of y_k , of $(y_k)^2$ and of $(y_k)^3$, we get the linear system

$$(5.19) \quad \begin{cases} \mathcal{Y}_{x^{i_1 x^{i_2}}} = \mathbf{P}(\mathcal{Y}_{x^l}) \\ \delta_{i_1}^k \mathcal{Y}_{x^{i_2 y}} + \delta_{i_2}^k \mathcal{Y}_{x^{i_1 y}} - \mathcal{X}_{x^{i_1 x^{i_2}}}^k = \mathbf{P}(\mathcal{Y}_y, \mathcal{X}_{x^{i_1}}^{l_2}, \mathcal{X}^l, \mathcal{Y}, \mathcal{Y}_{x^l}) \\ \delta_{i_1, i_2}^{k, k} \mathcal{Y}_{yy} - \delta_{i_1}^k \mathcal{X}_{x^{i_2 y}}^k - \delta_{i_2}^k \mathcal{X}_{x^{i_1 y}}^k = \mathbf{P}(\mathcal{Y}_y, \mathcal{X}_{x^{i_1}}^{l_2}, \mathcal{X}_y^l, \mathcal{X}^l, \mathcal{Y}, \mathcal{Y}_{x^l}) \\ \delta_{i_1, i_2}^{k, k} \mathcal{X}_{yy}^k = \mathbf{P}(\mathcal{Y}_y, \mathcal{X}_{x^{i_1}}^{l_2}, \mathcal{X}_y^l, \mathcal{X}^l, \mathcal{Y}, \mathcal{Y}_{x^l}), \end{cases}$$

upon which obvious linear combinations yield a known generalization of Proposition 5.9.

Proposition 5.20. ([Su2001, GM2003a]) An infinitesimal Lie symmetry $\sum_{k=1}^n \mathcal{X}^k \frac{\partial}{\partial x^k} + \mathcal{Y} \frac{\partial}{\partial y}$ is uniquely determined by the $n^2 + 4n + 3$ initial Taylor coefficients:

$$(5.21) \quad \mathcal{X}^l(0), \mathcal{Y}(0), \mathcal{X}_{x^{i_1}}^{l_2}(0), \mathcal{X}_y^l(0), \mathcal{Y}_{x^l}(0), \mathcal{Y}_y(0), \mathcal{Y}_{x^l y}(0), \mathcal{Y}_{yy}(0).$$

The bound $\dim \mathfrak{Sym}(\mathcal{E}_2) \leq n^2 + 4n + 3$ is attained with $F_{i_1, i_2} = 0$, whence all $\mathbf{P} = 0$ and

$$(5.22) \quad \begin{cases} A := \partial_y, & E := y \partial_y, \\ B_i := \partial_{x^i}, & F_i := y \partial_{x^i}, \\ C_i := x^i \partial_y, & G_i := x^i (x^1 \partial_{x^1} + \dots + x^n \partial_{x^n} + y \partial_y) + xy \partial_y, \\ D_{i, k} := x^i \partial_{x^k}, & H := y (x^1 \partial_{x^1} + \dots + x^n \partial_{x^n} + y \partial_y). \end{cases}$$

are infinitesimal generators of the group $\text{PGL}_{n+2}(\mathbb{K}) = \text{Aut}(P_{n+1}(\mathbb{K}))$ of projective transformations

$$(5.23) \quad (x, y) \mapsto \left(\frac{\alpha_1 x^1 + \dots + \alpha_n x^n + \beta y + \gamma}{\lambda_1 x^1 + \dots + \lambda_n x^n + \mu y + \nu}, \frac{\delta_1 x^1 + \dots + \delta_n x^n + \eta y + \epsilon}{\lambda_1 x^1 + \dots + \lambda_n x^n + \mu y + \nu} \right)$$

stabilizing the collections of all affine planes of \mathbb{K}^{n+1} , namely the solutions of the model equation $y_{x^{i_1 x^{i_2}}} = 0$. The model Lie algebra $\mathfrak{pgl}_{n+2}(\mathbb{K}) \simeq \mathfrak{sl}_{n+2}(\mathbb{K})$ is simple, hence rigid.

Theorem 5.24. The bound $\dim \mathfrak{Sym}(\mathcal{E}_2) \leq n^2 + 4n + 3$ is attained if and only if (\mathcal{E}_2) is equivalent, through a diffeomorphism $(x^i, y) \mapsto (X^k, Y)$, to $Y_{X^{k_1} X^{k_2}} = 0$.

The proof, similar to that of Theorem 5.13, is skipped.

The study of (\mathcal{E}_3) also leads to the model algebra $\mathfrak{pgl}_{n+2}(\mathbb{K}) \simeq \mathfrak{sl}_{n+2}(\mathbb{K})$ and an analog to Theorem 5.13 holds. Details are similar.

§6. TRANSFER OF LIE SYMMETRIES TO THE PARAMETER SPACE

6.1. Stabilization of foliations. As announced in §2.38, we now transfer the theory of Lie symmetries to submanifolds of solutions.

Restarting from §4.1, let φ a Lie symmetry of (\mathcal{E}) , namely $\varphi_{\Delta_{\mathcal{E}}}$ stabilizes $F_{\Delta_{\mathcal{E}}}$. The diffeomorphism A defined by (2.9) transforms F_v to $F_{\Delta_{\mathcal{E}}}$. Conjugating, we get the self-transformation $A^{-1} \circ \varphi_{\Delta_{\mathcal{E}}} \circ A$ of the (x, a, b) -space that must stabilize also the foliation F_v . Equivalently, it must have expression:

$$(6.2) \quad [A^{-1} \circ \varphi_{\Delta_{\mathcal{E}}} \circ A](x, a, b) = (\theta(x, a, b), f(a, b), g(a, b)) \in \mathbb{K}^n \times \mathbb{K}^p \times \mathbb{K}^m,$$

where, importantly, the last two components are independent of the coordinate x , because the leaves of F_v are just $\{a = \text{cst.}, b = \text{cst.}\}$.

Lemma 6.3. *To every Lie symmetry φ of (\mathcal{E}) , there corresponds a transformation of the parameters*

$$(6.4) \quad (a, b) \longmapsto (f(a, b), g(a, b)) =: h(a, b)$$

meaning that φ transforms the local solution $y_{a,b}(x) := \Pi(x, a, b)$ to the local solution $y_{h(a,b)}(x) = \Pi(x, h(a, b))$.

Unfortunately, the expression of $A^{-1} \circ \varphi_{\Delta_{\mathcal{E}}} \circ A$ does not clearly show that f and g are independent of x . Indeed, reminding the expressions of A and of Φ , we have:

$$(6.5) \quad \varphi_{\Delta_{\mathcal{E}}} \circ A(x, a, b) = \left(\varphi(x, \Pi(x, a, b)), \Phi_{\beta(q)}^{j(q)}(x^{i_1}, \Pi^{j_1}(x, a, b), \Pi_{x, \beta(q_1)}^{j(q_1)}(x, a, b)) \right).$$

To compose with A^{-1} whose expression is given by (2.21), it is useful to split $\varphi = (\phi, \psi) \in \mathbb{K}^n \times \mathbb{K}^m$, so above we write

$$(6.6) \quad \varphi(x, \Pi(x, a, b)) = (\phi(x, \Pi(x, a, b)), \psi(x, \Pi(x, a, b))),$$

and finally, dropping the arguments:

$$(6.7) \quad [A^{-1} \circ \varphi_{\Delta_{\mathcal{E}}} \circ A](x, a, b) = \left(\phi^i, A^q(\phi^{i_1}, \psi^{j_1}, \Phi_{\beta(q_1)}^{j(q_1)}), B^j(\phi^{i_1}, \psi^{j_1}, \Phi_{\beta(q_1)}^{j(q_1)}) \right).$$

In case $(\mathcal{E}) = (\mathcal{E}_1)$, is an exercise to verify by computations that the $A^q(\cdot)$ and $B^j(\cdot)$ are independent of x . In general however, the explicit expression of $\Phi_{i_1, \dots, i_\lambda}^j$ is unknown. Unfortunately also, nothing shows how $(f(a, b), g(a, b))$ is uniquely associated to $\varphi(x, y)$. Further explanations are needed.

6.8. Determination of parameter transformations. At first, we state a geometric reformulation of the preceding lemma.

Lemma 6.9. *Every Lie symmetry $(x, y) \mapsto \varphi(x, y)$ of (\mathcal{E}) induces a local \mathbb{K} -analytic diffeomorphism*

$$(6.10) \quad (x, y, a, b) \longmapsto (\varphi(x, y), h(a, b))$$

of $\mathbb{K}_x^n \times \mathbb{K}_y^m \times \mathbb{K}_a^p \times \mathbb{K}_b^m$ that maps to itself the associated submanifold of solutions

$$(6.11) \quad \mathcal{M}_{\mathcal{E}} = \{(x, y, a, b) : y = \Pi(x, a, b)\}.$$

Proof. In fact, we know that the n -dimensional leaf $\{(x, \Pi(x, a, b)) : x \in \mathbb{K}^n\}$ is sent $\{(x, \Pi(x, h(a, b))) : x \in \mathbb{K}^n\}$. \square

Equivalently, setting $c := (a, b)$ and writing $(\varphi, h) = (\phi, \psi, h)$, we have $\psi = \Pi(\phi, h)$ when $y = \Pi(x, c)$, namely

$$(6.12) \quad \psi(x, \Pi(x, c)) \equiv \Pi(\phi(x, \Pi(x, c)), h(c))$$

Proposition 6.13. *There exists a universal rational map \widehat{H} such that*

$$(6.14) \quad h(c) \equiv \widehat{H}\left(J_{x,a,b}^{\kappa+1} \Pi(x, c), J_{x,y}^{\kappa} \varphi(x, \Pi(x, c))\right)$$

This shows unique determination of h from φ , given (\mathcal{E}) or equivalently, given Π .

Proof. Differentiating a function $\chi(x, \Pi(x, c))$ with respect to x^k , $k = 1, \dots, n$, corresponds to applying to χ the vector field

$$(6.15) \quad L_k := \frac{\partial}{\partial x_k} + \sum_{j=1}^m \frac{\partial \Pi^j}{\partial x_k}(x, c) \frac{\partial}{\partial y^j}, \quad k = 1, \dots, n.$$

Thus, applying L_k to the m scalar equations (6.12), we get

$$(6.16) \quad L_k \psi^j = \sum_{l=1}^n \frac{\partial \Pi^j}{\partial x^l} L_k \phi^l,$$

for $1 \leq k \leq n$ and $1 \leq j \leq m$. It follows from the assumption that φ is a local diffeomorphism that $\det(L_k \phi^l(0))_{1 \leq l \leq n}^{1 \leq l \leq n} \neq 0$ also. So we may solve the first derivatives Π_x above: there exist universal polynomials S_l^j such that

$$(6.17) \quad \frac{\partial \Pi^j}{\partial x^l} = \frac{S_l^j \left(\{L_{k'} \varphi^{i'}\}_{1 \leq i' \leq n+m}^{1 \leq k' \leq n} \right)}{\det(L_{k'} \phi^{l'})_{1 \leq l' \leq n}^{1 \leq k' \leq n}}.$$

Again, we apply the L_k to these equations, getting, thanks to the chain rule:

$$(6.18) \quad \sum_{l_2=1}^n \frac{\partial^2 \Pi^j}{\partial x^{l_1} \partial x^{l_2}} L_k \phi^{l_2} = \frac{R_{l_1, k}^j \left(\{L_{k'_1} L_{k'_2} \varphi^{i'}\}_{1 \leq i' \leq n+m}^{1 \leq k'_1, k'_2 \leq n} \right)}{\left[\det(L_{k'} \phi^{l'})_{1 \leq l' \leq n}^{1 \leq k' \leq n} \right]^2}.$$

Here, $R_{l_1, k}^j$ are universal polynomials. Solving the second derivatives $\Pi_{x^{l_1} x^{l_2}}^j$, we get

$$(6.19) \quad \frac{\partial^2 \Pi^j}{\partial x^{l_1} \partial x^{l_2}} = \frac{S_{l_1, l_2}^j \left(\{L_{k'_1} L_{k'_2} \varphi^{i'}\}_{1 \leq i' \leq n+m}^{1 \leq k'_1, k'_2 \leq n} \right)}{\left[\det(L_{k'} \phi^{l'})_{1 \leq l' \leq n}^{1 \leq k' \leq n} \right]^3}.$$

By induction, for every $\beta \in \mathbb{N}^n$:

$$(6.20) \quad \frac{\partial^{|\beta|} \Pi^j}{\partial x^\beta} = \frac{S_\beta^j \left(\{L^{\beta'} \varphi^{i'}\}_{|\beta'| \leq |\beta|}^{1 \leq i' \leq n+m} \right)}{\left[\det(L_{k'} \phi^{l'})_{1 \leq l' \leq n}^{1 \leq k' \leq n} \right]^{2|\beta|+1}},$$

where S_β^j are universal polynomials. Here, for $\beta' \in \mathbb{N}^n$, we denote by $L^{\beta'}$ the derivation of order $|\beta'|$ defined by $(L_1)^{\beta'_1} \dots (L_n)^{\beta'_n}$.

Next, thanks to the assumption that \mathcal{M} is solvable with respect to the parameters, there exist integers $j(1), \dots, j(p)$ with $1 \leq j(q) \leq m$ and multiindices $\beta(1), \dots, \beta(p) \in \mathbb{N}^n$ with $|\beta(q)| \geq 1$ and $\max_{1 \leq q \leq p} |\beta(q)| = \kappa$ such that the local \mathbb{K} -analytic map

$$(6.21) \quad \mathbb{K}^{p+m} \ni c \mapsto \left((\Pi^j(0, c))^{1 \leq j \leq m}, \left(\frac{\partial^{|\beta(q)|} \Pi^{j(q)}}{\partial x^{\beta(q)}}(0, c) \right)_{1 \leq q \leq p} \right) \in \mathbb{K}^{p+m}$$

has rank $p + m$ at $c = 0$. We then consider in (6.20) only the $(p + m)$ equations written for $(j, 0)$, $(j(q), \beta(q))$ and we solve $h(c)$ by means of the analytic implicit function theorem: (6.22)

$$h = \widehat{H} \left(\phi, \frac{\mathcal{S}_{\beta(1)}^{j(1)} \left(\left\{ \mathbb{L}^{\beta'} \varphi^{i'} \right\}_{\substack{1 \leq i' \leq n+m \\ |\beta'| \leq |\beta(1)|}} \right)}{\det \left[\left(\mathbb{L}_{k'} \phi^{l'} \right)_{\substack{1 \leq l' \leq n \\ 1 \leq k' \leq n}} \right]^{2|\beta(1)|+1}}, \dots, \frac{\mathcal{S}_{\beta(p)}^{j(p)} \left(\left\{ \mathbb{L}^{\beta'} \varphi^{i'} \right\}_{\substack{1 \leq i' \leq n+m \\ |\beta'| \leq |\beta(p)|}} \right)}{\det \left[\left(\mathbb{L}_{k'} \phi^{l'} \right)_{\substack{1 \leq l' \leq n \\ 1 \leq k' \leq n}} \right]^{2|\beta(p)|+1}} \right).$$

Finally, by developping every derivative $\mathbb{L}^{\beta'} \varphi^{i'}$ (including $\mathbb{L}_{k'} \phi^{l'}$ as a special case), taking account of the fact that the coefficients of the $\mathbb{L}_{k'}$ depend directly on Π , we get some universal polynomial $P_{\beta'}(J_x^{|\beta'|+1} \Pi, J_{x,y}^{|\beta'|} \varphi^{i'})$. Inserting above, we get the map \widehat{H} . \square

6.23. Pseudogroup of twin transformations. The previous considerations lead to introducing the following.

Definition 6.24. By $G_{v,p}$, we denote the infinite-dimensional (pseudo)group of local \mathbb{K} -analytic diffeomorphisms

$$(6.25) \quad (x, y, a, b) \longmapsto (\varphi(x, y), h(a, b))$$

that respect the separation between the variables and the parameters.

A converse to Lemma 6.3 holds.

Lemma 6.26. *Let \mathcal{M} be a submanifold $y = \Pi(x, a, b)$ that is solvable with respect to the parameters (a, b) . If a local \mathbb{K} -analytic diffeomorphism $(x, y, a, b) \longmapsto (\varphi(x, y), h(a, b))$ of $\mathbb{K}_x^n \times \mathbb{K}_y^m \times \mathbb{K}_a^p \times \mathbb{K}_b^m$ belonging to $G_{v,p}$ sends \mathcal{M} to \mathcal{M} , then $(x, y) \mapsto \varphi(x, y)$ is a Lie symmetry of the PDE system $\mathcal{E}_{\mathcal{M}}$ associated to \mathcal{M} .*

Proof. In fact, since (φ, h) respects the separation of variables and stabilizes \mathcal{M} , it respects the fundamental pair of foliations (F_v, F_p) , namely $\{(a, b) = (a_0, b_0)\} \cap \mathcal{M}$ is sent to $\{(a, b) = h(a_0, b_0)\} \cap \mathcal{M}$ and $\{(x, y) = (x_0, y_0)\} \cap \mathcal{M}$ is sent to $\{(x, y) = \varphi(x_0, y_0)\} \cap \mathcal{M}$. Hence $\varphi_{\Delta_{\mathcal{E}_{\mathcal{M}}}}$ also stabilizes $F_{\Delta_{\mathcal{E}}}$. \square

Corollary 6.27. *Through the one-to-one correspondence $(\mathcal{E}) \longleftrightarrow \mathcal{M}$ of Proposition 2.17, Lie symmetries of (\mathcal{E}) correspond to elements of $G_{v,p}$ which stabilize \mathcal{M} .*

Definition 6.28. Let $\text{Aut}_{v,p}(\mathcal{M})$ denote the local (pseudo)group of $(\varphi, h) \in G_{v,p}$ stabilizing \mathcal{M} . Let $\text{Lie}(\mathcal{E})$ denote the local (pseudo)group of Lie symmetries of (\mathcal{E}) .

In summary:

$$(6.29) \quad \boxed{\text{Lie}(\mathcal{E}) \simeq \text{Aut}_{v,p}(\mathcal{M}_{(\mathcal{E})})} \quad \text{and} \quad \boxed{\text{Aut}_{v,p}(\mathcal{M}) \simeq \text{Lie}(\mathcal{E}_{\mathcal{M}})}.$$

6.30. Transfer of infinitesimal Lie symmetries. Let $\mathcal{L} \in \mathfrak{G}\mathfrak{Y}\mathfrak{M}(\mathcal{E})$, i.e. $\mathcal{L}_{\Delta_{\mathcal{E}}}$ is tangent to $\Delta_{\mathcal{E}}$. Through the diffeomorphism A , the push-forward of $\mathcal{L}_{\Delta_{\mathcal{E}}}$ must be of the form

$$(6.31) \quad A_*^{-1}(\mathcal{L}_{\Delta_{\mathcal{E}}}) = \sum_{k=1}^n \Theta^k(x, a, b) \frac{\partial}{\partial x^k} + \sum_{q=1}^p \mathcal{F}^q(a, b) \frac{\partial}{\partial a^q} + \sum_{j=1}^m \mathcal{G}^j(a, b) \frac{\partial}{\partial b^j},$$

where the last two families of \mathbb{K} -analytic coefficients \mathcal{F}^q and \mathcal{G}^j depend only on (a, b) .

Lemma 6.32. *To every infinitesimal symmetry \mathcal{L} of (\mathcal{E}) , we can associate an infinitesimal symmetry*

$$(6.33) \quad \mathcal{L}^* := \sum_{q=1}^p \mathcal{F}^q(a, b) \frac{\partial}{\partial a^q} + \sum_{j=1}^m \mathcal{G}^j(a, b) \frac{\partial}{\partial b^j}$$

of the space of parameters which tells how the flow of \mathcal{L} acts infinitesimally on the leaves of $F_{\Delta\mathcal{E}}$. Furthermore, $\mathcal{L} + \mathcal{L}^*$ is tangent to the submanifold of solutions $\mathcal{M}_{(\mathcal{E})}$.

Considering the flow of $\mathcal{L} + \mathcal{L}^*$ reduces these assertions and the next to the arguments of the preceding paragraphs. So we summarize.

Lemma 6.34. *Let \mathcal{M} be a submanifold $y = \Pi(x, a, b)$ that is solvable with respect to the parameters (a, b) . If a vector field that respects the separation between variables and parameters, namely of the form*

$$(6.35) \quad \mathcal{L} + \mathcal{L}^* = \sum_{i=1}^n \mathcal{X}^i(x, y) \frac{\partial}{\partial x^i} + \sum_{j=1}^m \mathcal{Y}^j(x, y) \frac{\partial}{\partial y^j} + \sum_{q=1}^p \mathcal{F}^q(a, b) \frac{\partial}{\partial a^q} + \sum_{j=1}^m \mathcal{G}^j(a, b) \frac{\partial}{\partial b^j}$$

is tangent to \mathcal{M} , then \mathcal{L} is an infinitesimal Lie symmetry of $(\mathcal{E}_{\mathcal{M}})$

Corollary 6.36. *Through the one-to-one correspondence $(\mathcal{E}) \longleftrightarrow \mathcal{M}$ of Proposition 2.17, infinitesimal Lie symmetries of (\mathcal{E}) correspond to vector fields $\mathcal{L} + \mathcal{L}^*$ tangent to \mathcal{M} .*

Definition 6.37. Let $\mathfrak{Sym}(\mathcal{M})$ denote the Lie algebra of vector fields $\mathcal{L} + \mathcal{L}^*$ tangent to \mathcal{M} . Let $\mathfrak{Sym}(\mathcal{E})$ denote the Lie algebra of infinitesimal Lie symmetries of (\mathcal{E}) .

In summary:

$$(6.38) \quad \boxed{\mathfrak{Sym}(\mathcal{E}) \simeq \mathfrak{Sym}(\mathcal{M}_{(\mathcal{E})})} \quad \text{and} \quad \boxed{\mathfrak{Sym}(\mathcal{M}) \simeq \mathfrak{Sym}(\mathcal{E}_{\mathcal{M}})}.$$

6.39. Dual defining equations. As in §2.10, let $\mathcal{M} \subset \mathbb{K}_x^n \times \mathbb{K}_y^m \times \mathbb{K}_a^p \times \mathbb{K}_b^m$ given by $0 = -y^j + \Pi^j(x, a, b)$ and assume it to be solvable with respect to the parameters. In particular, we can solve the b^j , obtaining *dual defining equations*

$$(6.40) \quad b^j \equiv \Pi^{*j}(a, x, y), \quad j = 1, \dots, m,$$

for some local \mathbb{K} -analytic map $\Pi^* = (\Pi^{*1}, \dots, \Pi^{*m})$ satisfying

$$(6.41) \quad b \equiv \Pi^*(a, x, \Pi(x, a, b)) \quad \text{and} \quad y \equiv \Pi(x, a, \Pi^*(a, x, y)).$$

6.42. An algorithm for the computation of $\mathfrak{Sym}(\mathcal{M})$. The tangency to \mathcal{M} is expressed by applying the vector field (6.35) to $0 = -y^j + \Pi^j(x, a, b)$, which yields:

$$(6.43) \quad 0 = -\mathcal{Y}^j(x, y) + \sum_{i=1}^n \mathcal{X}^i(x, y) \Pi_{x^i}^j(x, a, b) + \sum_{q=1}^p \mathcal{F}^q(a, b) \Pi_{a^q}^j(x, a, b) + \sum_{l=1}^m \mathcal{G}^l(a, b) \Pi_{b^l}^j(x, a, b),$$

for $j = 1, \dots, m$ and for $(x, y, a, b) \in \mathcal{M}$. In fact, after replacing the variable y by $\Pi(x, a, b)$, these equations should be interpreted as power series identities in $\mathbb{K}\{x, a, b\}$.

Denote by $\Delta(x, a, b)$ the determinant of the (invertible) matrix $(\Pi_{b^l}^j(x, a, b))_{1 \leq l, j \leq m}$ and by $D(x, a, b)$ its matrix of cofactors, so that $\Pi_b^{-1} = [\Delta]^{-1} D$. Hence we can solve \mathcal{G} from (6.43):

$$(6.44) \quad \left\{ \begin{array}{l} \mathcal{G}(a, b) \equiv \frac{D(x, a, b)}{\Delta(x, a, b)} \left[\mathcal{Y}(x, \Pi(x, a, b)) - \sum_{i=1}^n \mathcal{X}^i(x, \Pi(x, a, b)) \Pi_{x^i}(x, a, b) - \right. \\ \left. - \sum_{q=1}^p \mathcal{F}^q(a, b) \Pi_{a^q}(x, a, b) \right]. \end{array} \right.$$

Next, we aim to solve the $\mathcal{F}^q(a, b)$. Consequently, we gather all the other terms in the brackets as $\Psi_0(J_{x,a,b}^1 \Pi, \mathcal{X}, \mathcal{Y})$:

$$(6.45) \quad \mathcal{G}(a, b) \equiv \frac{D(x, a, b)}{\Delta(x, a, b)} \left[- \sum_{q=1}^p \mathcal{F}^q(a, b) \Pi_{a^q}(x, a, b) \right] + \frac{\Psi_0(J_{x,a,b}^1 \Pi, \mathcal{X}, \mathcal{Y})}{\Delta(x, a, b)}.$$

Here, Ψ_0 is linear with respect to $(\mathcal{X}, \mathcal{Y})$, with polynomial coefficients of degree one in $J_{x,a,b}^1 \Pi$.

Next, for $k = 1, \dots, n$, we differentiate this identity with respect to x_k . Then $\mathcal{G}(a, b)$ disappears and we chase the denominator Δ^2 :

$$(6.46) \quad \left\{ \begin{array}{l} 0 \equiv [\Delta D] \left[- \sum_{q=1}^p \mathcal{F}^q(a, b) \Pi_{a^q x^k}(x, a, b) \right] + \\ \quad + [\Delta D_{x_k} - \Delta_{x_k} D] \left[- \sum_{q=1}^p \mathcal{F}^q(a, b) \Pi_{a^q}(x, a, b) \right] + \\ \quad + \Psi_k(J_{x,a,b}^2 \Pi, J_{x,y}^1 \mathcal{X}, J_{x,y}^1 \mathcal{Y}). \end{array} \right.$$

The Ψ_k are linear with respect to $(J_{x,y}^1 \mathcal{X}, J_{x,y}^1 \mathcal{Y})$, with polynomial coefficients in $J_{x,a,b}^2 \Pi$. Then we further differentiate with respect to x and by induction, for every $\beta \in \mathbb{N}^n$, we get:

$$(6.47) \quad \left\{ \begin{array}{l} 0 \equiv [\Delta D] \left[- \sum_{q=1}^p \mathcal{F}^q(a, b) \Pi_{a^q x^\beta}(x, a, b) \right] + \\ \quad + \sum_{|\beta_1| < |\beta|} D_{\beta, \beta_1} (J^{|\beta_1|+1} \Pi) \left[- \sum_{q=1}^p \mathcal{F}^q(a, b) \Pi_{a^q x^{\beta_1}}(x, a, b) \right] + \\ \quad + \Psi_\beta (J_{x,a,b}^{|\beta|+1} \Pi, J_{x,y}^{|\beta|} \mathcal{X}, J_{x,y}^{|\beta|} \mathcal{Y}), \end{array} \right.$$

where the expressions D_{β, β_1} are certain $m \times m$ matrices with polynomial coefficients in the jet $J_{x,a,b}^{|\beta_1|+1} \Pi$, and where the terms $\Psi_\beta (J_{x,a,b}^{|\beta|+1} \Pi, J_{x,y}^{|\beta|} \mathcal{X}, J_{x,y}^{|\beta|} \mathcal{Y})$ are linear with respect to $(J_{x,y}^{|\beta|} \mathcal{X}, J_{x,y}^{|\beta|} \mathcal{Y})$, with polynomial coefficients in $J_{x,a,b}^{|\beta|+1} \Pi$.

Writing these identity for $(j, \beta) = (j(q), \beta(q))$, $q = 1, \dots, p$, reminding $\max_{1 \leq q \leq p} |\beta(q)| = \kappa$, it follows from the assumption of solvability with respect to the parameters (a boring technical check is needed) that we may solve

$$(6.48) \quad \mathcal{F}^q(a, b) \equiv \Phi^q (J_{x,a,b}^{\kappa+1} \Pi(x, a, b), J_{x,y}^\kappa \mathcal{X}(x, \Pi(x, a, b)), J_{x,y}^\kappa \mathcal{Y}(x, \Pi(x, a, b))),$$

for $q = 1, \dots, p$, where each local \mathbb{K} -analytic function Φ_q is linear with respect to $(J^\kappa \mathcal{X}, J^\kappa \mathcal{Y})$ and rational with respect to $J^{\kappa+1} \Pi$, with denominator not vanishing at $(x, a, b) := (0, 0, 0)$.

Pursuing, we differentiate (6.48) with respect to x^l for $l = 1, \dots, n$. Then $\mathcal{F}^q(a, b)$ disappears and we get:

$$(6.49) \quad 0 \equiv \Phi_{q,l} (J_{x,a,b}^{\kappa+2} \Pi(x, a, b), J_{x,y}^{\kappa+1} \mathcal{X}(x, \Pi(x, a, b)), J_{x,y}^{\kappa+1} \mathcal{Y}(x, \Pi(x, a, b))),$$

for $1 \leq q \leq p$ and $1 \leq l \leq n$. In (6.46), we then replace the functions \mathcal{F}^q by their values Φ^q :

$$(6.50) \quad 0 \equiv \Psi_{k,j} (J_{x,a,b}^{\kappa+1} \Pi(x, a, b), J_{x,y}^\kappa \mathcal{X}(x, \Pi(x, a, b)), J_{x,y}^\kappa \mathcal{Y}(x, \Pi(x, a, b))),$$

for $1 \leq k \leq n$ and $1 \leq j \leq m$. Then we replace the variable b by $\Pi^*(a, x, y)$ in the two obtained systems (6.49) and (6.50); taking account of the functional identity $y \equiv \Pi(x, a, \Pi^*(a, x, y))$ written in (6.41), we get

$$(6.51) \quad \begin{cases} 0 \equiv \Phi_{q,l}(J_{x,a,b}^{\kappa+2}\Pi(x, a, \Pi^*(a, x, y)), J_{x,y}^{\kappa+1}\mathcal{X}(x, y), J_{x,y}^{\kappa+1}\mathcal{Y}(x, y)), \\ 0 \equiv \Psi_{k,j}(J_{x,a,b}^{\kappa+1}\Pi(x, a, \Pi^*(a, x, y)), J_{x,y}^{\kappa}\mathcal{X}(x, y), J_{x,y}^{\kappa}\mathcal{Y}(x, y)). \end{cases}$$

Finally, we developpe these equations in power series with respect to a :

$$(6.52) \quad \begin{cases} 0 \equiv \sum_{\gamma \in \mathbb{N}^p} a^\gamma \Phi_{q,l,\gamma}(x, y, J_{x,y}^{\kappa+1}\mathcal{X}(x, y), J_{x,y}^{\kappa+1}\mathcal{Y}(x, y)), \\ 0 \equiv \sum_{\gamma \in \mathbb{N}^p} a^\gamma \Psi_{k,j,\gamma}(x, y, J_{x,y}^{\kappa}\mathcal{X}(x, y), J_{x,y}^{\kappa}\mathcal{Y}(x, y)), \end{cases}$$

where the terms $\Phi_{q,l,\gamma}$ and $\Psi_{k,j,\gamma}$ are linear with respect to the jets of \mathcal{X}, \mathcal{Y} .

Proposition 6.53. *A vector field (6.35) belongs to $\mathfrak{S}\mathfrak{Y}\mathfrak{M}(\mathcal{M})$ if and only if $\mathcal{X}^i, \mathcal{Y}^j$ satisfy the linear PDE system*

$$(6.54) \quad \begin{cases} 0 \equiv \Phi_{q,l,\gamma}(x, y, J_{x,y}^{\kappa+1}\mathcal{X}(x, y), J_{x,y}^{\kappa+1}\mathcal{Y}(x, y)), \\ 0 \equiv \Pi_{k,j,\gamma}(x, y, J_{x,y}^{\kappa}\mathcal{X}(x, y), J_{x,y}^{\kappa}\mathcal{Y}(x, y)), \end{cases}$$

where $1 \leq q \leq p, 1 \leq l \leq n, 1 \leq k \leq n$ and $\gamma \in \mathbb{N}^p$. Then \mathcal{F}^q defined by (6.48) and \mathcal{G}^j defined by (6.45) are independent of x .

This provides a second algorithm, essentially equivalent to Sophus Lie's.

Example 6.55. For $y_{xx}(x) = F(x, y(x), y_x(x))$, the first line of (6.54) is (the second one is redundant):

$$(6.56) \quad \left\{ \begin{aligned} 0 \equiv & \mathcal{X} [-\Pi_{xa}\Pi_{xxx}\Pi_b + \Pi_a\Pi_{xb}\Pi_{xxx} - \Pi_x\Pi_{xxa}\Pi_{xb} + \Pi_{xa}\Pi_x\Pi_{xxb} + \\ & + \Pi_{xxa}\Pi_{xx}\Pi_b - \Pi_a\Pi_{xxb}\Pi_{xx}] + \\ & + \mathcal{Y} [-\Pi_{xa}\Pi_{xxb} + \Pi_{xxa}\Pi_{xb}] + \\ & + \mathcal{X}_x [-2\Pi_{xx}\Pi_{xa}\Pi_b + 2\Pi_{xx}\Pi_a\Pi_{xb} + \Pi_x\Pi_b\Pi_{xxa} - \Pi_x\Pi_a\Pi_{xxb}] + \\ & + \mathcal{Y}_x [-\Pi_b\Pi_{xxa} + \Pi_a\Pi_{xxb}] + \\ & + \mathcal{X}_y [-3\Pi_x\Pi_{xx}\Pi_{xa}\Pi_b + 3\Pi_x\Pi_a\Pi_{xx}\Pi_{xb} + (\Pi_x)^2\Pi_b\Pi_{xxa} - (\Pi_x)^2\Pi_a\Pi_{xxb}] + \\ & + \mathcal{Y}_y [\Pi_{xx}\Pi_b\Pi_{xa} - \Pi_{xx}\Pi_a\Pi_{xb} - \Pi_x\Pi_b\Pi_{xxa} + \Pi_x\Pi_a\Pi_{xxb}] + \\ & + \mathcal{X}_{xx} [-\Pi_x\Pi_b\Pi_{xa} + \Pi_x\Pi_a\Pi_{xb}] + \\ & + \mathcal{X}_{xy} [-2(\Pi_x)^2\Pi_b\Pi_{xa} + 2(\Pi_x)^2\Pi_a\Pi_{xb}] + \\ & + \mathcal{X}_{y^2} [-(\Pi_x)^3\Pi_b\Pi_{xa} + (\Pi_x)^3\Pi_a\Pi_{xb}] + \\ & + \mathcal{Y}_{xx} [\Pi_b\Pi_{xa} - \Pi_a\Pi_{xb}] + \\ & + \mathcal{Y}_{xy} [2\Pi_x\Pi_b\Pi_{xa} - 2\Pi_x\Pi_a\Pi_{xb}] + \\ & + \mathcal{Y}_{y^2} [(\Pi_x)^2\Pi_b\Pi_{xa} - (\Pi_x)^2\Pi_a\Pi_{xb}]. \end{aligned} \right.$$

We observe the similarity with (4.19): the expression is linear in the partial derivatives of \mathcal{X}, \mathcal{Y} of order ≤ 2 , but the coefficients in the equation above are more complicated. In fact, after dividing by $-\Pi_b\Pi_{xa} + \Pi_a\Pi_{xb}$, this equation coincides with (4.21), thanks to $\Pi_x = y_1$ and to the formulas (2.34) for F_x, F_y, F_{y_1} .

6.57. Infinitesimal CR automorphisms of generic submanifolds. If the system (\mathcal{E}) is associated to the complexification $\mathcal{M} = (M)^c$ of a generic $M \subset \mathbb{C}^{n+m}$ as in §1.16, then $a = (\bar{z})^c = \zeta$, $b = (\bar{w})^c = \xi$, and the vector field \mathcal{L}^* associated to an infinitesimal Lie symmetry

$$(6.58) \quad \mathcal{L} = \sum_{i=1}^n \mathcal{X}^i(z, w) \frac{\partial}{\partial z^i} + \sum_{j=1}^m \mathcal{Y}^j(z, w) \frac{\partial}{\partial w^j}$$

of (\mathcal{E}) is simply the complexification $\underline{\mathcal{L}}$ of its conjugate $\bar{\mathcal{L}}$, namely

$$(6.59) \quad \mathcal{L}^* = \underline{\mathcal{L}} = \sum_{i=1}^n \bar{\mathcal{X}}^i(\zeta, \xi) \frac{\partial}{\partial \zeta^i} + \sum_{j=1}^m \bar{\mathcal{Y}}^j(\zeta, \xi) \frac{\partial}{\partial \xi^j}.$$

Then the sum $\mathcal{L} + \underline{\mathcal{L}}$ is tangent to \mathcal{M} and its flow stabilizes the two invariant foliations, obtained by intersecting \mathcal{M} by $\{(z, w) = \text{cst.}\}$ or by $\{(\zeta, \xi) = \text{cst.}\}$. In [Me2005a, Me2005b], these two foliations, denoted \mathcal{F} , $\underline{\mathcal{F}}$, are called (conjugate) Segre foliations, since its leaves are the complexifications of the (conjugate) classical Segre varieties ([Se1931, Pi1975, Pi1978, We1977, DW1980, BJT1985, DF1988, BER1999, Su2001, Su2002, Su2003, GM2003a]) associated to M , viewed in its ambient space \mathbb{C}^{n+m} . The next definition is also classical ([Be1979, Lo1981, EKV1985, Kr1987, KV1987, Be1988, Vi1990, St1996, Be1997, BER1999, Lo2002, FK2005a, FK2005b]):

Definition 6.60. By $\mathfrak{hol}(M)$ is meant the Lie algebra of local holomorphic vector fields $\mathcal{L} = \sum_{i=1}^n \mathcal{X}^i(z, w) \frac{\partial}{\partial z^i} + \sum_{j=1}^m \mathcal{Y}^j(z, w) \frac{\partial}{\partial w^j}$ whose real flow $\exp(t\mathcal{L})(z, w)$ induces one-parameter families of local biholomorphic transformations of \mathbb{C}^{n+m} stabilizing M . Equivalently,

$$(6.61) \quad 2 \operatorname{Re} \mathcal{L} = \mathcal{L} + \bar{\mathcal{L}}$$

is tangent to M . Again equivalently, $\mathcal{L} + \underline{\mathcal{L}}$ is tangent to $\mathcal{M} = M^c$.

Then obviously $\mathfrak{hol}(M)$ is a real Lie algebra.

Theorem 6.62. ([Ca1932a, BER1999, GM2004]) *The complexification $\mathfrak{hol}(M) \otimes \mathbb{C}$ identifies with $\mathfrak{S}\mathfrak{M}(\mathcal{E}(M^c))$. Furthermore, if M is finitely nondegenerate and minimal at the origin, both are finite-dimensional and $\mathfrak{hol}(M)$ is totally real in $\mathfrak{S}\mathfrak{M}(\mathcal{E}(M^c))$.*

The minimality assumption is sometimes presented by saying that the Lie algebra generated by $T^c M$ generates TM at the origin ([BER1999]). However, it is more natural to proceed with the fundamental pair of foliations associated to \mathcal{M} ([Me2001, GM2004, Me2005a, Me2005b]). Anticipating Sections 10 and 11 to which the reader is referred, we set.

Definition 6.63. A real analytic generic submanifold $M \subset \mathbb{C}^{n+m}$ is *minimal* at one of its points p if the fundamental pair of foliations of its complexification \mathcal{M} is covering at p (Definition 10.17).

Further informations may be found in Section 10. We conclude by formulating applications of Theorems 5.13 and 5.24.

Corollary 6.64. *The bound $\dim \mathfrak{hol}(M) \leq 8$ for a Levi nondegenerate hypersurface $M \subset \mathbb{C}^2$ is attained if and only if it is locally biholomorphic to the sphere $S^3 \subset \mathbb{C}^2$.*

Corollary 6.65. *The bound $\dim \mathfrak{hol}(M) \leq n^2 + 4n + 3$ for a Levi nondegenerate hypersurface $M \subset \mathbb{C}^{n+1}$ is attained if and only if it is locally biholomorphic to the sphere $S^{2n+1} \subset \mathbb{C}^{n+1}$.*

§7. EQUIVALENCE PROBLEMS AND NORMAL FORMS

7.1. Equivalences of submanifolds of solutions. As in §3.1, let (\mathcal{E}) and (\mathcal{E}') be two PDE systems and assume that φ transforms (\mathcal{E}) to (\mathcal{E}') . Defining A' similarly as A , it follows that

$$(7.2) \quad A'^{-1} \circ \Phi_{\mathcal{E}, \mathcal{E}'} \circ A(x, a, b) \equiv (\theta(x, a, b), f(a, b), g(a, b)) =: (x', a', b')$$

transforms F_v to F'_v , hence induces a map $(a, b) \mapsto (a', b')$. The arguments of Section 6 apply here with minor modifications to provide two fundamental lemmas.

Lemma 7.3. *Every equivalence $(x, y) \mapsto (x', y')$ between two PDE systems (\mathcal{E}) and (\mathcal{E}') comes with an associated transformation $(a, b) \mapsto (a', b')$ of the parameter spaces such that*

$$(7.4) \quad (x, y, a, b) \longmapsto (x', y', a', b')$$

is an equivalence between the associated submanifolds of solutions $\mathcal{M}_{(\mathcal{E})} \rightarrow \mathcal{M}'_{(\mathcal{E}')}$.

Conversely, let \mathcal{M} and \mathcal{M}' be two submanifolds of $\mathbb{K}_x^n \times \mathbb{K}_y^m \times \mathbb{K}_a^p \times \mathbb{K}_b^m$ and of $\mathbb{K}_{x'}^n \times \mathbb{K}_{y'}^m \times \mathbb{K}_{a'}^p \times \mathbb{K}_{b'}^m$ represented by $y = \Pi(x, a, b)$ and by $y' = \Pi'(x', a', b')$, in the same dimensions. Assume both are solvable with respect to the parameters.

Lemma 7.5. *Every equivalence*

$$(7.6) \quad (x, y, a, b) \longmapsto (\varphi(x, y), h(a, b))$$

between \mathcal{M} and \mathcal{M}' belonging to $G_{v,p}$ induces by projection the equivalence $(x, y) \mapsto \varphi(x, y)$ between the associated PDE systems $(\mathcal{E}_{\mathcal{M}})$ and $(\mathcal{E}'_{\mathcal{M}'})$.

7.7. Classification problems. Consequently, classifying PDE systems under point transformations (Section 3) is equivalent to the following.

Equivalence problem 7.8. Find an algorithm to decide whether two given submanifolds (of solutions) \mathcal{M} and \mathcal{M}' are equivalent through an element of $G_{v,p}$.

Classification problem 7.9. Classify submanifolds (of solutions) \mathcal{M} , namely provide a complete list of all possible such equations, including their automorphism group $\text{Aut}_{v,p}(\mathcal{M}) \subset G_{v,p}$.

7.10. Partial normal forms. Both problems above are of high complexity. At least as a preliminary step, it is useful to try to simplify somehow the defining equations of \mathcal{M} , by appropriate changes of coordinates belonging to $G_{v,p}$. To begin with, the next lemma holds for \mathcal{M} defined by $y = \Pi(x, a, b)$ with the only assumption that $b \mapsto \Pi(0, 0, b)$ has rank m at $b = 0$.

Lemma 7.11. ([CM1974, BER1999, Me2005a], [*]) *In coordinates $x' = (x'^1, \dots, x'^n)$ and $y' = (y'^1, \dots, y'^m)$ an arbitrary submanifold \mathcal{M}' defined by $y' = \Pi'(x', a', b')$ or dually by $b' = \Pi'^*(a', x', y')$ is equivalent to*

$$(7.12) \quad y = \Pi(x, a, b) \quad \text{or dually to} \quad b = \Pi^*(a, x, y)$$

with

$$(7.13) \quad \Pi(0, a, b) \equiv \Pi(x, 0, b) \equiv b \quad \text{or dually} \quad \Pi^*(0, x, y) \equiv \Pi^*(a, 0, y) \equiv y,$$

namely $\Pi = b + O(xa)$ and $\Pi^ = y + O(ax)$.*

Proof. We developpe

$$(7.14) \quad y' = \Pi'(0, a', b') + \Lambda'(x') + O(x'a').$$

Since $b' \mapsto \Pi'(0, a', b')$ has rank m at $b' = 0$, the coordinate change

$$(7.15) \quad b'' := \Pi'(0, a', b'), \quad a'' := a', \quad x'' := x', \quad y'' := y',$$

transforms \mathcal{M}' to \mathcal{M}'' defined by

$$(7.16) \quad y'' = \Pi''(x'', a'', b'') := b'' + \Lambda'(x'') + O(x''a'').$$

Solving b'' by means of the implicit function theorem, we get

$$(7.17) \quad b'' = \Pi''^*(a'', x'', y'') = y'' - \Lambda'(x'') + O(a''x''),$$

and it suffices to set $y := y'' - \Lambda'(x'')$, $x := x''$ and $a := a''$, $b := b''$. \square

Taking account of solvability with respect to the parameters, finer normalizations holds.

Lemma 7.18. *With $n = m = \kappa = 1$, every submanifold of solutions $y' = b' + x'a' [1 + O_1]$ of $y'_{x'x'} = F'(x', y', y'_{x'})$ is equivalent to*

$$(7.19) \quad y_{xx} = b + xa + O(x^2a^2).$$

Proof. Writing $y' = b' + x'[a' + a' \Lambda'(a', b') + O(x'a')]$, where $\Lambda' = O_1$, we set $a'' := a' + a' \Lambda'(a', b')$, $b'' := b'$, $x'' := x'$, $y'' := y'$, whence $y'' = b'' + x''[a'' + O(x''a'')]$. Dually $b'' = y'' - a''[x'' + x''x'' \Lambda''(x'', y'') + O(x''x''a'')]$, so we set $x := x'' + x''x'' \Lambda''(x'', y'')$, $y := y''$, $a := a''$, $b := b''$. \square

Corollary 7.20. *Every second order ordinary differential equation $y'_{x'x'} = F'(x', y', y'_{x'})$ is equivalent to*

$$(7.21) \quad y_{xx} = (y_x)^2 R(x, y, y_x).$$

7.22. Complete normal forms. The Moser theory of normal forms may be transferred with minor modifications to submanifolds of solutions associated to (\mathcal{E}_1) and to (\mathcal{E}_2) .

Theorem 7.23. ([CM1974, Ja1990], [*]) *A local \mathbb{K} -analytic submanifold of solutions associated to (\mathcal{E}_1) :*

$$(7.24) \quad y' = b' + x'a' + O_3 = \sum_{k' \geq 0} \sum_{l' \geq 0} \Pi'_{k', l'}(b') x'^{k'} a'^{l'}$$

can be mapped, by a transformation $(x', y', a', b') \mapsto (x, y, a, b)$ belonging to $\mathbb{G}_{v, p}$, to a submanifold of solutions of the specific form

$$(7.25) \quad y = b + xa + \Pi_{2,4}(b) x^2 a^4 + \Pi_{4,2}(b) a^2 x^4 + \sum_{k \geq 2} \sum_{l \geq 2} \sum_{k+l \geq 7} \Pi_{k,l}(b) x^k a^l.$$

Solving (a, b) from $y = \Pi$ and $y_x = \Pi_x$ with Π as above, we deduce the following.

Corollary 7.26. *Every $y'_{x'x'} = F'(x', y', y'_{x'})$ is equivalent to*

$$(7.27) \quad y_{xx} = (y_x)^2 [x^2 F_{2,2}(y) + x^3 r(x, y)] + (y_x)^4 [F_{0,4}(y) + x r(x, y)] + \sum_{k \geq 0} \sum_{l \geq 0} \sum_{k+l \geq 5} F_{k,l}(y) x^k (y_x)^l.$$

For the completely integrable system (\mathcal{E}_2) having several dependent variables (x^1, \dots, x^n) , $n \geq 2$, we have the following.

Theorem 7.28. ([CM1974], [*]) *A local \mathbb{K} -analytic submanifold of solutions associated to (\mathcal{E}_2) :*

$$(7.29) \quad y' = b' + \sum_{1 \leq k \leq n} x'^k a'^k + O_3$$

can be mapped, by a transformation $(x', y', a', b') \mapsto (x, y, a, b)$ belonging to $G_{v,p}$, to a submanifold of solutions of the specific form:

$$(7.30) \quad y = b + \sum_{1 \leq k \leq n} x^k a^k + \sum_{k \geq 2} \sum_{l \geq 2} \Pi_{k,l}(x, a, b)$$

where

$$(7.31) \quad \Pi_{k,l}(x, a, b) := \sum_{k_1 + \dots + k_n = k} \sum_{l_1 + \dots + l_n = l} \Pi_{k_1, \dots, k_n, l_1, \dots, l_n}(b) (x^1)^{k_1} \dots (x^n)^{k_n} (a^1)^{l_1} \dots (a^n)^{l_n}$$

with the terms $\Pi_{2,2}$, $\Pi_{2,3}$ and $\Pi_{3,3}$ satisfying:

$$(7.32) \quad 0 = \Delta \Pi_{2,2} = \Delta \Delta \Pi_{2,3} = \Delta \Delta \Pi_{3,2} = \Delta \Delta \Delta \Pi_{3,3},$$

where

$$(7.33) \quad \Delta := \sum_{1 \leq k \leq n} \frac{\partial^2}{\partial x^k \partial a^k}.$$

Exercise: solving (a^k, b) from $y = \Pi$ and $y_{x^l} = \Pi_{x^l}$, with Π as above, deduce a complete normal form for (\mathcal{E}_2) .

Open problem 7.34. *Find complete normal forms for submanifolds of solutions associated to (\mathcal{E}_4) and to (\mathcal{E}_5) .*

§8. STUDY OF TWO SPECIFIC EXAMPLES

8.1. Study of the Lie symmetries of (\mathcal{E}_4) . Its submanifold of solutions possesses two equations:

$$(8.2) \quad y^1 = \Pi^1(x, a, b^1, b^2) \quad y^2 = \Pi^2(x, a, b^1, b^2).$$

For instance, a generic submanifold $M \subset \mathbb{C}^3$ of CR dimension 1 and of codimension 3 has equations of such a form.

Assuming $\mathcal{V}_S(\mathcal{E}_4)$ to be twin solvable and having covering submanifold of solutions (see Definition 10.17), it may be verified (for $M \subset \mathbb{C}^3$, see [Be1997]) that at a Zariski-generic point, its equations are of the form:

$$(8.3) \quad \begin{aligned} y^1 &= b^1 + xa + O(x^2) + O(b^1) + O(b^2), \\ y^2 &= b^2 + xa(x+a) + O(x^3) + O(b^1) + O(b^2). \end{aligned}$$

The model has zero remainders with associated system

$$(8.4) \quad y_1^2 = 2x y_1^1 + (y_1^1)^2, \quad y_2^1 = 0,$$

the third equation $y_2^2 = 2y_1^1$ being obtained by differentiating the first.

We may put the submanifold in partial normal form. Proceeding as in [BES2005], some partial normalizations belonging to $G_{v,p}$ yield:

$$(8.5) \quad \begin{aligned} y^1 &= b^1 + ax + a^2 [\Pi_{3,2}^1(b) x^3 + \Pi_{4,2}^1(b) x^4 + \dots] + O(a^3 x^2), \\ y^2 &= b^2 + a [x^2 + \Pi_{4,1}^2(b) x^4 + \dots] + a^2 [x + \Pi_{3,2}^2(b) x^3 + \dots] + O(a^3 x^2). \end{aligned}$$

Redifferentiating, we get an appropriate, partially normalized system (\mathcal{E}_4):

$$(8.6) \quad \begin{cases} y_1^2 = y_1^1(2x + \mathbf{g}^1) + (y_1^1)^2(1 + \mathbf{g}^2) + (y_1^1)^3 \mathbf{s} + (y_1^1)^4 \mathbf{s} + (y_1^1)^5 \mathbf{s} + (y_1^1)^6 \mathbf{R}, \\ y_2^1 = (y_1^1)^2 \mathbf{h} + (y_1^1)^3 \mathbf{R}, \\ y_2^2 = y_1^1(2 + \mathbf{g}_x^1) + (y_1^1)^2(\mathbf{g}_x^2 + (2x + \mathbf{g}^1)\mathbf{h}) + (y_1^1)^3 \mathbf{r} + (y_1^1)^4 \mathbf{r} + (y_1^1)^5 \mathbf{r} + (y_1^1)^6 \mathbf{R}, \end{cases}$$

where, precisely:

- \mathbf{g}^1 , \mathbf{g}^2 and \mathbf{h} are functions of (x, y^1, y^2) satisfying $\mathbf{g}^j = \mathcal{O}(xx) + \mathcal{O}(y^1) + \mathcal{O}(y^2)$, $j = 1, 2$ and $\mathbf{h} = \mathcal{O}(x) + \mathcal{O}(y^1) + \mathcal{O}(y^2)$;
- \mathbf{r} and \mathbf{s} are unspecified functions, varying in the context, of (x, y^1, y^2) with $\mathbf{s} = \mathcal{O}(x) + \mathcal{O}(y^1) + \mathcal{O}(y^2)$, but possibly $\mathbf{r}(0) \neq 0$;
- \mathbf{R} is a remainder function of all the variables (x, y^1, y^2, y_1^1) parametrizing $\Delta_{\mathcal{E}_4}$.

Letting $\mathcal{L} = \mathcal{X} \frac{\partial}{\partial x} + \mathcal{Y}^1 \frac{\partial}{\partial y^1} + \mathcal{Y}^2 \frac{\partial}{\partial y^2}$ be a candidate infinitesimal Lie symmetry and applying $\mathcal{L}^{(2)} = \mathcal{L} + \mathbf{Y}_1^1 \frac{\partial}{\partial y_1^1} + \mathbf{Y}_1^2 \frac{\partial}{\partial y_1^2} + \mathbf{Y}_2^1 \frac{\partial}{\partial y_2^1} + \mathbf{Y}_2^2 \frac{\partial}{\partial y_2^2}$ to $\Delta_{\mathcal{E}_4}$, we obtain firstly, computing $\text{mod}(y_1^1)^5$:

$$(8.7) \quad \begin{aligned} 0 \equiv & -\mathbf{Y}_1^2 + [\mathcal{X}](y_1^1(2 + \mathbf{g}_x^1) + (y_1^1)^2 \mathbf{g}_x^2 + (y_1^1)^3 \mathbf{r} + (y_1^1)^4 \mathbf{r}) + \\ & + [\mathcal{Y}^1](y_1^1 \mathbf{r} + (y_1^1)^2 \mathbf{r} + (y_1^1)^3 \mathbf{r} + (y_1^1)^4 \mathbf{r}) + \\ & + [\mathcal{Y}^2](y_1^1 \mathbf{r} + (y_1^1)^2 \mathbf{r} + (y_1^1)^3 \mathbf{r} + (y_1^1)^4 \mathbf{r}) + \\ & + \mathbf{Y}_1^1(2x + \mathbf{g}^1 + y_1^1(2 + 2\mathbf{g}^2) + (y_1^1)^2 \mathbf{s} + (y_1^1)^3 \mathbf{s} + (y_1^1)^4 \mathbf{s}), \end{aligned}$$

and secondly, computing $\text{mod}(y_1^1)^2$:

$$(8.8) \quad 0 \equiv -\mathbf{Y}_2^1 + 2y_1^1 \mathbf{Y}_1^1 h.$$

The third Lie equation involving \mathbf{Y}_2^2 will be superfluous. Specializing (4.6)(II) to $m = 2$, we get \mathbf{Y}_1^1 and \mathbf{Y}_1^2 :

$$(8.9) \quad \begin{aligned} \mathbf{Y}_1^1 &= \mathcal{Y}_x^1 + [\mathcal{Y}_{y^1}^1 - \mathcal{X}_x] y_1^1 + [\mathcal{Y}_{y^2}^1] y_1^2 + [-\mathcal{X}_{y^1}] (y_1^1)^2 + [-\mathcal{X}_{y^2}] y_1^1 y_1^2 \\ \mathbf{Y}_1^2 &= \mathcal{Y}_x^2 + [\mathcal{Y}_{y^1}^2] y_1^1 + [\mathcal{Y}_{y^2}^2 - \mathcal{X}_x] y_1^2 + [-\mathcal{X}_{y^1}] y_1^2 y_1^1 + [-\mathcal{X}_{y^2}] (y_1^1)^2. \end{aligned}$$

and also \mathbf{Y}_2^1 and \mathbf{Y}_2^2 (in fact superfluous):

$$(8.10) \quad \begin{aligned} \mathbf{Y}_2^1 &= \mathcal{Y}_{xx}^1 + [2\mathcal{Y}_{xy^1}^1 - \mathcal{X}_{xx}] y_1^1 + [2\mathcal{Y}_{xy^2}^1] y_1^2 + [\mathcal{Y}_{y^1 y^1}^1 - 2\mathcal{X}_{xy^1}] (y_1^1)^2 + \\ &+ [2\mathcal{Y}_{y^1 y^2}^1 - 2\mathcal{X}_{xy^2}] y_1^1 y_1^2 + [\mathcal{Y}_{y^2 y^2}^1] (y_1^1)^2 + [-\mathcal{X}_{y^1 y^1}] (y_1^1)^3 + \\ &+ [-2\mathcal{X}_{y^1 y^2}] (y_1^1)^2 y_1^2 + [-\mathcal{X}_{y^2 y^2}] y_1^1 (y_1^1)^2 + [\mathcal{Y}_{y^1}^1 - 2\mathcal{X}_x] y_2^1 + \\ &+ [\mathcal{Y}_{y^2}^1] y_2^2 + [-3\mathcal{X}_{y^1}] y_1^1 y_2^1 + [-\mathcal{X}_{y^2}] y_1^1 y_2^2 + [-2\mathcal{X}_{y^2}] y_1^2 y_2^1, \\ \mathbf{Y}_2^2 &= \mathcal{Y}_{xx}^2 + [2\mathcal{Y}_{xy^1}^2] y_1^1 + [2\mathcal{Y}_{xy^2}^2 - \mathcal{X}_{xx}] y_1^2 + [\mathcal{Y}_{y^1 y^1}^2] (y_1^1)^2 + \\ &+ [2\mathcal{Y}_{y^1 y^2}^2 - 2\mathcal{X}_{xy^1}] y_1^1 y_1^2 + [\mathcal{Y}_{y^2 y^2}^2 - 2\mathcal{X}_{xy^2}] (y_1^1)^2 + [-\mathcal{X}_{y^1 y^1}] (y_1^1)^2 y_1^2 + \\ &+ [-2\mathcal{X}_{y^1 y^2}] y_1^1 (y_1^1)^2 + [-\mathcal{X}_{y^2 y^2}] (y_1^1)^3 + [\mathcal{Y}_{y^1}^2] y_2^1 + \\ &+ [\mathcal{Y}_{y^2}^2 - 2\mathcal{X}_x] y_2^2 + [-2\mathcal{X}_{y^1}] y_1^1 y_2^2 + [-\mathcal{X}_{y^1}] y_1^1 y_2^1 + [-3\mathcal{X}_{y^2}] y_1^2 y_2^2. \end{aligned}$$

Inserting Y_1^2 and Y_1^1 in the first Lie equation (8.7) in which y_1^2 is replaced by the value (8.6)₁ it has on $\Delta_{\mathcal{E}_4}$ and still computing mod $(y_1^1)^5$, we get, again with r, s being unspecified functions of (x, y^1, y^2) with $s(0) = 0$:

(8.11)

$$\begin{aligned}
0 \equiv & -\mathcal{Y}_x^2 + [-\mathcal{Y}_{y^1}^2] y_1^1 + \\
& + [-\mathcal{Y}_{y^2}^2 + \mathcal{X}_x] (y_1^1(2x + \mathbf{g}^1) + (y_1^1)^2(1 + \mathbf{g}^2) + (y_1^1)^3 s + (y_1^1)^4 s) + \\
& + [\mathcal{X}_{y^1}] ((y_1^1)^2(2x + \mathbf{g}^1) + (y_1^1)^3(1 + \mathbf{g}^2) + (y_1^1)^4 s) + \\
& + [\mathcal{X}_{y^2}] ((y_1^1)^2[2x + \mathbf{g}^1]^2 + (y_1^1)^3(4x + 2\mathbf{g}^1)(1 + \mathbf{g}^2) + (y_1^1)^4(1 + s)) + \\
& + [\mathcal{X}] (y_1^1(2 + \mathbf{g}_x^1) + (y_1^1)^2 \mathbf{g}_x^2 + (y_1^1)^3 r + (y_1^1)^4 r) + \\
& + [\mathcal{Y}^1] (y_1^1 r + (y_1^1)^2 r + (y_1^1)^3 r + (y_1^1)^4 r) + \\
& + [\mathcal{Y}^2] (y_1^1 r + (y_1^1)^2 r + (y_1^1)^3 r + (y_1^1)^4 r) + \\
& + [\mathcal{Y}_x^1] (2x + \mathbf{g}^1 + y_1^1(2 + 2\mathbf{g}^2) + (y_1^1)^2 s + (y_1^1)^3 s + (y_1^1)^4 s) + \\
& + [\mathcal{Y}_{y^1}^1 - \mathcal{X}_x] (y_1^1(2x + \mathbf{g}^1) + (y_1^1)^2(2 + 2\mathbf{g}^2) + (y_1^1)^3 s + (y_1^1)^4 s) + \\
& + [\mathcal{Y}_{y^2}^1] (y_1^1[2x + \mathbf{g}^1]^2 + (y_1^1)^2(2x + \mathbf{g}^1)(3 + 3\mathbf{g}^2) + (y_1^1)^3(2 + s) + (y_1^1)^4 s) + \\
& + [-\mathcal{X}_{y^1}] ((y_1^1)^2(2x + \mathbf{g}^1) + (y_1^1)^3(2 + 2\mathbf{g}^2) + (y_1^1)^4 s) + \\
& + [-\mathcal{X}_{y^2}] ((y_1^1)^2[2x + \mathbf{g}^1]^2 + (y_1^1)^3(2x + \mathbf{g}^1)(3 + 3\mathbf{g}^2) + (y_1^1)^4(2 + s)).
\end{aligned}$$

Collecting the coefficients of the monomials cst., y_1^1 , $(y_1^1)^2$, $(y_1^1)^3$, $(y_1^1)^4$, we get, after slight simplification (in the coefficient of $(y_1^1)^2$, the term $(2x + \mathbf{g}^1)\mathcal{X}_x$ annihilates with its opposite; in the coefficient of $(y_1^1)^3$, two pairs annihilate and then, we divide by $[1 + \mathbf{g}^2]$) a system of five linear PDE's:

$$\begin{aligned}
(8.12) \quad & 0 = -\mathcal{Y}_x^2 + (2x + \mathbf{g}^1)\mathcal{Y}_x^1, \\
& 0 = -\mathcal{Y}_{y^1}^2 - (2x + \mathbf{g}^1)\mathcal{Y}_{y^2}^2 + (2 + \mathbf{g}_x^1)\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + \\
& \quad + (2 + 2\mathbf{g}^2)\mathcal{Y}_x^1 + (2x + \mathbf{g}^1)\mathcal{Y}_{y^1}^1 + [2x + \mathbf{g}^1]^2\mathcal{Y}_{y^2}^1, \\
& 0 = -\mathcal{Y}_{y^2}^2 + \mathcal{X}_x + \mathbf{g}_x^2[1 + \mathbf{g}^2]^{-1}\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + \\
& \quad + s\mathcal{Y}_x^1 + 2\mathcal{Y}_{y^1}^1 - 2\mathcal{X}_x + (6x + 3\mathbf{g}^2)\mathcal{Y}_{y^2}^1, \\
& 0 = s\mathcal{Y}_{y^2}^2 + s\mathcal{X}_x + (1 + \mathbf{g}^2)\mathcal{X}_{y^1} + (2x + \mathbf{g}^1)(2 + 2\mathbf{g}^2)\mathcal{X}_{y^2} + \\
& \quad + r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + s\mathcal{Y}_x^1 + s\mathcal{Y}_{y^1}^1 + s\mathcal{X}_x + (2 + s)\mathcal{Y}_{y^2}^1 - \\
& \quad - (2 + 2\mathbf{g}^2)\mathcal{X}_{y^1} - (2x + \mathbf{g}^1)(3 + 3\mathbf{g}^2)\mathcal{X}_{y^2}, \\
& 0 = s\mathcal{Y}_{y^2}^2 + s\mathcal{X}_x + s\mathcal{X}_{y^1} + (1 + s)\mathcal{X}_{y^2} + r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + s\mathcal{Y}_x^1 + s\mathcal{Y}_{y^1}^1 + \\
& \quad + s\mathcal{X}_x + s\mathcal{Y}_{y^2}^1 + s\mathcal{X}_{y^1} - (2 + s)\mathcal{X}_{y^2}.
\end{aligned}$$

We then simplify the remainders using $s + s = s$, $r + s = r$ and $r + r = r$; we divide (8.12)₅ by $(1 + s)$; we replace \mathcal{X}_{y^2} obtained from (8.12)₅ in (8.12)₄; we divide (8.12)₄ by

$(1 + \mathbf{g}^2)$; we then solve \mathcal{X}_{y^1} from $(8.12)_4$ and finally we insert it in $(8.12)_5$; we get:

$$\begin{aligned}
(8.13) \quad & 0 = -\mathcal{Y}_x^2 + (2x + \mathbf{g}^1)\mathcal{Y}_x^1, \\
& 0 = -\mathcal{Y}_{y^1}^2 - (2x + \mathbf{g}^1)\mathcal{Y}_{y^2}^2 + (2 + \mathbf{g}_x^1)\mathcal{X} + (2 + 2\mathbf{g}^2)\mathcal{Y}_x^1 + (2x + \mathbf{g}^1)\mathcal{Y}_{y^1}^1 + \\
& \quad + [2x + \mathbf{g}^1]^2\mathcal{Y}_{y^2}^1 + r\mathcal{Y}^1 + r\mathcal{Y}^2, \\
& 0 = -\mathcal{Y}_{y^2}^2 - \mathcal{X}_x + 2\mathcal{Y}_{y^1}^1 + (6x + 3\mathbf{g}^2)\mathcal{Y}_{y^2}^1 + r\mathcal{Y}^1 + r\mathcal{Y}^2 + s\mathcal{Y}_x^1 + \\
& \quad + \mathbf{g}_x^2[1 + \mathbf{g}^2]^{-1}\mathcal{X}, \\
& 0 = -\mathcal{X}_{y^1} + (2 + s)\mathcal{Y}_{y^2}^1 + r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + s\mathcal{X}_x + s\mathcal{Y}_x^1 + s\mathcal{Y}_{y^1}^1 + s\mathcal{Y}_{y^2}^2, \\
& 0 = -\mathcal{X}_{y^2} + r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + s\mathcal{X}_x + s\mathcal{Y}_x^1 + s\mathcal{Y}_{y^1}^1 + s\mathcal{Y}_{y^2}^2 + s\mathcal{Y}_{y^2}^2.
\end{aligned}$$

Similarly, developing the second equation (8.8) and computing mod $(y_1^1)^2$, we get:

$$(8.14) \quad 0 \equiv -\mathcal{Y}_{xx}^1 + [-2\mathcal{Y}_{xy^1}^1 + \mathcal{X}_{xx}]y_1^1 + [-(4x + 2\mathbf{g}^1)\mathcal{Y}_{xy^2} - (2 + \mathbf{h})\mathcal{Y}_{y^2}^1] + \\
+ [2\mathbf{h}\mathcal{Y}_x^1]y_1^1.$$

Collecting the coefficients of the monomials cst., y_1^1 , we get two more linear PDE's:

$$(8.15) \quad \begin{aligned}
0 &= -\mathcal{Y}_{xx}^1, \\
0 &= -2\mathcal{Y}_{xy^1}^1 + \mathcal{X}_{xx} - (4x + 2\mathbf{g}^1)\mathcal{Y}_{xy^2}^1 - (2 + \mathbf{h})\mathcal{Y}_{y^2}^1 + 2\mathbf{h}\mathcal{Y}_x^1.
\end{aligned}$$

Proposition 8.16. *Setting as initial conditions the five specific differential coefficients*

$$(8.17) \quad \mathbf{P} := \mathbf{P}(\mathcal{X}, \mathcal{Y}^1, \mathcal{Y}^2, \mathcal{Y}_x^1, \mathcal{X}_x) = r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + r\mathcal{Y}_x^1 + r\mathcal{X}_x,$$

it follows by cross differentiations and by linear substitutions from the seven equations $(8.13)_i$, $i = 1, 2, 3, 4, 5$, $(8.15)_j$, $j = 1, 2$, that \mathcal{X}_{y^1} , \mathcal{X}_{y^2} , $\mathcal{Y}_{y^1}^1$, $\mathcal{Y}_{y^2}^1$, \mathcal{Y}_x^2 , $\mathcal{Y}_{y^1}^2$, $\mathcal{Y}_{y^2}^2$ and \mathcal{X}_{xx} , \mathcal{X}_{xy^1} , \mathcal{X}_{xy^2} , \mathcal{Y}_{xx}^1 , $\mathcal{Y}_{xy^1}^1$, $\mathcal{Y}_{xy^2}^1$ are uniquely determined as linear combinations of $(\mathcal{X}, \mathcal{Y}^1, \mathcal{Y}^2, \mathcal{Y}_x^1, \mathcal{X}_x)$, namely:

$$(8.18) \quad \left\{ \begin{array}{lllll}
& \mathcal{Y}_x^2 \stackrel{1}{=} \mathbf{P}, & \mathcal{X}_{xx} \stackrel{2}{=} \mathbf{P}, & \mathcal{Y}_{xx}^1 \stackrel{3}{=} \mathbf{P}, & \\
\mathcal{X}_{y^1} \stackrel{4}{=} \mathbf{P}, & \mathcal{Y}_{y^1}^1 \stackrel{5}{=} \mathbf{P}, & \mathcal{Y}_{y^1}^2 \stackrel{6}{=} \mathbf{P}, & \mathcal{X}_{xy^1} \stackrel{7}{=} \mathbf{P}, & \mathcal{Y}_{xy^1}^1 \stackrel{8}{=} \mathbf{P}, \\
\mathcal{X}_{y^2} \stackrel{9}{=} \mathbf{P}, & \mathcal{Y}_{y^2}^1 \stackrel{10}{=} \mathbf{P}, & \mathcal{Y}_{y^2}^2 \stackrel{11}{=} \mathbf{P}, & \mathcal{X}_{xy^2} \stackrel{12}{=} \mathbf{P}, & \mathcal{Y}_{xy^2}^1 \stackrel{13}{=} \mathbf{P}.
\end{array} \right.$$

Then the expressions \mathbf{P} are stable under differentiation:

$$(8.19) \quad \begin{aligned}
\mathbf{P}_x &= \mathbf{P} + r\mathcal{Y}_x^2 + r\mathcal{Y}_{xx}^1 + r\mathcal{X}_{xx} = \mathbf{P}, \\
\mathbf{P}_{y^1} &= \mathbf{P} + r\mathcal{X}_{y^1} + r\mathcal{Y}_{y^1}^1 + r\mathcal{Y}_{y^1}^2 + r\mathcal{Y}_{xy^1}^1 + r\mathcal{X}_{xy^1} = \mathbf{P}, \\
\mathbf{P}_{y^2} &= \mathbf{P} + r\mathcal{X}_{y^2} + r\mathcal{Y}_{y^2}^1 + r\mathcal{Y}_{y^2}^2 + r\mathcal{Y}_{xy^2}^1 + r\mathcal{X}_{xy^2} = \mathbf{P},
\end{aligned}$$

and moreover, all other, higher order partial derivatives of \mathcal{X} , of \mathcal{Y}^1 and of \mathcal{Y}^2 may be expressed as $\mathbf{P}(\mathcal{X}, \mathcal{Y}^1, \mathcal{Y}^2, \mathcal{Y}_x^1, \mathcal{X}_x)$.

Corollary 8.20. *An infinitesimal Lie symmetry of (\mathcal{E}_4) is uniquely determined by the five initial Taylor coefficients*

$$(8.21) \quad \mathcal{X}(0), \mathcal{Y}^1(0), \mathcal{Y}^2(0), \mathcal{Y}_x^1(0), \mathcal{X}_x(0).$$

Proof of the proposition. We notice that $(8.18)_1$ and $(8.18)_3$ are given for free by $(8.13)_1$ and by $(8.15)_1$. Differentiating $(8.13)_3$ with respect to x , we get:

$$(8.22) \quad 0 = -\mathcal{Y}_{xy^2} - \mathcal{X}_{xx} + 2\mathcal{Y}_{xy^1}^1 + (6 + 3\mathbf{g}_x^1)\mathcal{Y}_{y^2}^1 + (6x + 3\mathbf{g}^1)\mathcal{Y}_{xy^2}^1 + r\mathcal{Y}^1 + \\
+ r\mathcal{Y}_x^1 + r\mathcal{Y}^2 + r\mathcal{Y}_x^2 + r\mathcal{Y}_x^1 + s\mathcal{Y}_{xx}^1 + r\mathcal{X} + \mathbf{g}_x^2[1 + \mathbf{g}^2]^{-1}\mathcal{X}.$$

By (8.15)₁, $s\mathcal{Y}_{xx}^1$ vanishes. We replace \mathcal{Y}_x^2 thanks to (8.13)₁. Differentiating (8.13)₁ with respect to y^2 , we may subtract $0 = -\mathcal{Y}_{xy^2} + (2x + \mathbf{g}^1)\mathcal{Y}_{xy^2}^1 + r\mathcal{Y}_x^1$. We get:

$$(8.23) \quad \begin{aligned} 0 = & -\mathcal{X}_{xx} + 2\mathcal{Y}_{xy^1}^1 + (4x + 2\mathbf{g}^1)\mathcal{Y}_{xy^2}^1 + \\ & + (6 + 3\mathbf{g}_x^1)\mathcal{Y}_{y^2}^1 + r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + r\mathcal{Y}_x^1 + \mathbf{g}_x^2[1 + \mathbf{g}^2]^{-1}\mathcal{X}_x. \end{aligned}$$

By means of (8.15)₂, we replace the first three terms and then solve $\mathcal{Y}_{y^2}^1$:

$$(8.24) \quad \mathcal{Y}_{y^2}^1 = r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + r\mathcal{Y}_x^1 + \mathbf{k}^*\mathcal{X}_x,$$

introducing a notation for a new function that should be recorded:

$$(8.25) \quad \mathbf{k}^* := \mathbf{g}_x^2[1 + \mathbf{g}^2]^{-1}[4 + 3\mathbf{g}_x^1 - \mathbf{h}]^{-1}.$$

This is (8.18)₁₀. Next, we differentiate the obtained equation with respect to x , getting:

$$(8.26) \quad \mathcal{Y}_{xy^2}^1 = r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + r\mathcal{Y}_x^1 + r\mathcal{X}_x + \mathbf{k}^*\mathcal{X}_{xx}.$$

This is (8.18)₁₃. We replace the obtained value of $\mathcal{Y}_{y^2}^1$ in (8.13)₂, (8.13)₃, (8.15)₂ and the obtained value of $\mathcal{Y}_{xy^2}^1$ in (8.15)₂. This yields a new, simpler system of seven equations:

$$(8.27) \quad \begin{aligned} 0 = & -\mathcal{Y}_x^2 + (2x + \mathbf{g}^1)\mathcal{Y}_x^1, \\ 0 = & -\mathcal{Y}_{y^1}^2 - (2x + \mathbf{g}^1)\mathcal{Y}_{y^2}^2 + (2 + \mathbf{g}_x^1)\mathcal{X} + (2 + 2\mathbf{g}^2)\mathcal{Y}_x^1 + (2x + \mathbf{g}^1)\mathcal{Y}_{y^1}^1 + \\ & + s\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + s\mathcal{Y}_x^1 + \mathbf{k}^*[2x + \mathbf{g}^1]^2\mathcal{X}_x, \\ 0 = & -\mathcal{Y}_{y^2}^2 - \mathcal{X}_x + 2\mathcal{Y}_{y^1}^1 + s\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + s\mathcal{Y}_x^1 + \mathbf{k}^*(6x + 3\mathbf{g}^1)\mathcal{X}_x, \\ 0 = & -\mathcal{X}_{y^1} + r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + s\mathcal{Y}_x^1 + s\mathcal{X}_x + s\mathcal{Y}_{y^1}^1 + s\mathcal{Y}_{y^2}^2, \\ 0 = & -\mathcal{X}_{y^2} + r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + s\mathcal{Y}_x^1 + s\mathcal{X}_x + s\mathcal{Y}_{y^1}^1 + s\mathcal{Y}_{y^2}^2, \\ 0 = & -\mathcal{Y}_{xx}^1, \\ 0 = & -2\mathcal{Y}_{xy^1}^1 + \mathcal{X}_{xx}(1 - \mathbf{k}^*(4x + 2\mathbf{g}^1)) + r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + r\mathcal{Y}_x^1 + s\mathcal{X}_x. \end{aligned}$$

Restarting from this system, we differentiate (8.27)₃ with respect to x :

$$(8.28) \quad \begin{aligned} 0 = & -\mathcal{Y}_{xy^2}^2 - \mathcal{X}_{xx} + 2\mathcal{Y}_{xy^1}^1 + r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + \\ & + r\mathcal{X}_x + r\mathcal{Y}_x^1 + r\mathcal{Y}_x^2 + s\mathcal{Y}_{xx}^1 + \mathbf{k}^*(6x + 3\mathbf{g}^1)\mathcal{X}_{xx}. \end{aligned}$$

We replace \mathcal{Y}_x^2 , we erase \mathcal{Y}_{xx}^1 and we add (8.27)₇:

$$(8.29) \quad 0 = -\mathcal{Y}_{xy^2}^2 + \mathbf{k}^*(2x + \mathbf{g}^1)\mathcal{X}_{xx} + r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + r\mathcal{Y}_x^1 + r\mathcal{X}_x.$$

We differentiate (8.27)₂ with respect to x :

$$(8.30) \quad \begin{aligned} 0 = & -\mathcal{Y}_{xy^1}^2 - (2 + \mathbf{g}_x^1)\mathcal{Y}_{y^2}^2 - (2x + \mathbf{g}^1)\mathcal{Y}_{xy^2}^2 + r\mathcal{X} + (2 + \mathbf{g}_x^1)\mathcal{X}_x + \\ & + s\mathcal{Y}_x^1 + (2 + 2\mathbf{g}^2)\mathcal{Y}_{xx}^1 + (2 + \mathbf{g}_x^1)\mathcal{Y}_{y^1}^1 + (2x + \mathbf{g}^1)\mathcal{Y}_{xy^1}^1 + r\mathcal{X}_x + \\ & + s\mathcal{X}_x + r\mathcal{Y}^1 + r\mathcal{Y}_x^1 + r\mathcal{Y}^2 + r\mathcal{Y}_x^2 + r\mathcal{Y}_x^1 + s\mathcal{Y}_{xx}^1 + \mathbf{k}^*[2x + \mathbf{g}^1]^2\mathcal{X}_{xx}. \end{aligned}$$

Differentiating (8.27)₁ with respect to y^1 , we may subtract $0 = -\mathcal{Y}_{xy^1}^2 + (2x + \mathbf{g}^1)\mathcal{Y}_{xy^1}^1 + r\mathcal{Y}_x^1$; we replace \mathcal{Y}_x^2 and erase \mathcal{Y}_{xx}^1 ; we subtract (8.29) multiplied by $(2x + \mathbf{g}^1)$; we get:

$$(8.31) \quad 0 = -\mathcal{Y}_{y^2}^2 + (1 + s)\mathcal{X}_x + \mathcal{Y}_{y^1}^1 + r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + r\mathcal{Y}_x^1.$$

Comparing with (8.27)₃ yields:

$$(8.32) \quad \begin{aligned} \mathcal{Y}_{y^1}^1 = & (2 + s)\mathcal{X}_x + r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + r\mathcal{Y}_x^1, \\ \mathcal{Y}_{y^2}^2 = & (3 + s)\mathcal{X}_x + r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + r\mathcal{Y}_x^1. \end{aligned}$$

These are (8.18)₅ and (8.18)₁₁. Differentiating these two equations with respect to x , replacing \mathcal{Y}_x^2 and erasing \mathcal{Y}_{xx}^1 , we get:

$$(8.33) \quad \begin{aligned} \mathcal{Y}_{xy^1}^1 &= (2 + s)\mathcal{X}_{xx} + r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + r\mathcal{Y}_x^1 + r\mathcal{X}_x, \\ \mathcal{Y}_{xy^2}^2 &= (3 + s)\mathcal{X}_{xx} + r\mathcal{X} + r\mathcal{Y}^1 + r\mathcal{Y}^2 + r\mathcal{Y}_x^1 + r\mathcal{X}_x. \end{aligned}$$

We then replace this value of $\mathcal{Y}_{xy^2}^2$ in (8.29) and solve \mathcal{X}_{xx} : this yields (8.18)₂.

To conclude, we replace \mathcal{X}_{xx} so obtained in (8.27)₇: this yields (8.18)₈. We replace $\mathcal{Y}_{y^1}^1$ and $\mathcal{Y}_{y^2}^2$ from (8.32) in (8.27)₄ and in (8.27)₅: this yields (8.18)₄ and this yields (8.18)₉. Thanks to (8.18)₂ (got) we observe that

$$(8.34) \quad P_x = P + r\mathcal{Y}_{xx}^1 + r\mathcal{Y}_x^2 + r\mathcal{X}_{xx} = P.$$

Differentiating (8.18)₄ (got) and (8.18)₉ (got) with respect to x then yields (8.18)₇ and (8.18)₁₂. We replace $\mathcal{Y}_{y^1}^1$ and $\mathcal{Y}_{y^2}^2$ from (8.18)₅ (got) and (8.18)₁₁ (got) in (8.27)₂: this yields (8.18)₆. Finally, to obtain the very last (8.18)₁₃, we differentiate (8.18)₁₀ (got) with respect to x .

The proof of Proposition 8.16 is complete. \square

We claim that the bound $\dim \mathfrak{G}\mathfrak{Y}\mathfrak{M}(\mathcal{E}_4) \leq 5$ is attained for the model (8.4). Indeed, with $0 = r = s$ and $0 = \mathfrak{g}^1 = \mathfrak{g}^2 = \mathfrak{h}$ (whence $\mathfrak{k}^* = 0$) (8.24) is $\mathcal{Y}_{y^2}^1 = 0$ and then the seven equations (8.27) are:

$$(8.35) \quad \begin{cases} 0 = -\mathcal{Y}_x^2 + 2x\mathcal{Y}_x^1, \\ 0 = -\mathcal{Y}_{y^1}^2 - 2x\mathcal{Y}_{y^2}^2 + 2\mathcal{X} + 2\mathcal{Y}_x^1 + 2x\mathcal{Y}_{y^1}^1, \\ 0 = -\mathcal{Y}_{y^2}^2 - \mathcal{X}_x + 2\mathcal{Y}_{y^1}^1, \\ 0 = -\mathcal{X}_{y^1}, \\ 0 = -\mathcal{X}_{y^2}, \\ 0 = -\mathcal{Y}_{xx}^1, \\ 0 = -2\mathcal{Y}_{xy^1}^1 + \mathcal{X}_{xx}, \end{cases}$$

having the general solution

$$(8.36) \quad \begin{cases} \mathcal{X} = a - d + ex, \\ \mathcal{Y}^1 = b + dx + 2ey^1, \\ \mathcal{Y}^2 = c + 2ay^1 + 3ey^2 + dxx. \end{cases}$$

depending on five parameters $a, b, c, d, e \in \mathbb{K}$. Five generators of $\mathfrak{G}\mathfrak{Y}\mathfrak{M}(\mathcal{E}_4)$ are:

$$(8.37) \quad \begin{cases} \mathcal{D} := x\partial_x + 2y^1\partial_{y^1} + 3y^2\partial_{y^2}, \\ \mathcal{L}_1 := -\partial_x + x\partial_{y^1} + xx\partial_{y^2}, \\ \mathcal{L}'_1 := \partial_x + 2y^1\partial_{y^2}, \\ \mathcal{L}_2 := \partial_{y^1}, \\ \mathcal{L}_3 := \partial_{y^2}. \end{cases}$$

The commutator table

\mathcal{D}	\mathcal{L}_1	\mathcal{L}'_1	\mathcal{L}_2	\mathcal{L}_3	
\mathcal{D}	0	$-\mathcal{L}_1$	$-\mathcal{L}'_1$	$-2\mathcal{L}_2$	$-3\mathcal{L}_3$
\mathcal{L}_1	\mathcal{L}_1	0	$-\mathcal{L}_2$	0	0
\mathcal{L}'_1	\mathcal{L}'_1	\mathcal{L}_2	0	$-2\mathcal{L}_3$	0
\mathcal{L}_2	$2\mathcal{L}_2$	0	$2\mathcal{L}_3$	0	0
\mathcal{L}_3	$3\mathcal{L}_3$	0	0	0	0

Table 3.

shows that the subalgebra spanned by $\mathcal{L}_1, \mathcal{L}'_1, \mathcal{L}_2, \mathcal{L}_3$ is isomorphic to the unique irreducible 4-dimensional nilpotent Lie algebra \mathfrak{n}_4^1 ([OV1994, BES2005]). Then $\mathfrak{S}\mathfrak{M}(\mathcal{E}_4)$ is a semidirect product of \mathbb{K} with \mathfrak{n}_4^1 . The author ignores whether it is rigid. The following accessible research will be pursued in a subsequent publication.

Open problem 8.38. *Classify systems (\mathcal{E}_4) up to point transformations. Deduce a complete classification, up to local biholomorphisms, of all real analytic generic submanifolds of codimension 2 in \mathbb{C}^3 , valid at a Zariski-generic point.*

8.39. Almost everywhere rigid hypersurfaces. When studying and classifying differential objects, it is essentially no restriction to assume their Lie symmetry groups to be of dimension ≥ 1 , the study of objects having no infinitesimal symmetries being an independent field of research. In particular, if $M \subset \mathbb{C}^{n+1}$ ($n \geq 1$) is a connected real analytic hypersurface, we may suppose that $\dim \mathfrak{hol}(M) \geq 1$, at least. So let \mathcal{L} be a nonzero holomorphic vector field with $\mathcal{L} + \bar{\mathcal{L}}$ tangent to M .

Lemma 8.40. ([Ca1932a, St1996, BER1999]) *If in addition M is finitely nondegenerate, then*

$$(8.41) \quad \Sigma := \{p \in M : \mathcal{L}(p) \in T_p^c M\}$$

is a proper real analytic subset of M .

In other words, at every point p belonging to the Zariski-dense subset $M \setminus \Sigma$, the real nonzero vector $\mathcal{L}(p) + \bar{\mathcal{L}}(p) \in T_p M$ supplements $T_p^c M$. Straightening \mathcal{L} in a neighborhood of p , there exist local coordinates $t = (z_1, \dots, z_n, w)$ with $T_0^c M = \{w = 0\}$, $T_0 M = \{\text{Im } w = 0\}$, whence M is given by $\text{Im } w = h(z, \bar{z}, \text{Re } w)$, and with $\mathcal{L} = \frac{\partial}{\partial w}$. The tangency of $\frac{\partial}{\partial w} + \frac{\partial}{\partial \bar{w}} = \frac{\partial}{\partial u}$ to M entails that h is independent of u . Then the complex equation of M is of the precise form

$$(8.42) \quad w = \bar{w} + i\bar{\Theta}(z, \bar{z}),$$

with $\bar{\Theta} = 2h$ simply. The reality of h reads $\bar{\Theta}(z, \bar{z}) \equiv \Theta(\bar{z}, z)$.

Definition 8.43. A real analytic hypersurface $M \subset \mathbb{C}^{n+1}$ is called *rigid* at one of its points p if there exists $\mathcal{L} \in \mathfrak{hol}(M)$ with

$$(8.44) \quad T_p M = T_p^c M \oplus \mathbb{R}(\mathcal{L}(p) + \bar{\mathcal{L}}(p)).$$

Similar elementary facts hold for general submanifolds of solutions.

Lemma 8.45. *With $n \geq 1$ and $m = 1$, let \mathcal{M} be a (connected) submanifold of solutions that is solvable with respect to the parameters. If there exists a nonzero $\mathcal{L} + \mathcal{L}^* \in \mathfrak{S}\mathfrak{M}(\mathcal{M})$, then at Zariski-generic points $p \in \mathcal{M}$, we have $\mathcal{L}(p) \notin F_v(p)$ and there exist local coordinates centered at p in which $\mathcal{L} = \frac{\partial}{\partial y}$, $\mathcal{L}^* = \frac{\partial}{\partial b}$, whence \mathcal{M} has equation of the form*

$$(8.46) \quad y = b + \Pi(x, a),$$

with Π independent of b .

The associated system $(\mathcal{E}_{\mathcal{M}})$ has then equations F_α that are all independent of y .

8.47. Study of the Lie symmetries of (\mathcal{E}_5) . In Example 1.28, it is thus essentially no restriction to assume the hypersurface $M \subset \mathbb{C}^3$ to be rigid.

Theorem 8.48. ([GM2003b, FK2005a, FK2005b]) *The model hypersurface M_0 of equation*

$$(8.49) \quad w = \bar{w} + i \frac{2 z^1 \bar{z}^1 + z^1 z^1 \bar{z}^2 + \bar{z}^1 \bar{z}^1 z^2}{1 - z^2 \bar{z}^2}$$

has transitive Lie symmetry algebra $\mathfrak{ho}(M_0)$ isomorphic to $\mathfrak{so}(3, 2)$ and is locally biholomorphic to a neighborhood of every geometrically smooth point of the tube

$$(8.50) \quad (\operatorname{Re} w')^2 = (\operatorname{Re} z'_1)^2 + (\operatorname{Re} z'_1)^3$$

over the standard cone of \mathbb{R}^3 . Both are Levi-degenerate with Levi form of rank 1 at every point and are 2-nondegenerate. The associated PDE system (\mathcal{E}_{M_0})

$$(8.51) \quad y_{x^2} = \frac{1}{4} (y_{x^1})^2, \quad y_{x^1 x^1 x^1} = 0$$

(plus other equations obtained by cross differentiation) has infinitesimal Lie symmetry algebra isomorphic to $\mathfrak{so}(5, \mathbb{C})$, the complexification $\mathfrak{so}(3, 2) \otimes \mathbb{C}$.

Through tentative issues ([Eb2006, GM2006]), it has been suspected that M_0 is the right model in the category of real analytic hypersurfaces $M \subset \mathbb{C}^3$ having Levi form of rank 1 that are 2-nondegenerate everywhere. Based on the rigidity of the simple Lie algebra $\mathfrak{so}(5, \mathbb{C})$ (Theorem 5.15), Theorem 8.105 below will confirm this expectation.

8.52. Preparation. Thus, translating the considerations to the PDE language, with $n = 2$ and $m = 1$, consider a submanifold of solutions of the form

$$(8.53) \quad \begin{aligned} y &= b + \Pi(x, a) \\ &= b + \frac{2 x^1 a^1 + x^1 x^1 a^2 + a^1 a^1 x^2}{1 - x^2 a^2} + O_4, \end{aligned}$$

where O_4 is a function of (x, a) only. The term $2 x^1 a^1$ corresponds to a Levi form of rank ≥ 1 at every point. The term $x^1 x^1 a^2$ guarantees solvability with respect to the parameters (compare Definition 2.12). Let us developpe

$$(8.54) \quad \Pi(x, a) = \sum_{k_1, k_2 \geq 0} \sum_{l_1, l_2 \geq 0} \Pi_{k_1, k_2, l_1, l_2} (x^1)^{k_1} (x^2)^{k_2} (a^1)^{l_1} (a^2)^{l_2},$$

with $\Pi_{k_1, k_2, l_1, l_2} \in \mathbb{K}$. Of course, $\Pi_{1,0,1,0} = 2$, $\Pi_{2,0,0,1} = 1$ and $\Pi_{0,1,2,0} = 1$.

Lemma 8.55. *A transformation belonging to $G_{v,p}$ insures*

$$(8.56) \quad \begin{aligned} \Pi_{k_1, k_2, 0, 0} &= 0, & k_1 + k_2 &\geq 0, & \Pi_{0,0, l_1, l_2} &= 0, & l_1 + l_2 &\geq 0, \\ \Pi_{k_1, k_2, 1, 0} &= 0, & k_1 + k_2 &\geq 2, & \Pi_{1,0, l_1, l_2} &= 0, & l_1 + l_2 &\geq 2, \\ \Pi_{k_1, k_2, 2, 0} &= 0, & k_1 + k_2 &\geq 2, & \Pi_{2,0, l_1, l_2} &= 0, & l_1 + l_2 &\geq 2. \end{aligned}$$

Proof. Lemma 7.11 achieves the first line. The monomial x^1 being factored by $[a^1 + O_2(a)]$, we set $a^1 := a^1 + O_2(a)$ to achieve $\Pi_{1,0, l_1, l_2} = 0$, $l_1 + l_2 \geq 2$. As in the proof of Lemma 7.18, we pass to the dual equation $b = y - \Pi(x, a)$ to complete $\Pi_{k_1, k_2, 1, 0} = 0$, $k_1 + k_2 \geq 2$. Finally, $x^1 x^1$ is factored by $[a^2 + O_2(a)]$, so we proceed similarly to achieve the third line. \square

Since $\Pi(x, a)$ is assumed to be independent of b , the assumption that the Levi form of $M \subset \mathbb{C}^3$ has exactly rank 1 at every point translates to:

$$(8.57) \quad 0 \equiv \begin{vmatrix} \Pi_{x^1 a^1} & \Pi_{x^1 a^2} \\ \Pi_{x^2 a^1} & \Pi_{x^2 a^2} \end{vmatrix}.$$

For later use, it is convenient to develop somehow Π with respect to the powers of (a^1, a^2) :

$$(8.58) \quad y = b + \frac{2x^1 a^1 + x^1 x^1 a^2 + a^1 a^1 x^2}{1 - x^2 a^2} + \\ + a^2 \mathbf{b}(x) + a^1 a^2 \mathbf{d}(x) + a^2 a^2 \mathbf{e}(x) + a^1 a^1 a^1 \mathbf{f}(x) + a^1 a^1 a^2 \mathbf{g}(x) + \\ + (a^1)^4 \mathbf{R} + (a^1)^3 a^2 \mathbf{R} + a^1 (a^2)^2 \mathbf{R} + (a^2)^3 \mathbf{R},$$

with $\mathbf{R} = \mathbf{R}(x, a)$ being an unspecified remainder. Thanks to the previous lemma, the coefficients \mathbf{a} of a^1 and \mathbf{c} of $a^1 a^1$ must vanish. The function \mathbf{b} is an O_3 .

Lemma 8.59. *The function \mathbf{b} depends only on x^1 , is an $O_3(x^1)$ and the function \mathbf{g} satisfies $\mathbf{g}_{x^2 x^2}(0) = 0$.*

Proof. Developing $[1 - x^2 a^2]^{-1} = 1 + x^2 a^2 + (x^2 a^2)^2 + O_3(x^2 a^2)$, inserting the right hand side of

$$(8.60) \quad y - b = a^1 [2x^1] + a^2 [x^1 x^1 + \mathbf{b}(x)] + a^1 a^1 [x^2] + a^1 a^2 [2x^1 x^2 + \mathbf{d}(x)] + \\ + a^2 a^2 [x^1 x^1 x^2 + \mathbf{e}(x)] + a^1 a^1 a^1 [\mathbf{f}(x)] + a^1 a^1 a^2 [x^2 x^2 + \mathbf{g}(x)] + \\ + (a^1)^4 \mathbf{R} + (a^1)^3 a^2 \mathbf{R} + a^1 (a^2)^2 \mathbf{R} + (a^2)^3 \mathbf{R}$$

in the determinant (8.57) and selecting the coefficients of cst., of a^1 , of a^2 and of $a^1 a^1$, we get four PDES:

$$(8.61) \quad \begin{aligned} 0 &= 2 \mathbf{b}_{x^2}, \\ 0 &= 2 \mathbf{d}_{x^2} - 2 \mathbf{b}_{x^1}, \\ 0 &= 4 \mathbf{e}_{x^2} - 2x^1 \mathbf{d}_{x^2} - 2x^1 \mathbf{b}_{x^1} - \mathbf{d}_{x^2} \mathbf{b}_{x^1}, \\ 0 &= 2 \mathbf{g}_{x^2} - 2 \mathbf{d}_{x^1} - [6x^1 + 3 \mathbf{b}_{x^1}] \mathbf{f}_{x^2}. \end{aligned}$$

The first one yields $\mathbf{b} = \mathbf{b}(x^1)$, which must be an $O_3(x^1)$, because the whole remainder is an O_4 . Differentiating the fourth with respect to x^2 , it then follows that $\mathbf{g}_{x^2 x^2}(0) = 0$.

8.62. Associated PDE system (\mathcal{E}_5). Next, differentiating (8.60) with respect to x^1 , to $x^1 x^1$ and to $x^1 x^1 x^1$, we compute y_{x^1} and $y_{x^1 x^1}$, we substitute y_1 and $y_{1,1}$ and we push the monomials $a^2 a^2$, $a^1 a^1 a^1$ and $a^1 a^1 a^2$ in the remainder:

$$(8.63) \quad \begin{aligned} y_1 &= 2a^1 + a^2 [2x^1 + \mathbf{b}_{x^1}] + a^1 a^2 [2x^2 + \mathbf{d}_{x^1}] + (a^2)^2 \mathbf{R} + (a^1)^3 \mathbf{R} + (a^1)^2 a^2 \mathbf{R}, \\ y_{1,1} &= a^2 [2 + \mathbf{b}_{x^1 x^1}] + a^1 a^2 [\mathbf{d}_{x^1 x^1}] + (a^2)^2 \mathbf{R} + (a^1)^3 \mathbf{R} + (a^1)^2 a^2 \mathbf{R}, \\ y_{1,1,1} &= a^2 [\mathbf{b}_{x^1 x^1 x^1}] + a^1 a^2 [\mathbf{d}_{x^1 x^1 x^1}] + (a^2)^2 \mathbf{R} + (a^1)^3 \mathbf{R} + (a^1)^2 a^2 \mathbf{R}. \end{aligned}$$

Here, the written remainder *cannot* incorporate $a^1 a^1$, so it is said that the coefficient of $a^1 a^1$ does vanish in each equation above. Solving for a^1 and a^2 from the first two

equations, we get

(8.64)

$$\begin{cases} a^1 = \frac{1}{2} y_1 - y_{1,1} \left[\frac{2x^1 + \mathbf{b}_{x^1}}{4 + 2\mathbf{b}_{x^1 x^1}} \right] - y_1 y_{1,1} \left[\frac{2x^2 + \mathbf{d}_{x^1}}{8 + 4\mathbf{b}_{x^1 x^1}} \right] + \\ \quad + (y_{1,1})^2 R + (y_1)^3 R + (y_1)^2 y_{1,1} R, \\ a^2 = y_{1,1} \left[\frac{1}{2 + \mathbf{b}_{x^1 x^1}} \right] - y_1 y_{1,1} \left[\frac{\mathbf{d}_{x^1 x^1}}{2(2 + \mathbf{b}_{x^1 x^1})^2} \right] + (y_{1,1})^2 R + (y_1)^3 R + (y_1)^2 y_{1,1} R. \end{cases}$$

We then get (notice the change of remainder):

(8.65)

$$\begin{aligned} a^1 a^1 &= \frac{1}{4} (y_1)^2 - y_1 y_{1,1} \left[\frac{2x^1 + \mathbf{b}_{x^1}}{4 + 2\mathbf{b}_{x^1 x^1}} \right] - (y_1)^2 y_{1,1} \left[\frac{2x^2 + \mathbf{d}_{x^1}}{8 + 4\mathbf{b}_{x^1 x^1}} \right] + (y_1)^3 R + (y_{1,1})^2 R, \\ a^1 a^2 &= y_1 y_{1,1} \left[\frac{1}{4 + 2\mathbf{b}_{x^1 x^1}} \right] - (y_1)^2 y_{1,1} \left[\frac{\mathbf{d}_{x^1 x^1}}{(4 + 2\mathbf{b}_{x^1 x^1})^2} \right] + (y_1)^3 R + (y_{1,1})^2 R, \\ a^2 a^2 &= (y_1)^3 R + (y_{1,1})^2 R, \\ a^1 a^1 a^1 &= -(y_1)^2 y_{1,1} \left[\frac{6x^1 + 3\mathbf{b}_{x^1}}{16 + 8\mathbf{b}_{x^1 x^1}} \right] (y_1)^3 R + (y_{1,1})^2 R, \\ a^1 a^1 a^2 &= (y_1)^2 y_{1,1} \left[\frac{1}{8 + 4\mathbf{b}_{x^1 x^1}} \right] + (y_1)^3 R + (y_{1,1})^2 R. \end{aligned}$$

Differentiating (8.60) with respect to x^2 , substituting y_2 for y_{x^2} and replacing \mathbf{d}_{x^2} by \mathbf{b}_{x^1} thanks to (8.61)₂, we get

(8.66)

$$\begin{aligned} y_2 &= a^1 a^1 + a^1 a^2 [2x^1 + \mathbf{b}_{x^1}] + a^2 a^2 [x^1 x^1 + \mathbf{e}_{x^2}] + a^1 a^1 a^1 [\mathbf{f}_{x^2}] + a^1 a^1 a^2 [2x^2 + \mathbf{g}_{x^2}] + \\ &\quad + (a^1)^4 R + (a^1)^3 a^2 R + a^1 (a^2)^2 R + (a^2)^3 R. \end{aligned}$$

Replacing the monomials (8.65), we finally obtain:

(8.67)

$$\begin{aligned} y_2 &= \frac{1}{4} (y_1)^2 + (y_1)^2 y_{1,1} \left[\frac{2\mathbf{g}_{x^2} - 2\mathbf{d}_{x^1} - (6x^1 + 3\mathbf{b}_{x^1})\mathbf{f}_{x^2}}{16 + 8\mathbf{b}_{x^1 x^1}} - \frac{(2x^1 + \mathbf{b}_{x^1})\mathbf{d}_{x^1 x^1}}{(4 + 2\mathbf{b}_{x^1 x^1})^2} \right] + \\ &\quad + (y_1)^3 R + (y_{1,1})^2 R. \end{aligned}$$

Thanks to (8.61)₄, the first (big) coefficient of $(y_1)^2 y_{1,1}$ is zero; then the remainder coefficient is an $O(x^1)$, hence vanishes at $x = 0$, together with its partial first derivative with respect to x^2 . Accordingly, by $\mathbf{s}^* = \mathbf{s}^*(x^1, x^2)$, we will denote an unspecified function satisfying

(8.68)

$$\boxed{\mathbf{s}^*(0) = 0 \quad \text{and} \quad \mathbf{s}_{x^2}^*(0) = 0}.$$

Lemma 8.69. *The skeleton of the PDE system (\mathcal{E}_5) associated to the submanifold (8.58) possesses three main equations of the form*

$$(\Delta_{\mathcal{E}_5}) \quad \begin{cases} y_2 = \frac{1}{4} (y_1)^2 + (y_1)^3 r + (y_1)^4 r + (y_1)^5 r + (y_1)^6 R + \\ \quad + y_{1,1} [(y_1)^2 \mathbf{s}^* + (y_1)^3 r + (y_1)^4 r + (y_1)^5 r] + (y_{1,1})^2 R, \\ y_{1,2} = \frac{1}{2} y_1 y_{1,1} + (y_1)^3 r + (y_1)^4 r + (y_1)^5 r + (y_1)^6 R + \\ \quad + y_{1,1} [(y_1)^2 r + (y_1)^3 r + (y_1)^4 r + (y_1)^5 r] + (y_1)^6 R, \\ y_{1,1,1} = (y_1)^3 r + (y_1)^4 r + y_{1,1} [r + y_1 r + (y_1)^2 r + (y_1)^3 r] + \\ \quad + (y_{1,1})^2 [r + y_1 r + (y_1)^2 r + (y_1)^3 r] + (y_{1,1})^3 R, \end{cases}$$

where the letter r denotes an unspecified function of (x^1, x^2) , and where the coefficient s^* of $(y_1)^2 y_{1,1}$ in the first equation satisfies (8.68).

Proof. To get the second equation, we compute:

$$(8.70) \quad \begin{aligned} y_{1,2} &= a^1 a^2 [2 + \mathbf{b}_{x^1 x^1}] + a^1 a^1 a^2 [\mathbf{g}_{x^1 x^2}] + (a^1)^3 R + (a^2)^2 R \\ &= \frac{1}{2} y_1 y_{1,1} + (y_1)^2 y_{1,1} r + (y_1)^3 R + (y_{1,1})^2 R. \end{aligned}$$

The third equation is got similarly from (8.63)₃. To conclude, we develop the first two equations mod $[(y_1)^6, (y_{1,1})^2]$ and the third one mod $[(y_1)^4, (y_{1,1})^3]$. \square

This precise skeleton will be referred to as $\Delta_{\mathcal{E}_5}$ in the sequel. With the letter r , the computation rules are $\text{est}.r = r + r = r + s^* = r \cdot r = r$; sometimes, s^* may be replaced plainly by r .

8.71. Infinitesimal Lie symmetries of (\mathcal{E}_5) . Letting $\mathcal{L} = \mathcal{X}^1 \frac{\partial}{\partial x^1} + \mathcal{X}^2 \frac{\partial}{\partial x^2} + \mathcal{Y} \frac{\partial}{\partial y}$ be a candidate infinitesimal Lie symmetry and applying

$$(8.72) \quad \begin{aligned} \mathcal{L}^{(3)} &= \mathcal{X}^1 \frac{\partial}{\partial x^1} + \mathcal{X}^2 \frac{\partial}{\partial x^2} + \mathcal{Y} \frac{\partial}{\partial y} + \mathbf{Y}_1 \frac{\partial}{\partial y_1} + \mathbf{Y}_2 \frac{\partial}{\partial y_2} + \\ &+ \mathbf{Y}_{1,1} \frac{\partial}{\partial y_{1,1}} + \mathbf{Y}_{1,2} \frac{\partial}{\partial y_{1,2}} + \mathbf{Y}_{2,1} \frac{\partial}{\partial y_{2,1}} + \mathbf{Y}_{2,2} \frac{\partial}{\partial y_{2,2}} + \\ &+ \mathbf{Y}_{1,1,1} \frac{\partial}{\partial y_{1,1,1}} + \cdots + \mathbf{Y}_{2,2,2} \frac{\partial}{\partial y_{2,2,2}} \end{aligned}$$

to the skeleton $\Delta_{\mathcal{E}_5}$, we obtain firstly, computing mod $[(y_1)^5, y_{1,1}]$:

$$(8.73) \quad \begin{aligned} 0 &\equiv -\mathbf{Y}_2 + \frac{1}{2} y_1 \mathbf{Y}_1 + \\ &+ (y_1)^3 r \mathcal{X}^1 + (y_1)^4 r \mathcal{X}^1 + (y_1)^3 r \mathcal{X}^2 + (y_1)^4 r \mathcal{X}^2 + \\ &+ \mathbf{Y}_1 [(y_1)^2 r + (y_1)^3 r + (y_1)^4 r] + \\ &+ \mathbf{Y}_{1,1} [(y_1)^2 s^* + (y_1)^3 r + (y_1)^4 r], \end{aligned}$$

secondly, computing mod $[(y_1)^5, y_{1,1}]$:

$$(8.74) \quad \begin{aligned} 0 &\equiv -\mathbf{Y}_{1,2} + \frac{1}{2} y_1 \mathbf{Y}_{1,1} + \\ &+ (y_1)^3 r \mathcal{X}^1 + (y_1)^4 r \mathcal{X}^1 + (y_1)^3 r \mathcal{X}^2 + (y_1)^4 r \mathcal{X}^2 + \\ &+ \mathbf{Y}_1 [(y_1)^2 r + (y_1)^3 r + (y_1)^4 r] + \\ &+ \mathbf{Y}_{1,1} [(y_1)^2 r + (y_1)^3 r + (y_1)^4 r], \end{aligned}$$

and thirdly, computing mod $[(y_1)^3, (y_{1,1})^2]$:

$$(8.75) \quad \begin{aligned} 0 &\equiv -\mathbf{Y}_{1,1,1} + y_{1,1} \mathcal{X}^1 + y_{1,1} \mathcal{X}^2 + \\ &+ y_{1,1} y_1 \mathcal{X}^1 + y_{1,1} y_1 \mathcal{X}^2 + y_{1,1} (y_1)^2 \mathcal{X}^1 + y_{1,1} (y_1)^2 \mathcal{X}^2 + \\ &+ \mathbf{Y}_1 [(y_1)^2 r] + \mathbf{Y}_{1,1} [r + y_1 r + (y_1)^2 r] + y_{1,1} \mathbf{Y}_1 [r + y_1 r + (y_1)^2 r] + \\ &+ y_{1,1} \mathbf{Y}_{1,1} [r + y_1 r + (y_1)^2 r]. \end{aligned}$$

Specializing to $n = 2$ the formulas (3.9)(II), (3.20)(II) and (3.24)(II), we get \mathbf{Y}_1 , \mathbf{Y}_2 , $\mathbf{Y}_{1,1}$, $\mathbf{Y}_{1,2}$ and $\mathbf{Y}_{1,1,1}$:

$$(8.76) \quad \mathbf{Y}_1 = \mathcal{Y}_{x^1} + [\mathcal{Y}_y - \mathcal{X}_{x^1}^1] y_1 + [-\mathcal{X}_{x^1}^2] y_2 + [-\mathcal{X}_y^1] (y_1)^2 + [-\mathcal{X}_y^2] y_1 y_2.$$

$$(8.77) \quad \mathbf{Y}_2 = \mathcal{Y}_{x^2} + [-\mathcal{X}_{x^2}^1] y_1 + [\mathcal{Y}_y - \mathcal{X}_{x^2}^2] y_2 + [-\mathcal{X}_y^1] y_1 y_2 + [-\mathcal{X}_y^2] y_2 y_2.$$

$$(8.78) \quad \left\{ \begin{array}{l} \mathbf{Y}_{1,1} = \mathcal{Y}_{x^1 x^1} + [2\mathcal{Y}_{x^1 y} - \mathcal{X}_{x^1 x^1}^1] y_1 + [-\mathcal{X}_{x^1 x^1}^2] y_2 + [\mathcal{Y}_{yy} - 2\mathcal{X}_{x^1 y}^1] (y_1)^2 + \\ \quad + [-2\mathcal{X}_{x^1 y}^2] y_1 y_2 + [-\mathcal{X}_{yy}^1] (y_1)^3 + [-\mathcal{X}_{yy}^2] (y_1)^2 y_2 + \\ \quad + [\mathcal{Y}_y - 2\mathcal{X}_{x^1}^1] y_{1,1} + [-2\mathcal{X}_{x^1}^2] y_{1,2} + [-3\mathcal{X}_y^1] y_1 y_{1,1} + \\ \quad + [-\mathcal{X}_y^2] y_2 y_{1,1} + [-2\mathcal{X}_y^2] y_1 y_{1,2}. \end{array} \right.$$

$$(8.79) \quad \left\{ \begin{array}{l} \mathbf{Y}_{1,2} = \mathcal{Y}_{x^1 x^2} + [\mathcal{Y}_{x^2 y} - \mathcal{X}_{x^1 x^2}^1] y_1 + [\mathcal{Y}_{x^1 y} - \mathcal{X}_{x^1 x^2}^2] y_2 + \\ \quad + [-\mathcal{X}_{x^2 y}^1] (y_1)^2 + [\mathcal{Y}_{yy} - \mathcal{X}_{x^2 y}^2 - \mathcal{X}_{x^1 y}^1] y_1 y_2 + [-\mathcal{X}_{x^1 y}^2] y_2 y_2 + \\ \quad + [-\mathcal{X}_{yy}^1] (y_1)^2 y_2 + [-\mathcal{X}_{yy}^2] y_1 (y_2)^2 + \\ \quad + [-\mathcal{X}_{x^2}^1] y_{1,1} + [\mathcal{Y}_y - \mathcal{X}_{x^2}^2 - \mathcal{X}_{x^1}^1] y_{1,2} + [-\mathcal{X}_{x^1}^2] y_{2,2} + \\ \quad + [-2\mathcal{X}_y^1] y_1 y_{1,2} + [-2\mathcal{X}_y^2] y_2 y_{1,2}. \end{array} \right.$$

$$(8.80) \quad \left\{ \begin{array}{l} \mathbf{Y}_{1,1,1} = \mathcal{Y}_{x^1 x^1 x^1} + [3\mathcal{Y}_{x^1 x^1 y} - \mathcal{X}_{x^1 x^1 x^1}^1] y_1 + [-\mathcal{X}_{x^1 x^1 x^1}^2] y_2 + \\ \quad + [3\mathcal{Y}_{x^1 yy} - 3\mathcal{X}_{x^1 x^1 y}^1] (y_1)^2 + [-3\mathcal{X}_{x^1 x^1 y}^2] y_1 y_2 + \\ \quad + [\mathcal{Y}_{yyy} - 3\mathcal{X}_{x^1 yy}^1] (y_1)^3 + [-3\mathcal{X}_{x^1 yy}^2] (y_1)^2 y_2 + \\ \quad + [-\mathcal{X}_{yyy}^1] (y_1)^4 + [-\mathcal{X}_{yyy}^2] (y_1)^3 y_2 + \\ \quad + [3\mathcal{Y}_{x^1 y} - 3\mathcal{X}_{x^1 x^1}^1] y_{1,1} + [-3\mathcal{X}_{x^1 x^1}^2] y_{1,2} + \\ \quad + [3\mathcal{Y}_{yy} - 9\mathcal{X}_{x^1 y}^1] y_1 y_{1,1} + [-3\mathcal{X}_{x^1 y}^2] y_2 y_{1,1} + \\ \quad + [-6\mathcal{X}_{x^1 y}^2] y_1 y_{1,2} + [-6\mathcal{X}_{yy}^1] (y_1)^2 y_{1,1} + [-3\mathcal{X}_{yy}^2] y_1 y_2 y_{1,1} + \\ \quad + [-3\mathcal{X}_{yy}^2] (y_1)^2 y_{1,2} + [-3\mathcal{X}_y^1] (y_{1,1})^2 + [-3\mathcal{X}_y^2] y_{1,1} y_{1,2} + \\ \quad + [\mathcal{Y}_y - 3\mathcal{X}_{x^1}^1] y_{1,1,1} + [-3\mathcal{X}_{x^1}^2] y_{1,1,2} + [-4\mathcal{X}_y^1] y_1 y_{1,1,1} + \\ \quad + [-\mathcal{X}_y^2] y_2 y_{1,1,1} + [-3\mathcal{X}_y^2] y_1 y_{1,1,2}. \end{array} \right.$$

Inserting \mathbf{Y}_2 , $\mathbf{Y}_{1,2}$, $\mathbf{Y}_{1,1,1}$, \mathbf{Y}_1 , $\mathbf{Y}_{1,1}$ in the three Lie equations (8.73), (8.74), (8.75), replacing y_2 , $y_{1,2}$, $y_{1,1,1}$ by the values they have on $\Delta_{\mathcal{E}_5}$, we get firstly five linear PDEs by picking the coefficients of cst. , of y_1 , of $(y_1)^2$, of $(y_1)^3$, of $(y_1)^4$ in (8.73):

$$(8.81) \quad \left\{ \begin{array}{l} 0 = \mathcal{Y}_{x^2}, \\ 0 = \mathcal{X}_{x^2}^1 + \frac{1}{2} \mathcal{Y}_{x^1}, \\ 0 = \mathcal{Y}_y + \mathcal{X}_{x^2}^2 - 2\mathcal{X}_{x^1}^1 + r\mathcal{Y}_{x^1} + s^* \mathcal{Y}_{x^1 x^1}, \\ 0 = 2\mathcal{X}_y^1 + \mathcal{X}_{x^1}^2 + r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{X}_{x^1}^1 + \\ \quad + r\mathcal{Y}_{x^1 x^1} + s^* \mathcal{Y}_{x^1 y} + s^* \mathcal{X}_{x^1 x^1}^1, \\ 0 = \mathcal{X}_y^2 + r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{X}_{x^1}^1 + r\mathcal{X}_{x^1}^2 + r\mathcal{X}_y^1 + \\ \quad + r\mathcal{Y}_{x^1 x^1} + r\mathcal{Y}_{x^1 y} + r\mathcal{X}_{x^1 x^1}^1 + s^* \mathcal{X}_{x^1 x^1}^2 + s^* \mathcal{Y}_{yy} + s^* \mathcal{X}_{x^1 y}^1, \end{array} \right.$$

secondly, we get three more linear PDEs by picking the coefficients of $(y_1)^2$, of $(y_1)^3$, of $(y_1)^4$ in (8.74):

$$(8.82) \quad \begin{cases} 0 = 3\mathcal{Y}_{x^1y} + \mathcal{X}_{x^1x^2}^2 + 4\mathcal{X}_{x^2y}^1 - 2\mathcal{X}_{x^1x^1}^1 + r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_{x^1x^1}, \\ 0 = 2\mathcal{Y}_{yy} + 2\mathcal{X}_{x^2y}^2 - 6\mathcal{X}_{x^1y}^1 - \mathcal{X}_{x^1x^1}^2 + r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{X}_{x^1}^1 + \\ \quad + r\mathcal{Y}_{x^1x^1} + r\mathcal{Y}_{x^1y} + r\mathcal{X}_{x^1x^1}^1, \\ 0 = 4\mathcal{X}_{yy}^1 + 3\mathcal{X}_{x^1y}^2 + r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{X}_{x^1}^1 + r\mathcal{X}_{x^1}^2 + r\mathcal{X}_y^1 + \\ \quad + r\mathcal{Y}_{x^1x^1} + r\mathcal{Y}_{x^1y} + r\mathcal{X}_{x^1x^1}^1 + r\mathcal{X}_{x^1x^1}^2 + r\mathcal{Y}_{yy} + r\mathcal{X}_{x^1y}^1. \end{cases}$$

and thirdly, we get five more linear PDEs by picking the coefficients of cst., of y_1 , of $y_{1,1}$, of $y_1y_{1,1}$, of $(y_1)^2y_{1,1}$ in (8.75)

$$(8.83) \quad \begin{cases} 0 = \mathcal{Y}_{x^1x^1x^1} + r\mathcal{Y}_{x^1x^1}, \\ 0 = -3\mathcal{Y}_{x^1x^1y} + \mathcal{X}_{x^1x^1x^1}^1 + r\mathcal{Y}_{x^1x^1} + r\mathcal{Y}_{x^1y} + r\mathcal{X}_{x^1x^1}^1, \\ 0 = \mathcal{Y}_{x^1y} - \mathcal{X}_{x^1x^1}^1 + r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{X}_{x^1}^1 + r\mathcal{Y}_{x^1x^1} \\ 0 = -\frac{3}{2}\mathcal{X}_{x^1x^1}^2 + 3\mathcal{Y}_{yy} - 9\mathcal{X}_{x^1y}^1 + r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_{x^1}^1 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + \\ \quad + r\mathcal{X}_{x^1}^2 + r\mathcal{X}_y^1 + r\mathcal{Y}_{x^1x^1} + r\mathcal{Y}_{x^1y} + r\mathcal{X}_{x^1x^1}^1, \\ 0 = 6\mathcal{X}_{yy}^1 + \frac{15}{4}\mathcal{X}_{x^1y}^2 + r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_{x^1}^1 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{X}_{x^1}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_y^2 + \\ \quad + r\mathcal{Y}_{x^1x^1} + r\mathcal{Y}_{x^1y} + r\mathcal{X}_{x^1x^1}^1 + r\mathcal{X}_{x^1x^1}^2 + r\mathcal{Y}_{yy} + r\mathcal{X}_{x^1y}^1. \end{cases}$$

Proposition 8.84. *Setting as initial conditions the ten specific differential coefficients*

$$(8.85) \quad \begin{aligned} P &:= P(\mathcal{X}^1, \mathcal{X}^2, \mathcal{Y}, \mathcal{X}_y^1, \mathcal{X}_{x^2}^2, \mathcal{Y}_{x^1}, \mathcal{Y}_y, \mathcal{X}_{x^1x^2}^2, \mathcal{Y}_{x^1x^1}, \mathcal{Y}_{yy}) \\ &= r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{Y} + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{x^1x^1} + r\mathcal{Y}_{yy}, \end{aligned}$$

it follows by cross differentiations and by linear substitutions from the Lie equations (8.81)_i, $i = 1, 2, 3, 4, 5$, (8.82)_j, $j = 1, 2, 3$, (8.83)_i, $i = 1, 2, 3, 4, 5$, that $\mathcal{X}_{x^1}^1$, $\mathcal{X}_{x^1}^2$, \mathcal{Y}_{x^2} , $\mathcal{X}_{x^2}^1$, \mathcal{X}_y^2 , $\mathcal{X}_{x^1y}^1$, $\mathcal{X}_{x^2x^2}^2$, $\mathcal{Y}_{x^1x^2}$, $\mathcal{X}_{x^2y}^1$, $\mathcal{X}_{x^2y}^2$, \mathcal{Y}_{x^1y} , \mathcal{X}_{yy}^1 , \mathcal{Y}_{x^2y} , $\mathcal{X}_{x^1x^1x^2}^2$, $\mathcal{Y}_{x^1x^1x^1}$, $\mathcal{X}_{x^1x^2x^2}^2$

$\mathcal{Y}_{x^1x^1x^2}$, $\mathcal{X}_{x^1x^2y}^2$, $\mathcal{Y}_{x^1x^1y}$, \mathcal{Y}_{x^1yy} , \mathcal{Y}_{x^2yy} , \mathcal{Y}_{yyy} are uniquely determined as linear combinations of $(\mathcal{X}^1, \mathcal{X}^2, \mathcal{Y}, \mathcal{X}_y^1, \mathcal{X}_{x^2}^2, \mathcal{Y}_{x^1}, \mathcal{Y}_y, \mathcal{X}_{x^1x^1}^2, \mathcal{Y}_{x^1x^1}, \mathcal{Y}_{yy})$, namely:

$$(8.86) \quad \left\{ \begin{array}{lll} \mathcal{X}_{x^1}^1 \stackrel{1}{=} P, & \mathcal{X}_{x^1}^2 \stackrel{2}{=} P, & \mathcal{Y}_{x^2} \stackrel{3}{=} P, \\ \mathcal{X}_{x^2}^1 \stackrel{4}{=} P, & \mathcal{X}_y^2 \stackrel{5}{=} P, & \\ \mathcal{X}_{x^1y}^1 \stackrel{6}{=} P, & \mathcal{X}_{x^2x^2}^2 \stackrel{7}{=} P, & \mathcal{Y}_{x^1x^2} \stackrel{8}{=} P, \\ \mathcal{X}_{x^2y}^1 \stackrel{9}{=} P, & \mathcal{X}_{x^2y}^2 \stackrel{10}{=} P, & \mathcal{Y}_{x^1y} \stackrel{11}{=} P, \\ \mathcal{X}_{yy}^1 \stackrel{12}{=} P, & & \mathcal{Y}_{x^2y} \stackrel{13}{=} P, \\ & \mathcal{X}_{x^1x^1x^2}^2 \stackrel{14}{=} P, & \mathcal{Y}_{x^1x^1x^1} \stackrel{15}{=} P, \\ & \mathcal{X}_{x^1x^2x^2}^2 \stackrel{16}{=} P, & \mathcal{Y}_{x^1x^1x^2} \stackrel{17}{=} P, \\ & \mathcal{X}_{x^1x^2y}^2 \stackrel{18}{=} P, & \mathcal{Y}_{x^1x^1y} \stackrel{19}{=} P, \\ & & \mathcal{Y}_{x^1yy} \stackrel{20}{=} P, \\ & & \mathcal{Y}_{x^2yy} \stackrel{21}{=} P, \\ & & \mathcal{Y}_{yyy} \stackrel{22}{=} P. \end{array} \right.$$

Then the expressions P are stable under differentiation with respect to x^1 , to x^2 , to y and moreover, all other, higher order partial derivatives of \mathcal{X}^1 , of \mathcal{X}^2 , of \mathcal{Y} may be expressed as $P(\mathcal{X}^1, \mathcal{X}^2, \mathcal{Y}, \mathcal{X}_y^1, \mathcal{X}_{x^2}^2, \mathcal{Y}_{x^1}, \mathcal{Y}_y, \mathcal{X}_{x^1x^2}^2, \mathcal{Y}_{x^1x^1}, \mathcal{Y}_{yy})$.

Corollary 8.87. *Every infinitesimal Lie symmetry of the PDE system (\mathcal{E}_5) is uniquely determined by the ten initial Taylor coefficients*

$$(8.88) \quad \mathcal{X}^1(0), \mathcal{X}^2(0), \mathcal{Y}(0), \mathcal{X}_y^1(0), \mathcal{X}_{x^2}^2(0), \mathcal{Y}_{x^1}(0), \mathcal{Y}_y(0), \mathcal{X}_{x^1x^2}^2(0), \mathcal{Y}_{x^1x^1}(0), \mathcal{Y}_{yy}(0).$$

Proof of the proposition. At first, (8.83)₁ yields (8.86)₁₅; (8.81)₁ yields (8.86)₃; differentiating (8.81)₁ with respect to x^1 yields (8.86)₈; differentiating (8.81)₁ with respect to y yields (8.86)₁₃; differentiating (8.81)₁ with respect to x^1x^1 yields (8.86)₁₇; and differentiating (8.81)₁ with respect to yy yields (8.86)₂₁. Also, rewriting (8.81)₂ as

$$(8.89) \quad \mathcal{X}_{x^2}^1 = -\frac{1}{2} \mathcal{Y}_{x^1},$$

we get (8.86)₄; and rewriting (8.81)₃ as

$$(8.90) \quad \mathcal{X}_{x^1}^1 = \frac{1}{2} \mathcal{X}_{x^2}^2 + \frac{1}{2} \mathcal{Y}_y + r \mathcal{Y}_{x^1} + s^* \mathcal{Y}_{x^1x^1},$$

we get (8.86)₁.

Next, differentiating (8.81)₂ with respect to x^1 and (8.81)₃ with respect to x^2 , we get, taking account of $0 = \mathcal{Y}_{x^2y} = \mathcal{Y}_{x^1x^2} = \mathcal{Y}_{x^1x^1x^2}$, replacing $\mathcal{X}_{x^1x^2}$ by $-\frac{1}{2} \mathcal{Y}_{x^1x^1}$ and solving for $\mathcal{X}_{x^2x^2}^2$:

$$(8.91) \quad \begin{aligned} 0 &= \mathcal{X}_{x^1x^2}^1 + \frac{1}{2} \mathcal{Y}_{x^1x^1}, \\ \mathcal{X}_{x^2x^2}^2 &= -(1 + s_{x^2}^*) \mathcal{Y}_{x^1x^1} + r \mathcal{Y}_{x^1}. \end{aligned}$$

This is (8.86)₇. Differentiating (8.91)₂ with respect to x^1 , taking account of (8.83)₁, we get (8.86)₁₆:

$$(8.92) \quad \mathcal{X}_{x^1x^2x^2}^2 = r \mathcal{Y}_{x^1} + r \mathcal{Y}_{x^1x^1}.$$

We then replace $\mathcal{X}_{x^1}^1$ from (8.90) in (8.81)₄:

$$(8.93) \quad 0 = \mathcal{X}_{x^1}^2 + 2\mathcal{X}_y^1 + r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + s^*\mathcal{Y}_{x^1y} + s^*\mathcal{X}_{x^1x^1}^1.$$

We differentiate this equation with respect to x^2 , knowing $\mathcal{Y}_{x^2} = 0$:

$$(8.94) \quad 0 = \mathcal{X}_{x^1x^2}^2 + 2\mathcal{X}_{x^2y}^1 + r\mathcal{X}^1 + r\mathcal{X}_{x^2}^1 + r\mathcal{X}^2 + r\mathcal{X}_{x^2}^2 + r\mathcal{X}_{x^2x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + s_{x^2}^*\mathcal{Y}_{x^1y} + s_{x^2}^*\mathcal{X}_{x^1x^1}^1 + s^*\mathcal{X}_{x^1x^2}^1.$$

We replace: $\mathcal{X}_{x^2}^1$ from (8.89); $\mathcal{X}_{x^2x^2}^2$ from (8.91)₂; we differentiate (8.81)₂ with respect to x^1x^1 to replace $\mathcal{X}_{x^1x^1x^2}^1$ by $r\mathcal{Y}_{x^1x^1}$, thanks to (8.83)₁; and we reorganize:

$$(8.95) \quad 2\mathcal{X}_{x^2y}^1 + s_{x^2}^*\mathcal{Y}_{x^1y} + s_{x^2}^*\mathcal{X}_{x^1x^1}^1 = -\mathcal{X}_{x^1x^2}^2 + r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1}.$$

We differentiate (8.81)₂ with respect to y and (8.81)₃ with respect to x^1 :

$$(8.96) \quad \begin{aligned} \mathcal{X}_{x^2y}^1 + \frac{1}{2}\mathcal{Y}_{x^1y} &= 0, \\ \mathcal{Y}_{x^1y} - 2\mathcal{X}_{x^1x^1}^1 &= -\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_{x^1x^1}. \end{aligned}$$

For the three unknowns $\mathcal{X}_{x^1x^1}^1$, \mathcal{Y}_{x^1y} , $\mathcal{X}_{x^2y}^1$, we solve the three equations (8.95), (8.96)₁, (8.96)₂, reminding $s_{x^2}^*(0) = 0$:

$$(8.97) \quad \begin{aligned} \mathcal{X}_{x^1x^1}^1 &= r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2, \\ \mathcal{Y}_{x^1y} &= r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2, \\ \mathcal{X}_{x^2y}^1 &= r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2. \end{aligned}$$

We get (8.86)₁₁ and (8.86)₉.

Thus, we may replace $\mathcal{X}_{x^1x^1}^1$ and \mathcal{Y}_{x^1y} in (8.81)₄ to get (8.86)₂:

$$(8.98) \quad \mathcal{X}_{x^1}^2 = -2\mathcal{X}_y^1 + r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2.$$

Next, we differentiate (8.83)₃ with respect to x^1 and we replace: $\mathcal{X}_{x^1}^1$ from (8.90); $\mathcal{X}_{x^1}^2$ from (8.98); \mathcal{Y}_{x^1y} from (8.97)₂; $\mathcal{X}_{x^1x^1}^1$ from (8.97)₁; $\mathcal{Y}_{x^1x^1x^1}$ from (8.83)₁; and we compare with (8.83)₂; we differentiate (8.96)₁ with respect to x^1 and (8.96)₂ with respect to x^1 ; solving, we obtain four new relations:

$$(8.99) \quad \begin{aligned} \mathcal{X}_{x^1x^1x^1}^1 &= r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2, \\ \mathcal{Y}_{x^1x^1y} &= r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2, \\ \mathcal{X}_{x^1x^2y}^2 &= r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2, \\ \mathcal{X}_{x^1x^1x^2}^2 &= r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2. \end{aligned}$$

We get (8.86)₁₉ and (8.86)₁₄.

Next, in (8.81)₅, we replace: $\mathcal{X}_{x^1}^1$ from (8.90); $\mathcal{X}_{x^1}^2$ from (8.98); \mathcal{Y}_{x^1y} from (8.97)₂; we get:

$$(8.100) \quad \begin{aligned} \mathcal{X}_y^2 &= r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + \\ &+ s^*\mathcal{X}_{x^1x^1}^2 + s^*\mathcal{Y}_{yy} + s^*\mathcal{X}_{x^1y}^1. \end{aligned}$$

We differentiate (8.98) with respect to x^1 and we replace: $\mathcal{X}_{x^1}^1$ from (8.90); $\mathcal{X}_{x^1}^2$ from (8.98); \mathcal{Y}_{x^1y} from (8.97)₂; $\mathcal{Y}_{x^1x^1x^1}$ from (8.83)₁; $\mathcal{X}_{x^1x^1x^2}^2$ from (8.99)₄; we get:

$$(8.101) \quad \mathcal{X}_{x^1x^1}^2 + 2\mathcal{X}_{x^1y}^1 = r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2.$$

In (8.82)₂, we replace: $\mathcal{X}_{x^1}^1$ from (8.90); $\mathcal{X}_{x^1x^1}^1$ from (8.97)₁; \mathcal{Y}_{x^1y} from (8.97)₂; and we reorganize:

(8.102)

$$2\mathcal{X}_{x^2y}^2 - 6\mathcal{X}_{x^1y}^1 - \mathcal{X}_{x^1x^1}^2 = -2\mathcal{Y}_{yy} + r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2.$$

Differentiating (8.81)₃ with respect to y , we replace: \mathcal{Y}_{x^1y} from (8.97)₂; $\mathcal{Y}_{x^1x^1y}$ from (8.99)₂; and we reorganize:

$$(8.103) \quad \mathcal{X}_{x^2y}^2 - 2\mathcal{X}_{x^1y}^1 = -\mathcal{Y}_{yy} + r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2.$$

For the three unknowns $\mathcal{X}_{x^1x^1}^2$, $\mathcal{X}_{x^1y}^1$, $\mathcal{X}_{x^2y}^2$, we then solve the four equations (8.101), (8.102), (8.103), (8.83)₄ (in which we replace: $\mathcal{X}_{x^1}^1$ from (8.90); $\mathcal{X}_{x^1}^2$ from (8.98); \mathcal{Y}_{x^1y} from (8.97)₂; $\mathcal{X}_{x^1x^1}^1$ from (8.97)₁):

(8.104)

$$\mathcal{X}_{x^1x^1}^2 = r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{yy},$$

$$\mathcal{X}_{x^1y}^1 = r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{yy},$$

$$\mathcal{X}_{x^2y}^2 = r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{yy}.$$

We get (8.86)₆ and (8.86)₁₀. Replacing then $\mathcal{X}_{x^1x^1}^2$, $\mathcal{X}_{x^1y}^1$ in (8.100) gives

$$(8.105) \quad \mathcal{X}_y^2 = r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{yy}.$$

This is (8.86)₅.

Next, we differentiate (8.103) with respect to x^1 and we replace: $\mathcal{X}_{x^1}^1$ from (8.90); $\mathcal{X}_{x^1}^2$ from (8.98); \mathcal{Y}_{x^1y} from (8.97)₂; $\mathcal{Y}_{x^1x^1x^1}$ from (8.83)₁; $\mathcal{X}_{x^1x^1x^2}^2$ from (8.99)₄; we get:

$$(8.106) \quad \mathcal{Y}_{x^1yy} + \mathcal{X}_{x^1x^2y}^2 - 2\mathcal{X}_{x^1x^1y}^1 = r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2.$$

Also, we differentiate (8.83)₃ with respect to y and we replace: \mathcal{X}_y^2 from (8.105); \mathcal{Y}_{x^1y} from (8.97)₂; $\mathcal{X}_{x^1y}^1$ from (8.104)₂; $\mathcal{Y}_{x^1x^1y}$ from (8.99)₂; we get:

$$(8.107) \quad \mathcal{Y}_{x^1yy} - \mathcal{X}_{x^1x^1y}^1 = r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{yy}.$$

Also, we replace in (8.82)₃: $\mathcal{X}_{x^1}^1$ from (8.90); $\mathcal{X}_{x^1}^2$ from (8.98); \mathcal{Y}_{x^1y} from (8.97)₂; $\mathcal{X}_{x^1x^1}^1$ from (8.97)₁; $\mathcal{X}_{x^1x^1}^2$ from (8.104)₁; $\mathcal{X}_{x^1y}^1$ from (8.104)₂; we get:

$$(8.108) \quad 4\mathcal{X}_{yy}^1 + 3\mathcal{X}_{x^1y}^2 = r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{yy}.$$

We differentiate this equation with respect to x^2 and we replace: $4\mathcal{X}_{x^2yy}^1$ by $-2\mathcal{Y}_{x^1yy}$ from (8.89); $\mathcal{X}_{x^2}^1$ from (8.98); $\mathcal{X}_{x^2y}^1$ from (8.97)₃; (notice $0 = \mathcal{Y}_{x^1x^2} = \mathcal{Y}_{x^2y}$); $\mathcal{X}_{x^2x^2}^2$ from (8.91)₂; $\mathcal{X}_{x^1x^2x^2}^2$ from (8.92); we get:

(8.109)

$$-2\mathcal{Y}_{x^1yy} + 3\mathcal{X}_{x^1x^2y}^2 = r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{yy}.$$

For the three unknowns $\mathcal{X}_{x^1x^1y}^1$, \mathcal{Y}_{x^1yy} , $\mathcal{X}_{x^1x^2y}^2$, we solve the three equations (8.106), (8.107), (8.108); we get:

(8.110)

$$\mathcal{X}_{x^1x^1y}^1 = r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{yy},$$

$$\mathcal{Y}_{x^1yy} = r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{yy},$$

$$\mathcal{X}_{x^1x^2y}^2 = r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{yy}.$$

We get (8.86)₂₀ and (8.86)₁₈.

Next, in (8.93), we replace: \mathcal{Y}_{x^1y} from (8.97)₂; $\mathcal{X}_{x^1x^1}^1$ from (8.97)₁; we get:

$$(8.111) \quad \mathcal{X}_{x^1}^2 + 2\mathcal{X}_y^1 = r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2.$$

We differentiate this equation with respect to y and we replace: \mathcal{X}_y^2 from (8.105); $\mathcal{X}_{x^2y}^2$ from (8.104)₃; \mathcal{Y}_{xy} from (8.97)₂; $\mathcal{Y}_{x^1x^1y}$ from (8.99)₂; $\mathcal{X}_{x^1x^2y}^2$ from (8.99)₃; we get:

$$(8.112) \quad \mathcal{X}_{x^1y}^2 + 2\mathcal{X}_{yy}^1 = r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{yy}.$$

For the two unknowns \mathcal{X}_{yy}^1 and $\mathcal{X}_{x^1y}^2$, we solve the two equations (8.108) and (8.112); we get:

$$(8.113) \quad \begin{aligned} \mathcal{X}_{yy}^1 &= r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{yy}, \\ \mathcal{X}_{x^1y}^2 &= r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{yy}. \end{aligned}$$

We get (8.86)₁₂.

Next, we differentiate (8.113)₁ with respect to x^1 and we replace: $\mathcal{X}_{x^1}^1$, $\mathcal{X}_{x^1}^2$, $\mathcal{X}_{x^1y}^1$, \mathcal{Y}_{x^1y} , $\mathcal{Y}_{x^1x^1x^1}$, $\mathcal{X}_{x^1x^1x^2}^2$, \mathcal{Y}_{x^1yy} ; we get:

$$(8.114) \quad \mathcal{X}_{x^1yy}^1 = r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{yy}.$$

Also, we differentiate (8.113)₂ with respect to x^1 and we replace:

$$(8.115) \quad \mathcal{X}_{x^1x^1y}^2 = r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{yy}.$$

Also, we differentiate (8.83)₄ with respect to y ; we replace $\mathcal{X}_{x^1yy}^1$ from (8.114), we replace $\mathcal{X}_{x^1x^1y}^2$ from (8.115); and we achieve other evident replacements; we get:

$$(8.116) \quad \mathcal{Y}_{yyy} = r\mathcal{X}^1 + r\mathcal{X}^2 + r\mathcal{X}_y^1 + r\mathcal{X}_{x^2}^2 + r\mathcal{Y}_{x^1} + r\mathcal{Y}_y + r\mathcal{Y}_{x^1x^1} + r\mathcal{X}_{x^1x^2}^2 + r\mathcal{Y}_{yy}.$$

This is (8.96)₂₂, which completes the proof. \square

Theorem 8.117. *The bound $\dim \mathfrak{S}\mathfrak{Y}\mathfrak{M}(\mathcal{E}_5) \leq 10$ is attained if and only if (\mathcal{E}_5) is equivalent, through a diffeomorphism $(x^1, x^2, y) \mapsto (X^1, X^2, Y)$, to the model system*

$$(8.118) \quad Y_{X^2} = 0, \quad Y_{X^1X^1X^1} = 0.$$

Proof. Firstly, setting $r = s^* = 0$ everywhere, the solution to (8.81), (8.82), (8.83) is

$$(8.119) \quad \begin{aligned} \mathcal{X}^1 &= k + (c + j)x^1 - bx^2 - hy + ex^1x^1 - dx^1x^2 + fx^1y - ex^2y, \\ \mathcal{X}^2 &= g + 2hx^1 + 2jx^2 - dx^2x^2 + 2ex^1x^2 - fx^1x^1, \\ \mathcal{Y} &= a + 2bx^1 + 2cy + dx^1x^1 + 2ex^1y + fyy, \end{aligned}$$

where $a, b, c, d, e, f, g, h, j, k \in \mathbb{K}$ are arbitrary. Computing the third prolongations of the ten vector fields

$$(8.120) \quad \begin{aligned} &\frac{\partial}{\partial x^1}, \quad \frac{\partial}{\partial x^2}, \quad \frac{\partial}{\partial y}, \\ &-x^2 \frac{\partial}{\partial x^1} + 2x^1 \frac{\partial}{\partial y}, \quad x^1 \frac{\partial}{\partial x^1} + 2y \frac{\partial}{\partial y}, \quad x^1 \frac{\partial}{\partial x^1} + 2x^2 \frac{\partial}{\partial x^2}, \quad -y \frac{\partial}{\partial x^1} + 2x^1 \frac{\partial}{\partial x^2}, \\ &-x^1x^2 \frac{\partial}{\partial x^1} - x^2x^2 \frac{\partial}{\partial x^2} + x^1x^1 \frac{\partial}{\partial y}, \quad (x^1x^1 - x^2y) \frac{\partial}{\partial x^1} + 2x^1x^2 \frac{\partial}{\partial x^2} + 2x^1y \frac{\partial}{\partial y}, \\ &x^1y \frac{\partial}{\partial x^1} - x^1x^1 \frac{\partial}{\partial x^2} + yy \frac{\partial}{\partial y} \end{aligned}$$

one verifies that they all are tangent to the skeleton $y_2 = \frac{1}{4}(y_1)^2$, $y_{1,1,1} = 0$. Thus the bound is attained. One then verifies ([FK2005a]) that the spanned Lie algebra is isomorphic to $\mathfrak{so}(5, \mathbb{C})$.

Lemma 8.121. *Assuming the normalizations of Lemma 8.54, the remainder O_4 in (8.53) is an $O_3(x^1, a^1)$:*

$$(8.122) \quad y = b + \frac{2x^1a^1 + x^1x^1a^2 + a^1a^1x^2}{1 - x^2a^2} + (x^1)^3 R + (x^1)^2 a^1 R + x^1 (a^1)^2 R + (a^1)^3 R.$$

Proof. Indeed, writing

$$(8.123) \quad y = b + x^1 \Lambda^{1,0} + a^1 \Lambda^{0,1} + x^1 x^1 \Lambda^{2,0} + x^1 a^1 \Lambda^{1,1} + a^1 a^1 \Lambda^{0,2} + O_3(x^1, a^1),$$

with $\Lambda^{i,j} = \Lambda^{i,j}(x^2, a^2)$, and developing the determinant (8.54) with respect to the powers of (x^1, a^1) , the vanishing of the coefficients of cst. , of x^1 , of a^1 yields the system

$$(8.124) \quad \begin{cases} 0 \equiv \Lambda_{a^2}^{1,0} \Lambda_{x^2}^{0,1}, \\ 0 \equiv \Lambda^{1,1} \Lambda_{x^2 a^2}^{1,0} - 2 \Lambda_{a^2}^{2,0} \Lambda_{x^2}^{0,1} - \Lambda_{x^2}^{1,1} \Lambda_{a^2}^{1,0}, \\ 0 \equiv \Lambda^{1,1} \Lambda_{x^2 a^2}^{0,1} - \Lambda_{a^2}^{1,1} \Lambda_{x^2}^{0,1} - 2 \Lambda_{x^2}^{0,2} \Lambda_{a^2}^{1,0}. \end{cases}$$

If the first equation yields $\Lambda_{a^2}^{1,0} \equiv 0$, replacing in the second, using $\Lambda^{2,0} = a^2 + O_2$, we deduce that $\Lambda_{x^2}^{0,1} \equiv 0$ also. Similarly, $\Lambda_{x^2}^{0,1} \equiv 0$ implies $\Lambda_{a^2}^{1,0} \equiv 0$. Since the coordinate system satisfies the normalization $\Pi(0, a) \equiv \Pi(x, 0) \equiv 0$, necessarily $\Lambda^{1,0} = O(a^2)$ and $\Lambda^{0,1} = O(x^2)$. We deduce:

$$(8.125) \quad 0 \equiv \Lambda^{1,0} \equiv \Lambda^{0,1}.$$

Redeveloping the determinant, the vanishing of the coefficients of $x^1 x^1$, of $x^1 a^1$, of $a^1 a^1$ yields the system

$$(8.126) \quad \begin{cases} 0 \equiv \Lambda^{1,1} \Lambda_{x^2 a^2}^{2,0} - 2 \Lambda_{a^2}^{2,0} \Lambda_{x^2}^{1,1}, \\ 0 \equiv \Lambda^{1,1} \Lambda_{x^2 a^2}^{1,1} - \Lambda_{a^2}^{1,1} \Lambda_{x^2}^{1,1} - 4 \Lambda_{a^2}^{2,0} \Lambda_{x^2}^{0,2}, \\ 0 \equiv \Lambda^{1,1} \Lambda_{x^2 a^2}^{0,2} - 2 \Lambda_{x^2}^{0,2} \Lambda_{a^2}^{1,1}. \end{cases}$$

Since $\Lambda^{1,1}(0) = 2 \neq 0$, we may divide by $\Lambda^{1,1}$, obtaining a PDE system with the three functions $\Lambda_{x^2 a^2}^{2,0}$, $\Lambda_{x^2 a^2}^{1,1}$, $\Lambda_{x^2 a^2}^{0,2}$ in the left hand side. We observe that the normalizations of Lemma 8.55 entail

$$(8.127) \quad \Lambda^{2,0} = a^2 + O(x^2 a^2), \quad \Lambda^{1,1} = 2 + O(x^2 a^2), \quad \Lambda^{0,2} = x^2 + O(x^2 a^2).$$

By cross differentiations in the mentioned PDE system, it follows that all the Taylor coefficients of $\Lambda^{2,0}$, $\Lambda^{1,1}$, $\Lambda^{0,2}$ are uniquely determined. As already discovered in [GM2003b], the unique solution

$$(8.128) \quad \Lambda^{2,0} = \frac{a^2}{1 - x^2 a^2}, \quad \Lambda^{1,1} = \frac{2}{1 - x^2 a^2}, \quad \Lambda^{0,2} = \frac{x^2}{1 - x^2 a^2},$$

guarantees, when the remainder $O_3(x^1, a^1)$ vanishes, that the determinant (8.45) indeed vanishes identically. \square

Conversely, suppose that $\dim \mathfrak{S}\mathfrak{M}(\mathcal{E}_5) = 10$ is maximal.

With $\varepsilon \neq 0$ small, replacing $(x^1, x^2, y, a^1, a^2, b)$ by $(\varepsilon x^1, x^2, \varepsilon^2 y, \varepsilon a^1, a^2, \varepsilon \varepsilon b)$ in (8.122) and dividing by $\varepsilon \varepsilon$, the remainder terms become small:

$$(8.129) \quad y = b + \frac{2x^1a^1 + x^1x^1a^2 + a^1a^1x^2}{1 - x^2a^2} + O(\varepsilon).$$

Then all the remainders in the equations $\Delta_{\mathcal{E}_5}$ of the skeleton are $O(\varepsilon)$. We get ten generators similar to (8.120), plus an $O(\varepsilon)$ perturbation. Thanks to the rigidity of $\mathfrak{so}(5, \mathbb{C})$, Theorem 5.15 provides a change of generators, close to the 10×10 identity matrix, leading to the same structure constants as those of the ten vector fields (8.120). As in the end

of the proof of Theorem 5.13, we may then straighten some relevant vector fields (exercise) and finally check that their tangency to the skeleton implies that it is the model one. Theorem 8.117 is proved. \square

Corollary 8.130. *Let $M \subset \mathbb{C}^3$ be a connected real analytic hypersurface whose Levi form has uniform rank 1 that is 2-nondegenerate at every point. Then*

$$(8.131) \quad \dim \mathfrak{hol}(M) \leq 10,$$

and the bound is attained if and only if M is locally, in a neighborhood of Zariski-generic points, biholomorphic to the model M_0 .

§9. DUAL SYSTEM OF PARTIAL DIFFERENTIAL EQUATIONS

9.1. Solvability with respect to the variables. Let \mathcal{M} be as in §2.10 defined by $y = \Pi(x, a, b)$ or dually by $b = \Pi^*(a, x, y)$.

Definition 9.2. \mathcal{M} is *solvable with respect to the variables* if there exist an integer $\kappa^* \geq 1$ and multiindices $\delta(1), \dots, \delta(n) \in \mathbb{N}^p$ with $|\delta(l)| \geq 1$ for $l = 1, \dots, n$ and $\max_{1 \leq l \leq n} |\delta(l)| = \kappa^*$, together with integers $j(1), \dots, j(n)$ with $1 \leq j(l) \leq m$ such that the local \mathbb{K} -analytic map

$$(9.3) \quad \mathbb{K}^{n+m} \ni (x, y) \longmapsto \left((\Pi^{*j}(0, x, y))^{1 \leq j \leq m}, \left(\Pi_{a^{\delta(l)}}^{*j(l)}(0, x, y) \right)_{1 \leq l \leq n} \right) \in \mathbb{K}^{m+n}$$

is of rank equal to $n + m$ at $(x, y) = (0, 0)$

If \mathcal{M} is a complexified generic submanifold, solvability with respect to the parameters is equivalent to solvability with respect to the variables, because $\Pi^* = \overline{\Pi}$. This is untrue in general: with $n = 2, m = 1$, consider the system $y_{x^2} = 0, y_{x^1 x^1} = 0$, whose general solutions is $y(x) = b + x_1 a$ with x^2 absent.

To characterize generally such a degeneration property, we develop both

$$(9.4) \quad \begin{aligned} y^j &= \Pi^j(x, a, b) = \sum_{\beta \in \mathbb{N}^n} x^\beta \Pi_\beta^j(a, b) \quad \text{and} \\ b^j &= \Pi^{*j}(a, x, y) = \sum_{\delta \in \mathbb{N}^p} a^\delta \Pi_\delta^{*j}(x, y), \end{aligned}$$

with analytic functions $\Pi_\beta^j(a, b), \Pi_\delta^{*j}(x, y)$ and we introduce two \mathbb{K}^∞ -valued maps

$$(9.5) \quad \begin{aligned} \mathcal{Q}_\infty &: (a, b) \longmapsto \left(\Pi_\beta^j(a, b) \right)_{\beta \in \mathbb{N}^n}^{1 \leq j \leq m} \quad \text{and} \\ \mathcal{Q}_\infty^* &: (x, y) \longmapsto \left(\Pi_\delta^{*j}(x, y) \right)_{\delta \in \mathbb{N}^p}^{1 \leq j \leq m}. \end{aligned}$$

Since $b \mapsto \left(\Pi_0^j(0, b) \right)_{1 \leq j \leq m}$ and $y \mapsto \left(\Pi_0^{*j}(0, y) \right)_{1 \leq j \leq m}$ are already both of rank m at the origin, the generic ranks of these two maps, defined by testing the nonvanishing of minors of their infinite Jacobian matrices, satisfy

$$(9.6) \quad \begin{aligned} \text{genrk } \mathcal{Q}_\infty &= m + p_{\mathcal{M}} \quad \text{and} \\ \text{genrk } \mathcal{Q}_\infty^* &= m + n_{\mathcal{M}} \end{aligned}$$

for some two integers $0 \leq p_{\mathcal{M}} \leq p$ and $0 \leq n_{\mathcal{M}} \leq n$. So at a Zariski-generic point, the ranks are *equal* to $m + p_{\mathcal{M}}$ and to $m + n_{\mathcal{M}}$.

Proposition 9.7. *There exists a local proper \mathbb{K} -analytic subset $\Sigma_{\mathcal{M}}$ of $\mathbb{K}_x^n \times \mathbb{K}_y^m \times \mathbb{K}_a^p \times \mathbb{K}_b^m$ whose equations, of the specific form*

$$(9.8) \quad \Sigma_{\mathcal{M}} = \left\{ r_\nu(a, b) = 0, \nu \in \mathbb{N}, \quad r_\mu^*(x, y) = 0 \quad \mu \in \mathbb{N} \right\},$$

are obtained by equating to zero all $(m + p_{\mathcal{M}}) \times (m + p_{\mathcal{M}})$ minors of $\text{Jac } \mathcal{Q}_{\infty}$ and all $(m + n_{\mathcal{M}}) \times (m + n_{\mathcal{M}})$ minors of $\text{Jac } \mathcal{Q}_{\infty}^*$, such that for every point $p = (x_p, y_p, a_p, b_p) \notin \Sigma_{\mathcal{M}}$, there exists a local change of coordinates respecting the separation of the variables (x, y) and (a, b)

$$(9.9) \quad (x, y, a, b) \mapsto (\varphi(x, y), h(a, b)) =: (x', y', a', b')$$

by which \mathcal{M} is transformed to a submanifold \mathcal{M}' centered and localized at $p' = p$ having equations

$$(9.10) \quad y' = \Pi'(x', a', b') \quad \text{and dually} \quad b' = \Pi'^*(a', x', y')$$

with Π' and Π'^* independent of

$$(9.11) \quad (x'_{n_{\mathcal{M}+1}}, \dots, x'_n) \quad \text{and of} \quad (a'_{p_{\mathcal{M}+1}}, \dots, a'_p).$$

So \mathcal{M}' , may be considered to be living in $\mathbb{K}_{x'}^{n_{\mathcal{M}}} \times \mathbb{K}_{y'}^m \times \mathbb{K}_{a'}^{p_{\mathcal{M}}} \times \mathbb{K}_{b'}^m$ and in such a smaller space, at $p' = p$, it is solvable both with respect to the parameters and to the variables.

Interpretation: by forgetting some innocuous variables, at a Zariski-generic point, any \mathcal{M} is both solvable with respect to the parameters and to the variables. These two assumptions will be held up to the end of this Part I.

9.12. Dual system (\mathcal{E}^*) and isomorphisms $\mathfrak{G}\mathfrak{M}(\mathcal{E}) \simeq \mathfrak{G}\mathfrak{M}(\mathcal{V}_S(\mathcal{E})) = \mathfrak{G}\mathfrak{M}(\mathcal{V}_S(\mathcal{E}^*)) \simeq \mathfrak{G}\mathfrak{M}(\mathcal{E}^*)$. To a system (\mathcal{E}) , we associate its submanifold of solutions $\mathcal{M} := \mathcal{V}_S(\mathcal{E})$. Assuming it to be solvable with respect to the variables and proceeding as in §2.10, we can derive a *dual system of completely integrable partial differential equations* of the form

$$(9.12) \quad b_{a\gamma}^j(a) = G_{\gamma}^j(a, b(a), (b_{a\delta(l)}^{j(l)}(a))_{1 \leq l \leq n}),$$

where $(j, \gamma) \neq (j, 0)$ and $\neq (j(l), \delta(l))$. Its submanifold of solutions $\mathcal{V}_S(\mathcal{E}^*) \equiv \mathcal{V}_S(\mathcal{E})$ has equations dual to those of $\mathcal{V}_S(\mathcal{E})$.

Theorem 9.13. *Under the assumption of twin solvability, we have:*

$$(9.14) \quad \mathfrak{G}\mathfrak{M}(\mathcal{E}) \simeq \mathfrak{G}\mathfrak{M}(\mathcal{V}_S(\mathcal{E})) = \mathfrak{G}\mathfrak{M}(\mathcal{V}_S(\mathcal{E}^*)) \simeq \mathfrak{G}\mathfrak{M}(\mathcal{E}^*),$$

through $\mathcal{L} \longleftrightarrow \mathcal{L} + \mathcal{L}^* = \mathcal{L}^* + \mathcal{L} \longleftrightarrow \mathcal{L}^*$.

§10. FUNDAMENTAL PAIR OF FOLIATIONS AND COVERING PROPERTY

10.1. Fundamental pair of foliations on \mathcal{M} . As in §2, let (\mathcal{E}) and $\mathcal{M} = \mathcal{V}_S(\mathcal{E})$ be defined by $y = \Pi(x, a, b)$ or dually by $b = \Pi^*(a, x, y)$. Abbreviate

$$(10.2) \quad z := (x, y) \quad \text{and} \quad c := (a, b).$$

Every transformation $(z, c) \mapsto (\varphi(z), h(c))$ belonging to $\mathbb{G}_{v,p}$ stabilizes both $\{z = \text{cst.}\}$ and $\{c = \text{cst.}\}$. Accordingly, the two foliations of \mathcal{M}

$$(10.3) \quad F_v := \bigcup_{c_0} \mathcal{M} \cap \{c = c_0\} \quad \text{and} \quad F_p := \bigcup_{z_0} \mathcal{M} \cap \{z = z_0\}$$

are invariant under changes of coordinates. We call (F_v, F_p) the *fundamental pair of foliations* on \mathcal{M} . The leaves of the *foliation by variables* F_v are n -dimensional:


$$(10.4) \quad F_v(c_0) = \{(z, c_0) : y = \Pi(x, c_0)\}$$

The leaves of the *foliation by parameters* F_p are p -dimensional:

$$(10.5) \quad F_p(z_0) = \{(z_0, c) : b = \Pi^*(a, z_0)\}$$

We draw a diagram. In it, the positive codimension is invisible:

$$(10.6) \quad m = \dim \mathcal{M} - \dim F_v - \dim F_p \geq 1$$



10.7. Chains Γ_k and dual chains Γ_k^* . Similarly as in [GM2004, Me2005a, Me2005b, MP2005] (in a CR context), we introduce two collections $(L_k)_{1 \leq k \leq n}$ and $(L_q^*)_{1 \leq q \leq p}$ of vector fields whose integral manifolds coincide with the leaves of F_v and of F_p :

$$(10.8) \quad \begin{cases} L_k := \frac{\partial}{\partial x_k} + \sum_{j=1}^m \frac{\partial \Pi^j}{\partial x_k}(x, a, b) \frac{\partial}{\partial y^j}, & k = 1, \dots, n, \\ L_q^* := \frac{\partial}{\partial a^q} + \sum_{j=1}^m \frac{\partial \Pi^{*j}}{\partial a^q}(a, x, y) \frac{\partial}{\partial b^j}, & q = 1, \dots, p. \end{cases}$$

Let $(z_0, c_0) = (x_0, y_0, a_0, b_0) \in \mathcal{M}$ be a fixed point, let $x_1 := (x_1^1, \dots, x_1^n) \in \mathbb{K}^n$ and define the multiple flow map

$$(10.9) \quad \begin{cases} L_{x_1}(x_0, y_0, a_0, b_0) := \exp(x_1 L)(p_0) := \exp(x_1^n L_n(\dots(\exp(x_1^1 L_1(z_0, c_0)))) \dots) \\ \quad \quad \quad := (x_0 + x_1, \Pi(x_0 + x_1, a_0, b_0), a_0, b_0). \end{cases}$$

Similarly, for $a_1 = (a_1^1, \dots, a_1^p) \in \mathbb{K}^p$, define the multiple flow map

$$(10.10) \quad L_{a_1}^*(x_0, y_0, a_0, b_0) := (x_0, y_0, a_0 + a_1, \Pi^*(a_0 + a_1, x_0, y_0)).$$

Starting from the $(z_0, c_0) = (0, 0)$ and moving alternately along $F_v, F_p, F_v, etc.$, we obtain

$$(10.11) \quad \begin{cases} \Gamma_1(x_1) := L_{x_1}(0), \\ \Gamma_2(x_1, a_1) := L_{a_1}^*(L_{x_1}(0)), \\ \Gamma_3(x_1, a_1, x_2) := L_{x_2}(L_{a_1}^*(L_{x_1}(0))), \\ \Gamma_4(x_1, a_1, x_2, a_2) := L_{a_2}^*(L_{x_2}(L_{a_1}^*(L_{x_1}(0)))) \end{cases}$$

and so on. Generally, we get *chains* $\Gamma_k := \Gamma_k([xa]_k)$, where $[xa]_k := (x_1, a_1, x_2, a_2, \dots)$ with exactly k terms, where each $x_l \in \mathbb{K}^n$ and each $a_l \in \mathbb{K}^p$.

If, instead, the first movement consists in moving along F_p , we start with $\Gamma_1^*(a_1) := L_{a_1}^*(0)$, $\Gamma_2^*(a_1, x_1) := L_{x_1}(L_{a_1}^*(0))$, *etc.*, and generally we get *dual chains* $\Gamma_k^*([ax]_k)$, where $[ax]_k := (a_1, x_1, a_2, x_2, \dots)$, with exactly k terms. Both Γ_k and Γ_k^* have range in \mathcal{M} .

For $k = 1, 2, 3, \dots$, integers e_k and e_k^* are defined inductively by

$$(10.12) \quad \begin{cases} e_1 + e_2 + e_3 + \dots + e_k = \text{genrk}_{\mathbb{K}}(\Gamma_k), \\ e_1^* + e_2^* + e_3^* + \dots + e_k^* = \text{genrk}_{\mathbb{K}}(\Gamma_k^*). \end{cases}$$

By (10.9) and (10.10), it is clear that $e_1 = n$, $e_2 = p$, $e_1^* = p$, and $e_2^* = n$.

Example 10.13. For $y_{xx} = 0$, the submanifold of solutions \mathcal{M} is simply $y = b + xa$, whence

$$(10.14) \quad \begin{cases} \Gamma_1(x_1) = (x_1, 0, 0, 0), \\ \Gamma_2(x_1, a_1) = (x_1, 0, a_1, -x_1 a_1), \\ \Gamma_3(x_1, a_1, x_2) = (x_1 + x_2, x_2 a_1, a_1, -x_1 a_1). \end{cases}$$

The rank at $(0, 0, 0)$ of Γ_3 is equal to two, not more. However, its generic rank is equal to three. Similar observations hold for the two submanifolds of solutions $y = b + xxa + xaa$ and $y = b + xa^1 + xxa^2$ (in \mathbb{K}^5).

Lemma 10.15. *If $\text{genrk}_{\mathbb{K}}(\Gamma_{k+1}) = \text{genrk}_{\mathbb{K}}(\Gamma_k)$, then for each positive integer $l \geq 1$, we have $\text{genrk}_{\mathbb{K}}(\Gamma_{k+l}) = \text{genrk}_{\mathbb{K}}(\Gamma_k)$. The same stabilization property holds for Γ_k^* .*

10.16. Covering property. We now formulate a central concept.

Definition 10.17. The pair of foliations (F_v, F_p) is *covering at the origin* if there exists an integer k such that the generic rank of Γ_k is (maximal possible) equal to $\dim_{\mathbb{K}} \mathcal{M}$. Since for $a_1 = 0$, the dual $(k+1)$ -th chain Γ_{k+1}^* identifies with the k -th chain Γ_k , the same property holds for the dual chains.

Example 10.18. With $n = 1$, $m = 2$ and $p = 1$ the submanifold defined by $y^1 = b^1$ and $y^2 = b^2 + xa$ is twin solvable, but its pair of foliations is not covering at the origin. Then $\mathfrak{S}\mathfrak{M}(\mathcal{M})$ is infinite-dimensional, since for $a = a(y^1)$ an arbitrary function, it contains $a(y^1) \frac{\partial}{\partial y^1} + a(b^1) \frac{\partial}{\partial b^1}$.

Because we aim only to study finite-dimensional Lie symmetry groups of partial differential equations, in the remainder of this Part I, we will constantly assume the covering property to hold.

By Lemma 10.15, there exist two well defined integers μ and μ^* such that $e_3, e_4, \dots, e_{\mu+1} > 0$, but $e_{\mu+l} = 0$ for all $l \geq 2$ and similarly, $e_3^*, e_4^*, \dots, e_{\mu^*+1}^* > 0$, but $e_{\mu^*+l}^* = 0$ for all $l \geq 2$. Since the pair of foliations is covering, we have the two dimension equalities

$$(10.19) \quad \begin{cases} n + p + e_3 + \dots + e_{\mu+1} = \dim_{\mathbb{K}} \mathcal{M} = n + m + p, \\ p + n + e_3^* + \dots + e_{\mu^*+1}^* = \dim_{\mathbb{K}} \mathcal{M} = n + m + p. \end{cases}$$

By definition, the ranges of $\Gamma_{\mu+1}$ and of $\Gamma_{\mu^*+1}^*$ cover (at least; more is true, see: Theorem 10.28) an open subset of \mathcal{M} . Also, it is elementary to verify the four inequalities

$$(10.20) \quad \begin{aligned} \mu &\leq 1 + m, & \mu^* &\leq 1 + m, \\ \mu &\leq \mu^* + 1, & \mu^* &\leq \mu + 1. \end{aligned}$$

In fact, since Γ_{k+1} with $x_1 = 0$ identifies with Γ_k^* , the second line follows.

Definition 10.21. The *type of the covering pair of foliations* (F_v, F_p) is the pair of integers

$$(10.22) \quad (\mu, \mu^*), \quad \text{with} \quad \max(\mu, \mu^*) \leq 1 + m.$$

Example 10.23. (Continued) We write down the explicit expressions of Γ_4 and of Γ_5 :

$$(10.24) \quad \begin{cases} \Gamma_4(x_1, a_1, x_2, a_2; 0) = (x_1 + x_2, x_2 a_1, a_1 + a_2, -x_1 a_1 - x_1 a_2 - x_2 a_2), \\ \Gamma_5(x_1, a_1, x_2, a_2, x_3; 0) = (x_1 + x_2 + x_3, x_2 a_1 + x_3 a_1 + x_3 a_2, a_1 + a_2 \\ \hspace{15em} - x_1 a_1 - x_1 a_2 - x_2 a_2). \end{cases}$$

Here, $\dim \mathcal{M} = 3$. By computing its Jacobian matrix, Γ_5 is of rank 3 at every point $(x_1, a_1, 0, -a_1, -x_1) \in \mathbb{K}^5$ with $a_1 \neq 0$. Since (obviously)

$$(10.25) \quad \Gamma_5(x_1, a_1, 0, -a_1, -x_1) = 0 \in \mathcal{M},$$

we deduce that Γ_5 is submersive (“covering”) from a small neighborhood of $(x_1, a_1, 0, -a_1, -x_1)$ in \mathbb{K}^5 onto a neighborhood of the origin in \mathcal{M} .

10.26. Covering a neighborhood of the origin in \mathcal{M} . For $(z_0, c_0) \in \mathcal{M}$ fixed and close to the origin, we denote by $\Gamma_k([xa]_k; (z_0, c_0))$ and by $\Gamma_k^*([ax]_k; (z_0, c_0))$ the (dual) chains issued from (z_0, c_0) . For given parameters $[xa]_k = (x_1, a_1, x_2, \dots)$, we denote by $[-xa]_k$ the collection $(\dots, -x_2, -a_1, -x_2)$ with minus signs and reverse order; similarly, we introduce $[-ax]_k$. Notably, we have $L_{-x_1}(L_{x_1}(0)) = 0$ (because $L_{-x_1+x_1}(\cdot) = L_0(\cdot) = \text{Id}$), and also $L_{-x_1}(L_{-a_1}^*(L_{a_1}^*(L_{x_1}(0)))) = 0$ and generally:

$$(10.27) \quad \Gamma_k([-xa]_k; \Gamma_k([xa]_k; 0)) \equiv 0.$$

Geometrically speaking, by following backward the k -th chain Γ_k , we come back to 0.

Theorem 10.28. ([Me2005a, Me2005b], [*]) *The two maps $\Gamma_{2\mu+1}$ and $\Gamma_{2\mu^*+1}^*$ are submersive onto a neighborhood of the origin in \mathcal{M} . Precisely, there exist two points $[xa]_{2\mu+1}^0 \in \mathbb{K}^{(\mu+1)n+\mu p}$ and $[ax]_{2\mu^*+1}^0 \in \mathbb{K}^{\mu^*n+(\mu^*+1)p}$ arbitrarily close to the origin with $\Gamma_{2\mu+1}([xa]_{2\mu+1}^0) = 0$ and $\Gamma_{2\mu^*+1}^*([ax]_{2\mu^*+1}^0) = 0$ such that the two maps*

$$(10.29) \quad \begin{cases} \mathbb{K}^{(\mu+1)n+\mu p} \ni [xa]_{2\mu+1} \longmapsto \Gamma_{2\mu+1}([xa]_{2\mu+1}) \in \mathcal{M} & \text{and} \\ \mathbb{K}^{\mu^*n+(\mu^*+1)p} \ni [ax]_{2\mu^*+1} \longmapsto \Gamma_{2\mu^*+1}^*([ax]_{2\mu^*+1}) \in \mathcal{M} \end{cases}$$

are of rank $n + m + p = \dim_{\mathbb{K}} \mathcal{M}$ at the points $[xa]_{2\mu}^0$ and $[ax]_{2\mu^}^0$ respectively. In particular, the ranges of the two maps $\Gamma_{2\mu+1}$ and $\Gamma_{2\mu^*+1}^*$ cover a neighborhood of the origin in \mathcal{M} .*

Let $\pi_z(z, c) := z$ and $\pi_c(z, c) := c$ be the two canonical projections. The next corollary will be useful in Section 12. In the example above, it also follows that the map

$$(10.30) \quad [xa]_4 \mapsto \pi_c(\Gamma_4([xa]_4)) = (a_1 + a_2, -x_1a_1 - x_1a_2 - x_2a_2) \in \mathbb{K}^2$$

is of rank two at all points $[xa]_4^0 := (x_1^0, a_1^0, 0, -a_1^0)$ with $a_1^0 \neq 0$.

Corollary 10.31. ([Me2005a, Me2005b], [*]) *There exist two points $[xa]_{2\mu}^0 \in \mathbb{K}^{\mu(n+p)}$ and $[ax]_{2\mu^*}^0 \in \mathbb{K}^{\mu^*(n+p)}$ arbitrarily close to the origin with $\pi_c(\Gamma_{2\mu}([xa]_{2\mu}^0)) = 0$ and $\pi_z(\Gamma_{2\mu^*}^*([ax]_{2\mu^*}^0)) = 0$ such that the two maps*

$$(10.32) \quad \begin{cases} \mathbb{K}^{\mu(n+p)} \ni [xa]_{2\mu} \longmapsto \pi_c(\Gamma_{2\mu}([xa]_{2\mu})) \in \mathbb{K}^{m+p} & \text{and} \\ \mathbb{K}^{\mu^*(n+p)} \ni [ax]_{2\mu^*} \longmapsto \pi_z(\Gamma_{2\mu^*}^*([ax]_{2\mu^*})) \in \mathbb{K}^{n+m} \end{cases}$$

are of rank $m + p$ at the point $[xa]_{2\mu}^0 \in \mathbb{K}^{\mu(n+p)}$ and of rank $n + m$ at the point $[ax]_{2\mu^}^0 \in \mathbb{K}^{\mu^*(n+p)}$.*

In the case $m = 1$ (single dependent variable $y \in \mathbb{K}$), the covering property always hold with $\mu = \mu^* = 2$.

§11. FORMAL AND SMOOTH EQUIVALENCES BETWEEN SUBMANIFOLDS OF SOLUTIONS

11.1. Transformations of submanifolds of solutions. Lemma 7.3 shows that every equivalence φ between two PDE systems (\mathcal{E}) and (\mathcal{E}') lifts as a transformation which respects the separation between variables and parameters of the form

(11.2)


$$(x, y, a, b) \longmapsto (\phi(x, y), \psi(x, y), f(a, b), g(a, b)) = (\varphi(x, y), h(a, b)) =: (x', y', a', b')$$

from the source submanifolds of solutions $\mathcal{M} := \mathcal{V}_S(\mathcal{E})$ to the target $\mathcal{M}' := \mathcal{V}_S(\mathcal{E}')$, whose equations are

$$(11.3) \quad \begin{array}{ll} y = \Pi(x, c) & \text{or dually} \quad b = \Pi^*(a, z) \quad \text{and} \\ y' = \Pi'(x', c') & \text{or dually} \quad b' = \Pi'^*(a', z'). \end{array}$$

The study of transformations between submanifolds of solutions possesses strong similarities with the study of CR mappings between CR manifolds ([Pi1975, We1977, DW1980, BJT1985, DF1988, BER1999, Me2005a, Me2005b]). In fact, one may transfer the whole theory of the analytic reflection principle to this more general context. In the present §10 and in the next §11, we select and establish some of the results that are useful to the Lie theory. Some accessible open questions will also be formulated.

Maps of the form (11.2) send leaves of F_v and of F_p to leaves of F'_v , and of F'_p , respectively.



11.4. Regularity and jet parametrization. Some strong rigidity properties underly the above diagram. Especially, the smoothness of the two pairs (F_v, F_p) and (F'_v, F'_p) governs the smoothness of (φ, h) .

We shall study the regularity of a *purely formal* map $(z', c') = (\varphi(z), h(c))$, namely $\varphi(z) \in \mathbb{K}[[z]]^{n+m}$ and $h(c) \in \mathbb{K}[[c]]^{p+m}$, assuming (\mathcal{E}) and (\mathcal{E}') to be analytic. Concretely, the assumption that (φ, h) maps \mathcal{M} to \mathcal{M}' reads as one of the four equivalent identities:

$$(11.5) \quad \begin{cases} \psi(x, \Pi(x, c)) \equiv \Pi'(\phi(x, \Pi(x, c)), h(c)), \\ \psi(z) \equiv \Pi'(\phi(z), h(a, \Pi^*(a, z))), \\ g(a, \Pi^*(a, z)) \equiv \Pi'^*(f(a, \Pi^*(a, z)), \varphi(z)), \\ g(c) \equiv \Pi'^*(f(c), \varphi(x, \Pi(x, c))), \end{cases}$$

in $\mathbb{K}[[x, c]]^m$ and in $\mathbb{K}[[a, z]]^m$.

Theorem 11.6. *Let $(\varphi, h) := \mathcal{M} \rightarrow \mathcal{M}'$ be a purely formal equivalence between two local \mathbb{K} -analytic submanifolds of solutions. Assume that the fundamental pair of foliations (F_ν, F_ρ) is covering at the origin, with type (μ, μ^*) at the origin. Assume that \mathcal{M}' is both κ -solvable with respect to the parameters and κ^* -solvable with respect to the variables. Set $\ell := \mu^*(\kappa + \kappa^*)$ and $\ell^* := \mu(\kappa^* + \kappa)$. Then there exist two \mathbb{K}^{n+m} -valued and \mathbb{K}^{p+m} -valued local \mathbb{K} -analytic maps Φ_ℓ and H_{ℓ^*} , constructible only by means of Π, Π^*, Π', Π'^* , such that the following two formal power series identities hold:*

$$(11.7) \quad \begin{cases} \varphi(z) \equiv \Phi_\ell(z, J_z^\ell \varphi(0)), \\ h(c) \equiv H_{\ell^*}(c, J_c^{\ell^*} h(0)), \end{cases}$$

in $\mathbb{K}[[z]]^{n+m}$ and in $\mathbb{K}[[c]]^{p+m}$, where $J_z^\ell \varphi(0)$ denotes the ℓ -th jet of φ at the origin and similarly for $J_c^{\ell^*} h(0)$. In particular, as a corollary, we have the following two automatic regularity properties:

- $\varphi(z) \in \mathbb{K}\{z\}^{n+m}$ and $h(c) \in \mathbb{K}\{c\}^{p+m}$ are in fact convergent;
- if in addition \mathcal{M} and \mathcal{M}' are \mathbb{K} -algebraic in the sense of Nash, then Φ_ℓ and H_{ℓ^*} are also \mathbb{K} -algebraic, whence $\varphi(z) \in \mathcal{A}_{\mathbb{K}}\{z\}^{n+m}$ and $h(c) \in \mathcal{A}_{\mathbb{K}}\{c\}^{p+m}$ are in fact \mathbb{K} -algebraic.

Proof. We remind the explicit expressions of the two collections of vector fields spanning the leaves of the two foliations F_ν and F_ρ :

$$(11.8) \quad \begin{cases} L_k := \frac{\partial}{\partial x_k} + \sum_{j=1}^m \frac{\partial \Pi^j}{\partial x_k}(x, c) \frac{\partial}{\partial y^j}, & k = 1, \dots, n, \\ L_q^* := \frac{\partial}{\partial a^q} + \sum_{j=1}^m \frac{\partial \Pi^{*j}}{\partial a^q}(a, z) \frac{\partial}{\partial b^j}, & q = 1, \dots, p. \end{cases}$$

Observe that differentiating the first line of (11.5) with respect to x^k amounts to applying the derivation L_k . Similarly, differentiating the third line of (11.5) with respect to a^q amounts to applying L_q^* . We thus get for $(z, c) \in \mathcal{M}$

$$(11.9) \quad \begin{cases} L_k \psi(z) = \sum_{l=1}^n \frac{\partial \Pi^l}{\partial x^k}(\phi(z), h(c)) L_l \phi^l(z) & \text{and} \\ L_q^* g(c) = \sum_{r=1}^p \frac{\partial \Pi^{*r}}{\partial a^q}(f(c), \varphi(z)) L_r^* f^r(c), \end{cases}$$

It follows from $\det \left(\frac{\partial \varphi^k}{\partial z^l} \right)(0) \neq 0$ and $\det \left(\frac{\partial h^k}{\partial c^l} \right)(0) \neq 0$ that the two formal determinants

$$(11.10) \quad \det (L_k \phi^l(z))_{\substack{1 \leq l \leq n \\ 1 \leq k \leq n}}^{1 \leq l \leq n} \quad \text{and} \quad \det (L_q^* f^r(c))_{\substack{1 \leq r \leq p \\ 1 \leq q \leq p}}^{1 \leq r \leq p}$$

have nonvanishing constant term. Consequently, these two matrices are invertible in $\mathbb{K}[[z]]$ and in $\mathbb{K}[[c]]$. So there exist universal polynomials S_l^j and S_r^{*j} such that

$$(11.11) \quad \begin{cases} \frac{\partial \Pi^j}{\partial x^l}(\varphi(z), h(c)) = \frac{S_l^j \left(\{L_{k'} \varphi^{i'}(z)\}_{\substack{1 \leq i' \leq n+m \\ 1 \leq k' \leq n}} \right)}{\det (L_{k'} \phi^{l'}(z))_{\substack{1 \leq l' \leq n \\ 1 \leq k' \leq n}}^{1 \leq l' \leq n}} & \text{and} \\ \frac{\partial \Pi^{*j}}{\partial a^{r'}}(f(c), \varphi(z)) = \frac{S_r^{*j} \left(\{L_{q'}^* h^{i'}(c)\}_{\substack{1 \leq i' \leq p+m \\ 1 \leq q' \leq p}} \right)}{\det (L_{q'}^* f^{r'}(c))_{\substack{1 \leq r' \leq p \\ 1 \leq q' \leq p}}^{1 \leq r' \leq p}}, \end{cases}$$

for $1 \leq j \leq m$, for $1 \leq l \leq n$, for $1 \leq r \leq p$ and for $(z, c) \in \mathcal{M}$.

Again, we apply the vector fields L_k to the obtained first line and the vector fields L_q^* to the obtained second line, getting, thanks to the chain rule:

(11.12)

$$\left\{ \begin{array}{l} \sum_{l_2=1}^n \frac{\partial^2 \Pi^j}{\partial x'^{l_1} x'^{l_2}} (\phi(z), h(c)) L_k \phi^{l_2}(z) = \frac{R_{l_1, k}^j \left(\{L_{k'_1} L_{k'_2} \varphi^{i'}(z)\}_{1 \leq i' \leq n+m}^{1 \leq k'_1, k'_2 \leq n} \right)}{\left[\det(L_{k'} \phi^{l'}(z))_{1 \leq k' \leq n}^{1 \leq l' \leq n} \right]^2} \quad \text{and} \\ \sum_{r_2=1}^p \frac{\partial^2 \Pi^{*j}}{\partial a'^{r_1} a'^{r_2}} (f(c), \varphi(z)) L_q^* f^{r_2}(c) = \frac{R_{r_1, q}^{*j} \left(\{L_{q'_1}^* L_{q'_2}^* h^{i'}(c)\}_{1 \leq i' \leq p+m}^{1 \leq q'_1, q'_2 \leq p} \right)}{\left[\det(L_{q'}^* f^{r'}(c))_{1 \leq q' \leq p}^{1 \leq r' \leq p} \right]^2}, \end{array} \right.$$

for $1 \leq j \leq m$, for $1 \leq l_1, l_2 \leq n$, for $1 \leq r_1, r_2 \leq p$ and for $(z, c) \in \mathcal{M}$. Here, $R_{l_1, k}^j$ and $R_{r_1, q}^{*j}$ are universal polynomials. Then applying once more Cramer's rule, we get

$$(11.13) \quad \left\{ \begin{array}{l} \frac{\partial^2 \Pi^j}{\partial x'^{l_1} x'^{l_2}} (\phi(z), h(c)) = \frac{S_{l_1, l_2}^j \left(\{L_{k'_1} L_{k'_2} \varphi^{i'}(z)\}_{1 \leq i' \leq n+m}^{1 \leq k'_1, k'_2 \leq n} \right)}{\left[\det(L_{k'} \phi^{l'}(z))_{1 \leq k' \leq n}^{1 \leq l' \leq n} \right]^3} \quad \text{and} \\ \frac{\partial^2 \Pi^{*j}}{\partial a'^{r_1} a'^{r_2}} (f(c), \varphi(z)) = \frac{S_{r_1, r_2}^{*j} \left(\{L_{q'_1}^* L_{q'_2}^* h^{i'}(c)\}_{1 \leq i' \leq p+m}^{1 \leq q'_1, q'_2 \leq p} \right)}{\left[\det(L_{q'}^* f^{r'}(c))_{1 \leq q' \leq p}^{1 \leq r' \leq p} \right]^3}. \end{array} \right.$$

By induction, for every j with $1 \leq j \leq m$ and every two multiindices $\beta \in \mathbb{N}^n$ and $\delta \in \mathbb{N}^p$, there exists two universal polynomials S_β^j and S_δ^{*j} such that

$$(11.14) \quad \left\{ \begin{array}{l} \frac{\partial^{|\beta|} \Pi^j}{\partial x'^{\beta}} (\phi(z), h(c)) = \frac{S_\beta^j \left(\{L^{\beta'} \varphi^{i'}(z)\}_{|\beta'| \leq |\beta|}^{1 \leq i' \leq n+m} \right)}{\left[\det(L_{k'} \phi^{l'}(z))_{1 \leq k' \leq n}^{1 \leq l' \leq n} \right]^{2|\beta|+1}} \quad \text{and} \\ \frac{\partial^{|\delta|} \Pi^{*j}}{\partial a'^{\delta}} (f(c), \varphi(z)) = \frac{S_\delta^{*j} \left(\{L^{*\delta'} h^{i'}(c)\}_{|\delta'| \leq |\delta|}^{1 \leq i' \leq p+m} \right)}{\left[\det(L_{q'}^* f^{r'}(c))_{1 \leq q' \leq p}^{1 \leq r' \leq p} \right]^{2|\delta|+1}}. \end{array} \right.$$

Here, for $\beta' \in \mathbb{N}^n$, we denote by $L^{\beta'}$ the derivation of order $|\beta'|$ defined by $(L_1)^{\beta'_1} \dots (L_n)^{\beta'_n}$. Similarly, for $\delta' \in \mathbb{N}^p$, $L^{*\delta'}$ denotes the derivation of order $|\delta'|$ defined by $(L_1^*)^{\delta'_1} \dots (L_p^*)^{\delta'_p}$.

Next, by the assumption that \mathcal{M}' is solvable with respect to the parameters, there exist integers $j(1), \dots, j(p)$ with $1 \leq j(q) \leq m$ and multiindices $\beta(1), \dots, \beta(p) \in \mathbb{N}^n$ with $|\beta(q)| \geq 1$ and $\max_{1 \leq q \leq p} |\beta(q)| = \kappa$ such that the local \mathbb{K} -analytic map

$$(11.15) \quad \mathbb{K}^{p+m} \ni c' \mapsto \left((\Pi^{j(q)}(0, c'))^{1 \leq q \leq m}, \left(\frac{\partial^{|\beta(q)|} \Pi^{j(q)}}{\partial x'^{\beta(q)}}(0, c') \right)_{1 \leq q \leq p} \right) \in \mathbb{K}^{p+m}$$

is of rank $p+m$ at $c' = 0$. Similarly, by the assumption that \mathcal{M}' is solvable with respect to the variables, there exist integers $j^\sim(1), \dots, j^\sim(n)$ with $1 \leq j^\sim(l) \leq m$ and multiindices $\delta(1), \dots, \delta(p) \in \mathbb{N}^n$ with $|\delta(q)| \geq 1$ and $\max_{1 \leq q \leq p} |\delta(q)| = \kappa^*$ such that the local \mathbb{K} -analytic map

$$(11.16) \quad \mathbb{K}^{n+m} \ni z' \mapsto \left((\Pi^{*j^\sim(l)}(0, z'))^{1 \leq l \leq m}, \left(\frac{\partial^{|\delta(l)|} \Pi^{*j^\sim(l)}}{\partial a'^{\delta(l)}}(0, z') \right)_{1 \leq l \leq n} \right) \in \mathbb{K}^{n+m}$$

is of rank $n + m$ at $z' = 0$. We then consider from the first line of (11.14) only the $(p + m)$ equations written for $(j, 0)$, $(j(q), \beta(q))$ and we solve $h(c)$ by means of the analytic implicit function theorem; also, in the second line of (11.14), we consider the $(n + m)$ equations written for $(j, 0)$, $(j^\sim(l), \delta(l))$ and we solve $\varphi(z)$. We get:

$$(11.17) \quad \left\{ \begin{array}{l} h(c) = \widehat{H} \left(\phi(z), \frac{S_{\beta(1)}^{j(1)} \left(\left\{ L^{\beta'} \varphi^{i'}(z) \right\}_{\substack{1 \leq i' \leq n+m \\ |\beta'| \leq |\beta(1)|}} \right)}{\det \left[\left(L_{k'} \phi^{l'}(z) \right)_{\substack{1 \leq l' \leq n \\ 1 \leq k' \leq n}} \right]} \right)^{2|\beta(1)|+1}, \dots \\ \dots, \frac{S_{\beta(p)}^{j(p)} \left(\left\{ L^{\beta'} \varphi^{i'}(z) \right\}_{\substack{1 \leq i' \leq n+m \\ |\beta'| \leq |\beta(p)|}} \right)}{\det \left[\left(L_{k'} \phi^{l'}(z) \right)_{\substack{1 \leq l' \leq n \\ 1 \leq k' \leq n}} \right]} \right)^{2|\beta(p)|+1} \\ \varphi(z) = \widehat{\Phi} \left(f(c), \frac{S_{\delta(1)}^{*j^\sim(1)} \left(\left\{ L^{*\delta'} h^{i'}(c) \right\}_{\substack{1 \leq i' \leq p+m \\ |\delta'| \leq |\delta(1)|}} \right)}{\left[\det \left(L_{q'}^* f^{r'}(c) \right)_{\substack{1 \leq r' \leq p \\ 1 \leq q' \leq p}} \right]} \right)^{2|\delta(1)|+1}, \dots \\ \dots, \frac{S_{\delta(n)}^{*j^\sim(n)} \left(\left\{ L^{*\delta'} h^{i'}(c) \right\}_{\substack{1 \leq i' \leq p+m \\ |\delta'| \leq |\delta(n)|}} \right)}{\left[\det \left(L_{q'}^* f^{r'}(c) \right)_{\substack{1 \leq r' \leq p \\ 1 \leq q' \leq p}} \right]} \right)^{2|\delta(n)|+1} \end{array} \right.$$

for $(z, c) \in \mathcal{M}$. The maps \widehat{H} and $\widehat{\Phi}$ depend only on Π' , Π'^* .

Lemma 11.18. *For every $\beta' \in \mathbb{N}^n$, there exists a universal polynomial $P_{\beta'}$ in the jet variables $J_z^{|\beta'|}$ having \mathbb{K} -analytic coefficients in (z, c) which depends only on Π , Π^* such that, for $i' = 1, \dots, n + m$:*

$$(11.19) \quad L^{\beta'} \varphi^{i'}(z) \equiv P_{\beta'} \left(z, c, J_z^{|\beta'|} \varphi^{i'}(z) \right).$$

A similar property holds for $L^{*\delta'} h^{i'}(c)$.

We deduce that there exist two local \mathbb{K} -analytic mappings Φ_0^0 and H_0^0 such that we can write

$$(11.20) \quad \begin{cases} \varphi(z) = \Phi_0^0(z, c, J_c^{\kappa^*} h(c)), \\ h(c) = H_0^0(z, c, J_z^{\kappa} \varphi(z)), \end{cases}$$

for $(z, c) \in \mathcal{M}$. Concretely, this means that we have two equivalent pairs of formal identities

$$(11.21) \quad \begin{aligned} \varphi(z) &\equiv \Phi_0^0(z, a, \Pi^*(a, z), J_c^{\kappa^*} h(a, \Pi^*(a, z))) \\ \varphi(x, \Pi(x, c)) &\equiv \Phi_0^0(x, \Pi(x, c), c, J_c^{\kappa^*} h(c)) \\ h(c) &\equiv H_0^0(x, \Pi(x, c), c, J_z^{\kappa} \varphi(x, \Pi(x, c))) \\ h(a, \Pi^*(a, z)) &\equiv H_0^0(z, a, \Pi^*(a, z), J_z^{\kappa} \varphi(z)) \end{aligned}$$

in $\mathbb{K}[[a, z]]^{n+m}$ and in $\mathbb{K}[[x, c]]^{p+m}$. We notice that, whereas φ and h are *a priori* only purely formal, by construction, Φ_0^0 and H_0^0 are \mathbb{K} -analytic near $(0, 0, J_c^{\kappa^*} h(0))$ and near $(0, 0, J_z^{\kappa} \varphi(0))$.

Next, we introduce the following vector fields with \mathbb{K} -analytic coefficients tangent to \mathcal{M} :

$$(11.22) \quad \begin{cases} V_j := \frac{\partial}{\partial y^j} + \sum_{l=1}^m \frac{\partial \Pi^{*l}}{\partial y^j}(a, z) \frac{\partial}{\partial b^l}, & j = 1, \dots, m \quad \text{and} \\ V_j^* := \frac{\partial}{\partial b^j} + \sum_{l=1}^m \frac{\partial \Pi^l}{\partial b^j}(x, c) \frac{\partial}{\partial y^l}, & j = 1, \dots, m. \end{cases}$$

Indeed, we check that $V_{j_1}[b^{j_2} - \Pi^{*j_2}(a, z)] \equiv 0$ and that $V_{j_1}^*[y^{j_2} - \Pi^{j_2}(x, c)] \equiv 0$.

For $\delta' \in \mathbb{N}^m$, we observe that $V^{\delta'}\varphi = \frac{\partial^{|\delta'|}\varphi}{\partial y^{\delta'}}$. Applying then $L^{\beta'}$ with $\beta' \in \mathbb{N}^n$, we get for $i = 1, \dots, n + m$:

$$(11.23) \quad L^{\beta'}V^{\delta'}\varphi^i(z) = Q_{\beta', \delta'}(z, c, J_z^{|\beta'|+|\delta'|}\varphi^i(z)),$$

with $Q_{\beta', \delta'}$ universal. Since the $n + m$ vector fields L_k and V_j , having coefficients depending on (z, c) , span the tangent space to $\mathbb{K}_x^n \times \mathbb{K}_y^m$, the change of basis of derivations yields, by induction, the following.

Lemma 11.24. *For every $\alpha \in \mathbb{N}^{n+m}$, there exists a universal polynomial P_α in its last variables with coefficients being \mathbb{K} -analytic in (z, c) and depending only on Π, Π^* such that, for $i = 1, \dots, n + m$:*

$$(11.25) \quad \partial_z^\alpha \varphi^i(z) \equiv P_\alpha\left(z, c, (L^{\beta'}V^{\delta'}\varphi^i(z))_{|\beta'|+|\delta'| \leq |\alpha|}\right).$$

We are now in position to state and to prove the first fundamental technical lemma which generalizes the two formulas (11.20) to arbitrary jets.

Lemma 11.26. *For every $\lambda \in \mathbb{N}$, there exist two local \mathbb{K} -analytic maps, Φ_0^λ valued in $\mathbb{K}^{(n+m)C_{n+m+\lambda}^\lambda}$, and H_0^λ valued in $\mathbb{K}^{(p+m)C_{p+m+\lambda}^\lambda}$, such that:*

$$(11.27) \quad \begin{cases} J_z^\lambda \varphi(z) \equiv \Phi_0^\lambda(z, c, J_c^{\kappa^*+\lambda} h(c)), \\ J_c^\lambda h(c) \equiv H_0^\lambda(z, c, J_z^{\kappa+\lambda} \varphi(z)). \end{cases}$$

Proof. Consider for instance the first line. To obtain it, it suffices to apply the derivations $L^{\beta'}V^{\delta'}$ with $|\beta'| + |\delta'| \leq \lambda$ to the first line of (11.20), to use the chain rule and to apply Lemma 11.24. \square

Let $\theta \in \mathbb{K}^l$, $l \in \mathbb{N}$, let $Q(\theta) = (Q_1(\theta), \dots, Q_{n+2m+p}(\theta)) \in \mathbb{K}[[\theta]]^{n+2m+p}$ and let $a_1 \in \mathbb{K}^p$. As the multiple flow of L^* given by (10.10) does not act on the variables (x, y) , we have the trivial but crucial property:

$$(11.28) \quad \varphi(L_{a_1}^*(Q(\theta))) \equiv \varphi(\pi_z(L_{a_1}^*(Q(\theta)))) \equiv \varphi(\pi_z(Q(\theta))) \equiv \varphi(Q(\theta)).$$

At the end, we allow to suppress the projection π_z : this slight abuse of notation will lighten slightly the writing of further formulas. More generally, for $\lambda \in \mathbb{N}$, $a_1 \in \mathbb{K}^p$, $x_1 \in \mathbb{K}^n$:

$$(11.29) \quad \begin{aligned} J_z^\lambda \varphi(L_{a_1}^*(Q(\theta))) &\equiv J_z^\lambda \varphi(Q(\theta)) \quad \text{and} \\ J_c^\lambda h(L_{x_1}(Q(\theta))) &\equiv J_c^\lambda h(Q(\theta)). \end{aligned}$$

As a consequence, for $2k$ even and for $2k+1$ odd, we have the following four cancellation relations, useful below (we drop π_z and π_c after $J_z^\lambda \varphi$ and after $J_c^\lambda h$):

$$(11.30) \quad \begin{cases} J_z^\lambda \varphi(\Gamma_{2k}([xa]_{2k})) \equiv J_z^\lambda \varphi(\Gamma_{2k-1}([xa]_{2k-1})), \\ J_c^\lambda h(\Gamma_{2k}^*([ax]_{2k})) \equiv J_c^\lambda h(\Gamma_{2k-1}^*([ax]_{2k-1})), \\ J_z^\lambda \varphi(\Gamma_{2k+1}^*([ax]_{2k+1})) \equiv J_z^\lambda \varphi(\Gamma_{2k}^*([ax]_{2k})), \\ J_c^\lambda h(\Gamma_{2k+1}([xa]_{2k+1})) \equiv J_c^\lambda h(\Gamma_{2k}([xa]_{2k})). \end{cases}$$

We are now in position to state and to prove the second main technical proposition.

Proposition 11.31. *For every even chain-length $2k$ and for every jet-height λ , there exist two local \mathbb{K} -analytic maps, Φ_{2k}^λ valued in $\mathbb{K}^{(n+m)C_{n+m+\lambda}^\lambda}$, and H_{2k}^λ valued in $\mathbb{K}^{(p+m)C_{p+m+\lambda}^\lambda}$ such that:*

$$(11.32) \quad \begin{cases} J_z^\lambda \varphi(\Gamma_{2k}^*([ax]_{2k})) \equiv \Phi_{2k}^\lambda([ax]_{2k}, J_z^{k(\kappa+\kappa^*)+\lambda} \varphi(0)) \quad \text{and} \\ J_c^\lambda h(\Gamma_{2k}([xa]_{2k})) \equiv H_{2k}^\lambda([xa]_{2k}, J_c^{k(\kappa+\kappa^*)+\lambda} \varphi(0)). \end{cases}$$

Similarly, for every odd chain length $2k+1$ and for every jet height λ , there exist two local \mathbb{K} -analytic maps, Φ_{2k+1}^λ valued in $\mathbb{K}^{(n+m)C_{n+m+\lambda}^\lambda}$ and H_{2k+1}^λ valued in $\mathbb{K}^{(p+m)C_{p+m+\lambda}^\lambda}$, such that:

$$(11.33) \quad \begin{cases} J_z^\lambda \varphi(\Gamma_{2k+1}([xa]_{2k+1})) \equiv \Phi_{2k+1}^\lambda([xa]_{2k+1}, J_c^{k\kappa+(k+1)\kappa^*+\lambda} h(0)), \\ J_c^\lambda h(\Gamma_{2k+1}^*([ax]_{2k+1})) \equiv H_{2k+1}^\lambda([ax]_{2k+1}, J_z^{(k+1)\kappa+k\kappa^*+\lambda} \varphi(0)). \end{cases}$$

These maps depend only on Π, Π^*, Π', Π'^* .

Proof. For $2k+1=1$, we replace (z, c) by $\Gamma_1([xa]_1)$ in the first line of (11.27) and by $\Gamma_1^*([ax]_1)$ in the second line. Taking crucially account of the cancellation properties (11.29), we get:

$$(11.34) \quad \begin{cases} J_z^\lambda \varphi(\Gamma_1([xa]_1)) \equiv \Phi_0^\lambda(\Gamma_1([xa]_1), J_c^{\kappa^*+\lambda} h(\Gamma_1([xa]_1))) \\ \quad \equiv \Phi_0^\lambda(\Gamma_1([xa]_1), J_c^{\kappa^*+\lambda} h(0)) \\ \quad =: \Phi_1^\lambda([xa]_1, J_c^{\kappa^*+\lambda} h(0)), \\ J_c^\lambda h(\Gamma_1^*([ax]_1)) \equiv H_0^\lambda(\Gamma_1^*([ax]_1), J_z^{\kappa+\lambda} \varphi(\Gamma_1^*([ax]_1))) \\ \quad \equiv H_0^\lambda(\Gamma_1^*([ax]_1), J_z^{\kappa+\lambda} \varphi(0)) \\ \quad =: H_1^\lambda([ax]_1, J_z^{\kappa+\lambda} \varphi(0)). \end{cases}$$

Here, the third line defines Φ_1^λ and the sixth line defines H_1^λ . Thus, the proposition holds for $2k+1=1$.

The rest of the proof proceeds by induction. We treat only the induction step from an odd chain-length $2k+1$ to an even chain-length $2k+2$, the other induction step being similar.

To this aim, we replace the variables (z, c) in the first line of (11.27) by $\Gamma_{2k+2}^*([ax]_{2k+2})$. Taking account of the cancellation property and of the induction assumption:

$$(11.35) \quad \begin{aligned} J_z^\lambda \varphi(\Gamma_{2k+2}^*([ax]_{2k+2})) &\equiv \Phi_0^\lambda(\Gamma_{2k+2}^*([ax]_{2k+2}), J_c^{\kappa^*+\lambda} h(\Gamma_{2k+2}^*([ax]_{2k+2}))) \\ &\equiv \Phi_0^\lambda(\Gamma_{2k+2}^*([ax]_{2k+2}), J_c^{\kappa^*+\lambda} h(\Gamma_{2k+1}^*([ax]_{2k+1}))) \\ &\equiv \Phi_0^\lambda(\Gamma_{2k+2}^*([ax]_{2k+2}), H_{2k+1}^{\kappa^*+\lambda}([ax]_{2k+1}, J_c^{(k+1)(\kappa+\kappa^*)+\lambda} \varphi(0))) \\ &=: \Phi_{2k+2}^\lambda([ax]_{2k+2}, J_c^{(k+1)(\kappa+\kappa^*)+\lambda} \varphi(0)), \end{aligned}$$

The last line defines Φ_{2k+2}^λ . Similarly, we replace (z, c) in the second line of (11.27) by $\Gamma_{2k+2}([xa]_{2k+2})$. Taking account of the cancellation property and of the induction assumption:

(11.36)

$$\begin{aligned} J_c^\lambda h(\Gamma_{2k+2}([xa]_{2k+2})) &\equiv H_0^\lambda \left(\Gamma_{2k+2}([xa]_{2k+2}), J_c^{\kappa+\lambda} \varphi(\Gamma_{2k+2}([xa]_{2k+2})) \right) \\ &\equiv H_0^\lambda \left(\Gamma_{2k+2}([xa]_{2k+2}), J_c^{\kappa+\lambda} \varphi(\Gamma_{2k+1}([xa]_{2k+1})) \right) \\ &\equiv H_0^\lambda \left(\Gamma_{2k+2}([xa]_{2k+2}), \Phi_{2k+1}^{\kappa+\lambda} \left([xa]_{2k+1}, J_c^{(k+1)(\kappa+\kappa^*)+\lambda} h(0) \right) \right) \\ &=: H_{2k+2}^\lambda \left([xa]_{2k+2}, J_c^{(k+1)(\kappa+\kappa^*)+\lambda} h(0) \right). \end{aligned}$$

This completes the proof. \square

End of the proof of Theorem 11.6. With (μ, μ^*) being the type of (F_v, F_p) and with $[ax]_{2\mu^*}^0$ given by Corollary 10.31, the rank property (10.32) insures the existence of an affine $(n+m)$ -dimensional space $H \subset \mathbb{K}^{\mu^*(p+n)}$ passing through $[ax]_{2\mu^*}^0$ and equipped with a local parametrization

$$(11.37) \quad \mathbb{K}^{n+m} \ni s \mapsto [ax]_{2\mu^*}(s) \in H$$

satisfying $[ax]_{2\mu^*}(0) = [ax]_{2\mu^*}^0$, such that the map

$$(11.38) \quad \mathbb{K}^{n+m} \ni s \mapsto \pi_z \left(\Gamma_{2\mu^*}^*([ax]_{2\mu^*}(s)) \right) =: z(s) \in \mathbb{K}^{n+m}$$

is a local diffeomorphism fixing $0 \in \mathbb{K}^{n+m}$. Replacing z by $z(s)$ in $\varphi(z)$ and applying the formula in the first line of (11.32) with $\lambda = 0$ and with $k = 2\mu^*$, we obtain

$$\begin{aligned} (11.39) \quad \varphi(z(s)) &= \varphi \left(\pi_z \left(\Gamma_{2\mu^*}^*([ax]_{2\mu^*}(s)) \right) \right) \\ &= \varphi \left(\Gamma_{2\mu^*}^*([ax]_{2\mu^*}(s)) \right) \\ &\equiv \Phi_{2\mu^*}^0 \left([ax]_{2\mu^*}(s), J_z^{\mu^*(\kappa+\kappa^*)} \varphi(0) \right). \end{aligned}$$

Inverting $s \mapsto z = z(s)$ as $z \mapsto s = s(z)$, we finally get

$$(11.40) \quad \begin{aligned} \varphi(z) &= \varphi(z(s(z))) \equiv \Phi_{2\mu^*}^0 \left([ax]_{2\mu^*}(s(z)), J_z^{\mu^*(\kappa+\kappa^*)} \varphi(0) \right) \\ &=: \Phi_\ell \left(z, J_z^{\mu^*(\kappa+\kappa^*)} \varphi(0) \right), \end{aligned}$$

with $\ell := \mu^*(\kappa + \kappa^*)$, where the last line defines Φ_ℓ . In conclusion, we have derived the first line of (11.7). The second one is obtained similarly.

If Π, Π^*, Π', Π'^* are algebraic, so are $\Gamma_k, \Gamma_k^*, \widehat{H}, \widehat{\Phi}, \Phi_0^\lambda, H_0^\lambda, \Phi_k^\lambda, H_k^\lambda$ and Φ_ℓ, H_{ℓ^*} .

The proof of Theorem 11.6 is complete. \square

II: Explicit prolongations of infinitesimal Lie symmetries

Table of contents

1. Jet spaces and prolongations 57.
 2. One independent variable and one dependent variable 62.
 3. Several independent variables and one dependent variable 68.
 4. One independent variable and several dependent variables 84.
 5. Several independent variables and several dependent variables 90.

§1. JET SPACES AND PROLONGATIONS

1.1. Choice of notations for the jet space variables. Let $\mathbb{K} = \mathbb{R}$ or \mathbb{C} . Let $n \geq 1$ and $m \geq 1$ be two positive integers and consider two sets of variables $x = (x^1, \dots, x^n) \in \mathbb{K}^n$ and $y = (y^1, \dots, y^m)$. In the classical theory of Lie symmetries of partial differential equations, one considers certain differential systems whose (local) solutions should be mappings of the form $y = y(x)$. We refer to [Ol1986] and to [BK1989] for an exposition of the fundamentals of the theory. Accordingly, the variables x are usually called *independent*, whereas the variables y are called *dependent*. Not to enter in subtle regularity considerations (as in [Me2005b]), we shall assume C^∞ -smoothness of all functions throughout this paper.

Let $\kappa \geq 1$ be a positive integer. For us, in a very concrete way (without fiber bundles), the κ -th jet space $\mathcal{J}_{n,m}^\kappa$ consists of the space $\mathbb{K}^{n+m+m \frac{(n+m)!}{n! m!}}$ equipped with the affine coordinates

$$(1.2) \quad (x^i, y^j, y_{i_1}^j, y_{i_1, i_2}^j, \dots, y_{i_1, i_2, \dots, i_\kappa}^j),$$

having the symmetries

$$(1.3) \quad y_{i_1, i_2, \dots, i_\lambda}^j = y_{i_{\sigma(1)}, i_{\sigma(2)}, \dots, i_{\sigma(\lambda)}}^j,$$

for every λ with $1 \leq \lambda \leq \kappa$ and for every permutation σ of the set $\{1, \dots, \lambda\}$. The variable $y_{i_1, i_2, \dots, i_\lambda}^j$ is an independent coordinate corresponding to the λ -th partial derivative $\frac{\partial^\lambda y^j}{\partial x^{i_1} \partial x^{i_2} \dots \partial x^{i_\lambda}}$. So the symmetries (1.3) are natural.

In the classical Lie theory ([Ol1979], [Ol1986], [BK1989]), all the geometric objects: point transformations, vector fields, *etc.*, are local, defined in a neighborhood of some point lying in some affine space \mathbb{K}^N . However, in this paper, the original geometric motivations are rapidly forgotten in order to focus on combinatorial considerations. Thus, to simplify the presentation, we shall not introduce any special notation to speak of certain local open subsets of \mathbb{K}^{n+m} , or of the jet space $\mathcal{J}_{n,m}^\kappa = \mathbb{K}^{n+m+m \frac{(n+m)!}{n! m!}}$, *etc.*: we will always work in global affine spaces \mathbb{K}^N .

1.4. Prolongation $\varphi^{(\kappa)}$ of a local diffeomorphism φ to the κ -th jet space. In this paragraph, we recall how the prolongation of a diffeomorphism to the κ -th jet space is defined ([Ol1979], [Ol1986], [BK1989]).

Let $x_* \in \mathbb{K}^n$ be a central fixed point and let $\varphi : \mathbb{K}^{n+m} \rightarrow \mathbb{K}^{n+m}$ be a diffeomorphism whose Jacobian matrix is close to the identity matrix, at least in a small neighborhood of x_* . Let

$$(1.5) \quad J_{x_*}^\kappa := (x_*^i, y_{*i_1}^j, y_{*i_1, i_2}^j, \dots, y_{*i_1, i_2, \dots, i_\kappa}^j) \in \mathcal{J}_{n,m}^\kappa|_{x_*}$$

be an arbitrary κ -jet based at x_* . The goal is to define its transformation $\varphi^{(\kappa)}(J_{x_*}^\kappa)$ by φ .

To this aim, choose an arbitrary mapping $\mathbb{K}^n \ni x \mapsto g(x) \in \mathbb{K}^m$ defined at least in a neighborhood of x_* and representing this κ -th jet, *i.e.* satisfying

$$(1.6) \quad y_{*i_1, \dots, i_\lambda}^j = \frac{\partial^\lambda g^j}{\partial x^{i_1} \dots \partial x^{i_\lambda}}(x_*),$$

for every $\lambda \in \mathbb{N}$ with $0 \leq \lambda \leq \kappa$, for all indices i_1, \dots, i_λ with $1 \leq i_1, \dots, i_\lambda \leq n$ and for every $j \in \mathbb{N}$ with $1 \leq j \leq m$. In accordance with the splitting $(x, y) \in \mathbb{K}^n \times \mathbb{K}^m$ of coordinates, split the components of the diffeomorphism φ as $\varphi = (\phi, \psi) \in \mathbb{K}^n \times \mathbb{K}^m$. Write (\bar{x}, \bar{y}) the coordinates in the target space, so that the diffeomorphism φ is:

$$(1.7) \quad \mathbb{K}^{n+m} \ni (x, y) \longmapsto (\bar{x}, \bar{y}) = (\phi(x, y), \psi(x, y)) \in \mathbb{K}^{n+m}.$$

Restrict the variables (x, y) to belong to the graph of g , namely put $y := g(x)$ above, which yields

$$(1.8) \quad \begin{cases} \bar{x} = \phi(x, g(x)), \\ \bar{y} = \psi(x, g(x)). \end{cases}$$

As the differential of φ at x_* is close to the identity, the first family of n scalar equations may be solved with respect to x , by means of the implicit function theorem. Denote $x = \bar{\chi}(\bar{x})$ the resulting mapping, satisfying by definition

$$(1.9) \quad \bar{x} \equiv \phi(\bar{\chi}(\bar{x}), g(\bar{\chi}(\bar{x}))).$$

Replace x by $\bar{\chi}(\bar{x})$ in the second family of m scalar equations (1.8) above, which yields:

$$(1.10) \quad \bar{y} = \psi(\bar{\chi}(\bar{x}), g(\bar{\chi}(\bar{x}))).$$

Denote simply by $\bar{y} = \bar{g}(\bar{x})$ this last relation, where $\bar{g}(\cdot) := \psi(\bar{\chi}(\cdot), g(\bar{\chi}(\cdot)))$.

In summary, the graph $y = g(x)$ has been transformed to the graph $\bar{y} = \bar{g}(\bar{x})$ by the diffeomorphism φ .

Define then the *transformed jet* $\varphi^{(\kappa)}(J_{x_*}^\kappa)$ to be the κ -th jet of \bar{g} at the point $\bar{x}_* := \phi(x_*)$, namely:

$$(1.11) \quad \varphi^{(\kappa)}(J_{x_*}^\kappa) := \left(\frac{\partial^\lambda \bar{g}^j}{\partial \bar{x}^{i_1} \dots \partial \bar{x}^{i_\lambda}}(\bar{x}_*) \right)_{\substack{1 \leq j \leq m \\ 1 \leq i_1, \dots, i_\lambda \leq n, 0 \leq \lambda \leq \kappa}} \in \mathcal{J}_{n,m}^\kappa \Big|_{\bar{x}_*}.$$

It may be shown that this jet does not depend on the choice of a local graph $y = g(x)$ representing the κ -th jet $J_{x_*}^\kappa$ at x_* . Furthermore, if $\pi_\kappa := \mathcal{J}_{n,m}^\kappa \rightarrow \mathbb{K}^m$ denotes the canonical projection onto the first factor, the following diagram commutes:

$$\begin{array}{ccc} \mathcal{J}_{n,m}^\kappa & \xrightarrow{\varphi^{(\kappa)}} & \mathcal{J}_{n,m}^\kappa \\ \pi_\kappa \downarrow & & \downarrow \pi_\kappa \\ \mathbb{K}^{n+m} & \xrightarrow{\varphi} & \mathbb{K}^{n+m} \end{array}.$$

1.12. Inductive formulas for the κ -th prolongation $\varphi^{(\kappa)}$. To present them, we change our notations. Instead of (\bar{x}, \bar{y}) , as coordinates in the target space $\mathbb{K}^n \times \mathbb{K}^m$, we shall use capital letters:

$$(1.13) \quad (X^1, \dots, X^n, Y^1, \dots, Y^m).$$

In the source space \mathbb{K}^{n+m} equipped with the coordinates (x, y) , we use the jet coordinates (1.2) on the associated κ -th jet space. In the target space \mathbb{K}^{n+m} equipped with the coordinates (X, Y) , we use the coordinates

$$(1.14) \quad (X^i, Y^j, Y_{X^{i_1}}^j, Y_{X^{i_1} X^{i_2}}^j, \dots, Y_{X^{i_1} X^{i_2} \dots X^{i_\kappa}}^j)$$

on the associated κ -th jet space; to avoid confusion with $y_{i_1}, y_{i_1, i_2}, \dots$ in subsequent formulas, we do not write $Y_{i_1}, Y_{i_1, i_2}, \dots$. In these notations, the diffeomorphism φ whose first order approximation is close to the identity mapping in a neighborhood of x_* may be written under the form:

$$(1.15) \quad \varphi : (x^{i'}, y^{j'}) \mapsto (X^i, Y^j) = \left(X^i(x^{i'}, y^{j'}), Y^j(x^{i'}, y^{j'}) \right),$$

for some C^∞ -smooth functions $X^i(x^{i'}, y^{j'})$, $i = 1, \dots, n$, and $Y^j(x^{i'}, y^{j'})$, $j = 1, \dots, m$. The first prolongation $\varphi^{(1)}$ of φ may be written under the form:

$$(1.16) \quad \varphi^{(1)} : (x^{i'}, y^{j'}, y_{i'_1}^{j'}) \mapsto \left(X^i(x^{i'}, y^{j'}), Y^j(x^{i'}, y^{j'}), Y_{X^{i_1}}^j(x^{i'}, y^{j'}, y_{i'_1}^{j'}) \right),$$

for some functions $Y_{X^{i_1}}^j(x^{i'}, y^{j'}, y_{i'_1}^{j'})$ which depend on the pure first jet variables $y_{i'_1}^{j'}$. The way how these functions depend on the first order partial derivatives functions $X_{x^{i'}}^i, X_{y^{j'}}^i, Y_{x^{i'}}^j, Y_{y^{j'}}^j$ and on the pure first jet variables $y_{i'_1}^{j'}$ is provided (in principle) by the following compact formulas ([BK1989]):

$$(1.17) \quad \begin{pmatrix} Y_{X^1}^j \\ \vdots \\ Y_{X^n}^j \end{pmatrix} = \begin{pmatrix} D_1^1 X^1 & \dots & D_1^1 X^n \\ \vdots & \dots & \vdots \\ D_n^1 X^1 & \dots & D_n^1 X^n \end{pmatrix}^{-1} \begin{pmatrix} D_1^1 Y^j \\ \vdots \\ D_n^1 Y^j \end{pmatrix},$$

where, for $i' = 1, \dots, n$, the symbol $D_{i'}^1$ denotes the i' -th first order total differentiation operator:

$$(1.18) \quad D_{i'}^1 := \frac{\partial}{\partial x^{i'}} + \sum_{j'=1}^m y_{i'_1}^{j'} \frac{\partial}{\partial y^{j'}}.$$

Strictly speaking, these formulas (1.17) are not explicit, because an inverse matrix is involved and because the terms $D_{i'}^1 X^i, D_{i'}^1 Y^j$ are not developed. However, it would be feasible and elementary to write down the corresponding totally explicit complete formulas for the functions $Y_{X^{i_1}}^j = Y_{X^{i_1}}^j(x^{i'}, y^{j'}, y_{i'_1}^{j'})$.

Next, the second prolongation $\varphi^{(2)}$ is of the form

$$(1.19) \quad \varphi^{(2)} : (x^{i'}, y^{j'}, y_{i'_1}^{j'}, y_{i'_1, i'_2}^{j'}) \mapsto \left(\varphi^{(1)}(x^{i'}, y^{j'}, y_{i'_1}^{j'}), Y_{X^{i_1} X^{i_2}}^j(x^{i'}, y^{j'}, y_{i'_1}^{j'}, y_{i'_1, i'_2}^{j'}) \right),$$

for some functions $Y_{X^{i_1} X^{i_2}}^j(x^{i'}, y^{j'}, y_{i'_1}^{j'}, y_{i'_1, i'_2}^{j'})$ which depend on the pure first and second jet variables. For $i = 1, \dots, n$, the expressions of $Y_{X^{i_1} X^{i_2}}^j$ are given by the following compact formulas (again [BK1989]):

$$(1.20) \quad \begin{pmatrix} Y_{X^{i_1} X^1}^j \\ \vdots \\ Y_{X^{i_1} X^n}^j \end{pmatrix} = \begin{pmatrix} D_1^1 X^1 & \dots & D_1^1 X^n \\ \vdots & \dots & \vdots \\ D_n^1 X^1 & \dots & D_n^1 X^n \end{pmatrix}^{-1} \begin{pmatrix} D_1^2 Y_{X^{i_1}}^j \\ \vdots \\ D_n^2 Y_{X^{i_1}}^j \end{pmatrix},$$

where, for $i' = 1, \dots, n$, the symbol $D_{i'}^2$ denotes the i' -th second order total differentiation operator:

$$(1.21) \quad D_{i'}^2 := \frac{\partial}{\partial x^{i'}} + \sum_{j'=1}^m y_{i'_1}^{j'} \frac{\partial}{\partial y^{j'}} + \sum_{j'=1}^m \sum_{i'_1=1}^n y_{i'_1, i'_2}^{j'} \frac{\partial}{\partial y_{i'_1}^{j'}}.$$

Again, these formulas (1.20) are not explicit in the sense that an inverse matrix is involved and that the terms $D_{i'}^2 X^i, D_{i'}^2 Y_{X^{i_1}}^j$ are not developed. It would already be a nontrivial

computational task to develop these expressions and to find out some nice satisfying combinatorial formulas.

In order to present the general inductive non-explicit formulas for the computation of the κ -th prolongation $\varphi^{(\kappa)}$, we need some more notation. Let $\lambda \in \mathbb{N}$ be an arbitrary integer. For $i' = 1, \dots, n$, let $D_{i'}^\lambda$ denotes the i' -th λ -th order total differentiation operators, defined precisely by:

$$(1.22) \quad \left\{ \begin{aligned} D_{i'}^\lambda &:= \frac{\partial}{\partial x^{i'}} + \sum_{j'=1}^m y_{i'}^{j'} \frac{\partial}{\partial y^{j'}} + \sum_{j'=1}^m \sum_{i'_1=1}^n y_{i',i'_1}^{j'} \frac{\partial}{\partial y_{i'_1}^{j'}} + \sum_{j'=1}^m \sum_{i'_1,i'_2=1}^n y_{i',i'_1,i'_2}^{j'} \frac{\partial}{\partial y_{i'_1,i'_2}^{j'}} + \\ &+ \cdots + \sum_{j'=1}^m \sum_{i'_1,i'_2,\dots,i'_{\lambda-1}=1}^n y_{i',i'_1,i'_2,\dots,i'_{\lambda-1}}^{j'} \frac{\partial}{\partial y_{i'_1,i'_2,\dots,i'_{\lambda-1}}^{j'}}. \end{aligned} \right.$$

Then, for $i = 1, \dots, n$, the expressions of $Y_{X^{i_1} \dots X^{i_{\lambda-1}} X^i}^j$ are given by the following compact formulas (again [BK1989]):

$$(1.23) \quad \begin{pmatrix} Y_{X^{i_1} \dots X^{i_{\lambda-1}} X^i}^j \\ \vdots \\ Y_{X^{i_1} \dots X^{i_{\lambda-1}} X^n}^j \end{pmatrix} = \begin{pmatrix} D_1^1 X^1 & \cdots & D_1^1 X^n \\ \vdots & \cdots & \vdots \\ D_n^1 X^1 & \cdots & D_n^1 X^n \end{pmatrix}^{-1} \begin{pmatrix} D_1^\lambda Y_{X^{i_1} \dots X^{i_{\lambda-1}}}^j \\ \vdots \\ D_n^\lambda Y_{X^{i_1} \dots X^{i_{\lambda-1}}}^j \end{pmatrix}.$$

Again, these inductive formulas are incomplete and unsatisfactory.

Problem 1.24. Find totally explicit complete formulas for the κ -th prolongation $\varphi^{(\kappa)}$.

Except in the cases $\kappa = 1, 2$, we have not been able to solve this problem. The case $\kappa = 1$ is elementary. Complete formulas in the particular cases $\kappa = 2, n = 1, m \geq 1$ and $n \geq 1, m = 1$ are implicitly provided in [Me2004] and in Section 1.2.4, where one observes the appearance of some modifications of the Jacobian determinant of the diffeomorphism φ , inserted in a clearly understandable combinatorics. In fact, there is a nice dictionary between the formulas for $\varphi^{(2)}$ and the formulas for the second prolongation $\mathcal{L}^{(2)}$ of a vector field \mathcal{L} which were written in equation (43) of [GM2003a] (see also equations (2.6), (3.20), (4.6) and (5.3) in the next paragraphs). In the passage from $\varphi^{(2)}$ to $\mathcal{L}^{(2)}$, a sort of formal first order linearization may be observed and the reverse passage may be easily guessed. However, for $\kappa \geq 3$, the formulas for $\varphi^{(\kappa)}$ explode faster than the formulas for the κ -th prolongation $\mathcal{L}^{(\kappa)}$ of a vector field \mathcal{L} . Also, the dictionary between $\varphi^{(\kappa)}$ and $\mathcal{L}^{(\kappa)}$ disappears. In fact, to elaborate an appropriate dictionary, we believe that one should introduce before a sort of formal $(\kappa - 1)$ -th order linearizations of $\varphi^{(\kappa)}$, finer than the first order linearization $\mathcal{L}^{(\kappa)}$. To be optimistic, we believe that the final answer to Problem 1.24 is, nevertheless, accessible after hard work.

The present article is devoted to present totally explicit complete formulas for the κ -th prolongation $\mathcal{L}^{(\kappa)}$ of a vector field \mathcal{L} to $\mathcal{J}_{n,m}^\kappa$, for $n \geq 1$ arbitrary, for $m \geq 1$ arbitrary and for $\kappa \geq 1$ arbitrary.

1.25. Prolongation of a vector field to the κ -th jet space. Consider a vector field

$$(1.26) \quad \mathcal{L} = \sum_{i=1}^n \mathcal{X}^i(x, y) \frac{\partial}{\partial x^i} + \sum_{j=1}^m \mathcal{Y}^j(x, y) \frac{\partial}{\partial y^j},$$

defined in \mathbb{K}^{n+m} . Its flow:

$$(1.27) \quad \varphi_t(x, y) := \exp(t\mathcal{L})(x, y)$$

§2. ONE INDEPENDENT VARIABLE AND ONE DEPENDENT VARIABLE

2.1. Simplified adapted notations. Assume $n = 1$ and $m = 1$, let $\kappa \in \mathbb{N}$ with $\kappa \geq 1$ and simply denote the jet variables by:

$$(2.2) \quad (x, y, y_1, y_2, \dots, y_\kappa) \in \mathcal{J}_{1,1}^\kappa.$$

The κ -th prolongation of a vector field $\mathcal{L} = \mathcal{X} \frac{\partial}{\partial x} + \mathcal{Y} \frac{\partial}{\partial y}$ will be denoted by:

$$(2.3) \quad \mathcal{L}^{(\kappa)} = \mathcal{X} \frac{\partial}{\partial x} + \mathcal{Y} \frac{\partial}{\partial y} + \mathbf{Y}_1 \frac{\partial}{\partial y_1} + \mathbf{Y}_2 \frac{\partial}{\partial y_2} + \dots + \mathbf{Y}_\kappa \frac{\partial}{\partial y_\kappa}.$$

The coefficients $\mathbf{Y}_1, \mathbf{Y}_2, \dots, \mathbf{Y}_\kappa$ are computed by means of the inductive formulas:

$$(2.4) \quad \begin{cases} \mathbf{Y}_1 := D^1(\mathcal{Y}) - D^1(\mathcal{X}) y_1, \\ \mathbf{Y}_2 := D^2(\mathbf{Y}_1) - D^1(\mathcal{X}) y_2, \\ \dots \\ \mathbf{Y}_\kappa := D^\kappa(\mathbf{Y}_{\kappa-1}) - D^1(\mathcal{X}) y_\kappa, \end{cases}$$

where, for $1 \leq \lambda \leq \kappa$:

$$(2.5) \quad D^\lambda := \frac{\partial}{\partial x} + y_1 \frac{\partial}{\partial y} + y_2 \frac{\partial}{\partial y_1} + \dots + y_\lambda \frac{\partial}{\partial y_{\lambda-1}}.$$

By direct elementary computations, for $\kappa = 1$ and for $\kappa = 2$, we obtain the following two very classical formulas :

$$(2.6) \quad \begin{cases} \mathbf{Y}_1 = \mathcal{Y}_x + [\mathcal{Y}_y - \mathcal{X}_x] y_1 + [-\mathcal{X}_y] (y_1)^2, \\ \mathbf{Y}_2 = \mathcal{Y}_{x^2} + [2 \mathcal{Y}_{xy} - \mathcal{X}_{x^2}] y_1 + [\mathcal{Y}_{y^2} - 2 \mathcal{X}_{xy}] (y_1)^2 + [-\mathcal{X}_{y^2}] (y_1)^3 + \\ \quad + [\mathcal{Y}_y - 2 \mathcal{X}_x] y_2 + [-3 \mathcal{X}_y] y_1 y_2. \end{cases}$$

Our main objective is to *devise the general combinatorics*. Thus, to attain this aim, we have to achieve patiently formal computations of the next coefficients $\mathbf{Y}_3, \mathbf{Y}_4$ and \mathbf{Y}_5 . We systematically use parentheses $[\]$ to single out every coefficient of the polynomials $\mathbf{Y}_3, \mathbf{Y}_4$ and \mathbf{Y}_5 in the pure jet variables y_1, y_2, y_3, y_4 and y_5 , putting every sign inside these parentheses. We always put the monomials in the pure jet variables y_1, y_2, y_3, y_4 and y_5 after the parentheses. For completeness, let us provide the intermediate computation of the third coefficient \mathbf{Y}_3 . In detail:

$$\begin{aligned} & \mathbf{Y}_3 = D^3(\mathbf{Y}_2) - D^1(\mathcal{X}) y_3 \\ & = \left(\frac{\partial}{\partial x} + y_1 \frac{\partial}{\partial y} + y_2 \frac{\partial}{\partial y_1} + y_3 \frac{\partial}{\partial y_2} \right) \left(\mathcal{Y}_{x^2} + [2 \mathcal{Y}_{xy} - \mathcal{X}_{x^2}] y_1 + \right. \\ & \quad + [\mathcal{Y}_{y^2} - 2 \mathcal{X}_{xy}] (y_1)^2 + [-\mathcal{X}_{y^2}] (y_1)^3 + \\ & \quad \left. + [\mathcal{Y}_y - 2 \mathcal{X}_x] y_2 + [-3 \mathcal{X}_y] y_1 y_2 \right) \\ (2.7) \quad & = \underline{\mathcal{Y}_{x^3}}_{\boxed{1}} + \underline{[2 \mathcal{Y}_{x^2y} - \mathcal{X}_{x^3}] y_1}_{\boxed{2}} + \underline{[\mathcal{Y}_{xy^2} - 2 \mathcal{X}_{x^2y}] (y_1)^2}_{\boxed{3}} + \underline{[-\mathcal{X}_{xy^2}] (y_1)^3}_{\boxed{4}} + \\ & \quad + \underline{[\mathcal{Y}_{xy} - 2 \mathcal{X}_{x^2}] y_2}_{\boxed{6}} + \underline{[-3 \mathcal{X}_{xy}] y_1 y_2}_{\boxed{7}} + \underline{[\mathcal{Y}_{x^2y}] y_1}_{\boxed{2}} + \\ & \quad + \underline{[2 \mathcal{Y}_{xy^2} - \mathcal{X}_{x^2y}] (y_1)^2}_{\boxed{3}} + \underline{[\mathcal{Y}_{y^3} - 2 \mathcal{X}_{xy^2}] (y_1)^3}_{\boxed{4}} + \underline{[-\mathcal{X}_{y^3}] (y_1)^4}_{\boxed{5}} + \end{aligned}$$

$$\begin{aligned}
& + \underline{[\mathcal{Y}_{y^2} - 2\mathcal{X}_{xy}] y_1 y_2}_{\boxed{7}} + \underline{[-3\mathcal{X}_{y^2}] (y_1)^2 y_2}_{\boxed{8}} + \underline{[2\mathcal{Y}_{xy} - \mathcal{X}_{x^2}] y_2}_{\boxed{6}} + \\
& + \underline{[\mathcal{Y}_{y^2} - 2\mathcal{X}_{xy}] 2 y_1 y_2}_{\boxed{7}} + \underline{[-\mathcal{X}_{y^2}] 3(y_1)^2 y_2}_{\boxed{8}} + \underline{[-3\mathcal{X}_y] (y_2)^2}_{\boxed{9}} + \\
& + \underline{[\mathcal{Y}_y - 2\mathcal{X}_x] y_3}_{\boxed{10}} + \underline{[-3\mathcal{X}_y] y_1 y_3}_{\boxed{11}} - \\
& - \underline{[\mathcal{X}_x] y_3}_{\boxed{10}} - \underline{[\mathcal{X}_y] y_1 y_3}_{\boxed{11}}.
\end{aligned}$$

We have underlined all the terms with a number appended. Each number refers to the order of appearance of the terms in the final simplified expression of \mathbf{Y}_3 , also written in [BK1989] with different notations:

$$(2.8) \quad \left\{ \begin{aligned} \mathbf{Y}_3 &= \mathcal{Y}_{x^3} + [3\mathcal{Y}_{x^2 y} - \mathcal{X}_{x^3}] y_1 + [3\mathcal{Y}_{xy^2} - 3\mathcal{X}_{x^2 y}] (y_1)^2 + \\ & + [\mathcal{Y}_{y^3} - 3\mathcal{X}_{xy^2}] (y_1)^3 + [-\mathcal{X}_{y^3}] (y_1)^4 + [3\mathcal{Y}_{xy} - 3\mathcal{X}_{x^2}] y_2 + \\ & + [3\mathcal{Y}_{y^2} - 9\mathcal{X}_{xy}] y_1 y_2 + [-6\mathcal{X}_{y^2}] (y_1)^2 y_2 + [-3\mathcal{X}_y] (y_2)^2 + \\ & + [\mathcal{Y}_y - 3\mathcal{X}_x] y_3 + [-4\mathcal{X}_y] y_1 y_3. \end{aligned} \right.$$

After similar manual computations, the intermediate details of which we will not copy in this Latex file, we get the desired expressions of \mathbf{Y}_4 and of \mathbf{Y}_5 . Firstly:

$$(2.9) \quad \left\{ \begin{aligned} \mathbf{Y}_4 &= \mathcal{Y}_{x^4} + [4\mathcal{Y}_{x^3 y} - \mathcal{X}_{x^4}] y_1 + [6\mathcal{Y}_{x^2 y^2} - 4\mathcal{X}_{x^3 y}] (y_1)^2 + \\ & + [4\mathcal{Y}_{xy^3} - 6\mathcal{X}_{x^2 y^2}] (y_1)^3 + [\mathcal{Y}_{y^4} - 4\mathcal{X}_{xy^3}] (y_1)^4 + [-\mathcal{X}_{y^4}] (y_1)^5 + \\ & + [6\mathcal{Y}_{x^2 y} - 4\mathcal{X}_{x^3}] y_2 + [12\mathcal{Y}_{xy^2} - 18\mathcal{X}_{x^2 y}] y_1 y_2 + \\ & + [6\mathcal{Y}_{y^3} - 24\mathcal{X}_{xy^2}] (y_1)^2 y_2 + [-10\mathcal{X}_{y^3}] (y_1)^3 y_2 + \\ & + [3\mathcal{Y}_{y^2} - 12\mathcal{X}_{xy}] (y_2)^2 + [-15\mathcal{X}_{y^2}] y_1 (y_2)^2 + \\ & + [4\mathcal{Y}_{xy} - 6\mathcal{X}_{x^2}] y_3 + [4\mathcal{Y}_{y^2} - 16\mathcal{X}_{xy}] y_1 y_3 + [-10\mathcal{X}_{y^2}] (y_1)^2 y_3 + \\ & + [-10\mathcal{X}_y] y_2 y_3 + [\mathcal{Y}_y - 4\mathcal{X}_x] y_4 + [-5\mathcal{X}_y] y_1 y_4. \end{aligned} \right.$$

Secondly:

$$(2.10) \quad \left\{ \begin{aligned} \mathbf{Y}_5 &= \mathcal{Y}_{x^5} + [5\mathcal{Y}_{x^4 y} - \mathcal{X}_{x^5}] y_1 + [10\mathcal{Y}_{x^3 y^2} - 5\mathcal{X}_{x^4 y}] (y_1)^2 + \\ & + [10\mathcal{Y}_{x^2 y^3} - 10\mathcal{X}_{x^3 y^2}] (y_1)^3 + [5\mathcal{Y}_{xy^4} - 10\mathcal{X}_{x^2 y^3}] (y_1)^4 + \\ & + [\mathcal{Y}_{y^5} - 5\mathcal{X}_{xy^4}] (y_1)^5 + [-\mathcal{X}_{y^5}] (y_1)^6 + [10\mathcal{Y}_{x^3 y} - 5\mathcal{X}_{x^4}] y_2 + \\ & + [30\mathcal{Y}_{x^2 y^2} - 30\mathcal{X}_{x^3 y}] y_1 y_2 + [30\mathcal{Y}_{xy^3} - 60\mathcal{X}_{x^2 y^2}] (y_1)^2 y_2 + \\ & + [10\mathcal{Y}_{y^4} - 50\mathcal{X}_{xy^3}] (y_1)^3 y_2 + [-15\mathcal{X}_{y^4}] (y_1)^4 y_2 + \\ & + [15\mathcal{Y}_{xy^2} - 30\mathcal{X}_{x^2 y}] (y_2)^2 + [15\mathcal{Y}_{y^3} - 75\mathcal{X}_{xy^2}] y_1 (y_2)^2 + \\ & + [-45\mathcal{X}_{y^3}] (y_1)^2 (y_2)^2 + [-15\mathcal{X}_{y^2}] (y_2)^3 + \\ & + [10\mathcal{Y}_{x^2 y} - 10\mathcal{X}_{x^3}] y_3 + [20\mathcal{Y}_{xy^2} - 40\mathcal{X}_{x^2 y}] y_1 y_3 + \\ & + [10\mathcal{Y}_{y^3} - 50\mathcal{X}_{xy^2}] (y_1)^2 y_3 + [-20\mathcal{X}_{y^3}] (y_1)^3 y_3 + \\ & + [10\mathcal{Y}_{y^2} - 50\mathcal{X}_{xy}] y_2 y_3 + [-60\mathcal{X}_{y^2}] y_1 y_2 y_3 + [-10\mathcal{X}_y] (y_3)^2 + \\ & + [5\mathcal{Y}_{xy} - 10\mathcal{X}_{x^2}] y_4 + [5\mathcal{Y}_{y^2} - 25\mathcal{X}_{xy}] y_1 y_4 + [-15\mathcal{X}_{y^2}] (y_1)^2 y_4 + \\ & + [-15\mathcal{X}_y] y_2 y_4 + [\mathcal{Y}_y - 5\mathcal{X}_x] y_5 + [-6\mathcal{X}_y] y_1 y_5. \end{aligned} \right.$$

2.11. Formal inspection, formal intuition and formal induction. Now, we have to comment these formulas. We have written in length the five polynomials \mathbf{Y}_1 , \mathbf{Y}_2 , \mathbf{Y}_3 , \mathbf{Y}_4 and \mathbf{Y}_5 in the pure jet variables y_1, y_2, y_3, y_4 and y_5 . Except the first “constant” term \mathcal{Y}_{x^κ} , all the monomials in the expression of \mathbf{Y}_κ are of the general form

$$(2.12) \quad (y_{\lambda_1})^{\mu_1} (y_{\lambda_2})^{\mu_2} \dots (y_{\lambda_d})^{\mu_d},$$

for some positive integer $d \geq 1$, for some collection of strictly increasing jet indices:

$$(2.13) \quad 1 \leq \lambda_1 < \lambda_2 < \cdots < \lambda_d \leq \kappa,$$

and for some positive integers $\mu_1, \dots, \mu_d \geq 1$. This and the next combinatorial facts may be confirmed by reading the formulas giving $\mathbf{Y}_1, \mathbf{Y}_2, \mathbf{Y}_3, \mathbf{Y}_4$ and \mathbf{Y}_5 . It follows that the integer d satisfies the inequality $d \leq \kappa + 1$. To include the first “constant” term \mathcal{Y}_{x^κ} , we shall make the convention that putting $d = 0$ in the monomial (2.12) yields the constant term 1.

Furthermore, by inspecting the formulas giving $\mathbf{Y}_1, \mathbf{Y}_2, \mathbf{Y}_3, \mathbf{Y}_4$ and \mathbf{Y}_5 , we see that the following inequality should be satisfied:

$$(2.14) \quad \mu_1 \lambda_1 + \mu_2 \lambda_2 + \cdots + \mu_d \lambda_d \leq \kappa + 1.$$

For instance, in the expression of \mathbf{Y}_4 , the two monomials $(y_1)^3 y_2$ and $y_1 (y_2)^2$ do appear, but the two monomials $(y_1)^4 y_2$ and $(y_1)^2 (y_2)^2$ cannot appear. All coefficients of the pure jet monomials are of the general form:

$$(2.15) \quad [A \mathcal{Y}_{x^\alpha y^{\beta+1}} - B \mathcal{X}_{x^{\alpha+1} y^\beta}],$$

for some nonnegative integers $A, B, \alpha, \beta \in \mathbb{N}$. Sometimes A is zero, but B is zero only for the (constant, with respect to pure jet variables) term \mathcal{Y}_{x^κ} . Importantly, \mathcal{X} is differentiated once more with respect to x and \mathcal{Y} is differentiated once more with respect to y . Again, this may be confirmed by reading all the terms in the formulas for $\mathbf{Y}_1, \mathbf{Y}_2, \mathbf{Y}_3, \mathbf{Y}_4$ and \mathbf{Y}_5 .

In addition, we claim that there is a link between the couple (α, β) and the collection $\{\mu_1, \lambda_1, \dots, \mu_d, \lambda_d\}$. To discover it, let us write some of the monomials appearing in the expressions of \mathbf{Y}_4 (first column) and of \mathbf{Y}_5 (second column), for instance:

$$(2.16) \quad \left\{ \begin{array}{ll} [6 \mathcal{Y}_{x^2 y^2} - 4 \mathcal{X}_{x^3 y}] (y_1)^2, & [5 \mathcal{Y}_{x y^4} - 10 \mathcal{X}_{x^2 y^3}] (y_1)^4, \\ [12 \mathcal{Y}_{x y^2} - 18 \mathcal{X}_{x^2 y}] y_1 y_2, & [30 \mathcal{Y}_{x y^3} - 60 \mathcal{X}_{x^2 y^2}] (y_1)^2 y_2, \\ [-10 \mathcal{X}_{y^3}] (y_1)^3 y_2, & [-15 \mathcal{X}_{y^4}] (y_1)^4 y_2, \\ [4 \mathcal{Y}_{y^2} - 16 \mathcal{X}_{x y}] y_1 y_3, & [10 \mathcal{Y}_{y^2} - 50 \mathcal{X}_{x y}] y_2 y_3, \\ [-10 \mathcal{X}_{y^2}] (y_1)^2 y_3, & [-60 \mathcal{X}_{y^2}] y_1 y_2 y_3. \end{array} \right.$$

After some reflection, we discover the hidden intuitive rule: the partial derivatives of \mathcal{Y} and of \mathcal{X} associated with the monomial $(y_{\lambda_1})^{\mu_1} \cdots (y_{\lambda_d})^{\mu_d}$ are, respectively:

$$(2.17) \quad \left\{ \begin{array}{l} \mathcal{Y}_{x^{\kappa - \mu_1 \lambda_1 - \cdots - \mu_d \lambda_d} y^{\mu_1 + \cdots + \mu_d}}, \\ \mathcal{X}_{x^{\kappa - \mu_1 \lambda_1 - \cdots - \mu_d \lambda_d + 1} y^{\mu_1 + \cdots + \mu_d - 1}}. \end{array} \right.$$

This may be checked on each of the 10 examples (2.16) above.

Now that we have explored and discovered the combinatorics of the pure jet monomials, of the partial derivatives and of the complete sum giving \mathbf{Y}_κ , we may express that it is of the following general form:

$$(2.18) \quad \left\{ \begin{array}{l} \mathbf{Y}_\kappa = \mathcal{Y}_{x^\kappa} + \sum_{d=1}^{\kappa+1} \sum_{1 \leq \lambda_1 < \cdots < \lambda_d \leq \kappa} \sum_{\mu_1 \geq 1, \dots, \mu_d \geq 1} \sum_{\mu_1 \lambda_1 + \cdots + \mu_d \lambda_d \leq \kappa + 1} \\ \left[A_{\kappa}^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)} \cdot \mathcal{Y}_{x^{\kappa - \mu_1 \lambda_1 - \cdots - \mu_d \lambda_d} y^{\mu_1 + \cdots + \mu_d}} - \right. \\ \left. - B_{\kappa}^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)} \cdot \mathcal{X}_{x^{\kappa - \mu_1 \lambda_1 - \cdots - \mu_d \lambda_d + 1} y^{\mu_1 + \cdots + \mu_d - 1}} \right] \cdot \\ \cdot (y_{\lambda_1})^{\mu_1} \cdots (y_{\lambda_d})^{\mu_d}. \end{array} \right.$$

Here, we separate the first term \mathcal{Y}_{x^κ} from the general sum; it is the constant term in \mathbf{Y}_κ , which itself is a polynomial with respect to the jet variables y_λ . In this general formula, the only remaining unknowns are the nonnegative integer coefficients $A_\kappa^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)} \in \mathbb{N}$ and $B_\kappa^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)} \in \mathbb{N}$. In Section 3 below, we shall explain how we have discovered their exact value.

At present, even if we are unable to devise their explicit expression, we may observe that the value of the special integer coefficients $A_{\mu_1}^{(\mu_1, 1)}$ and $B_{\mu_1}^{(\mu_1, 1)}$ which are attached to the monomials ct. , y_1 , $(y_1)^2$, $(y_1)^3$, $(y_1)^4$ and $(y_1)^5$ are simple. Indeed, by inspecting the first terms in the expressions of \mathbf{Y}_1 , \mathbf{Y}_2 , \mathbf{Y}_3 , \mathbf{Y}_4 and \mathbf{Y}_5 , we of course recognize the binomial coefficients. In general:

Lemma 2.19. For $\kappa \geq 1$,

$$(2.20) \quad \left\{ \begin{array}{l} \mathbf{Y}_\kappa = \mathcal{Y}_{x^\kappa} + \sum_{\lambda=1}^{\kappa} \left[\binom{\kappa}{\lambda} \mathcal{Y}_{x^{\kappa-\lambda} y^\lambda} - \binom{\kappa}{\lambda-1} \mathcal{X}_{x^{\kappa-\lambda+1} y^{\lambda-1}} \right] (y_1)^\lambda + \\ \quad + [-\mathcal{X}_{y^\kappa}] (y_1)^\kappa + \text{remainder,} \end{array} \right.$$

where the term **remainder** collects all remaining monomials in the pure jet variables.

In addition, let us remind what we have observed and used in a previous co-signed work.

Lemma 2.21. ([GM2003a], p. 536) For $\kappa \geq 4$, nine among the monomials of \mathbf{Y}_κ are of the following general form:

$$(2.22) \quad \left\{ \begin{array}{l} \mathbf{Y}_\kappa = \mathcal{Y}_{x^\kappa} + [C_\kappa^1 \mathcal{Y}_{x^{\kappa-1} y} - \mathcal{X}_{x^\kappa}] y_1 + [C_\kappa^2 \mathcal{Y}_{x^{\kappa-2} y} - C_\kappa^1 \mathcal{X}_{x^{\kappa-1}}] y_2 + \\ \quad + [C_\kappa^2 \mathcal{Y}_{x^2 y} - C_\kappa^3 \mathcal{X}_{x^3}] y_{\kappa-2} + [C_\kappa^1 \mathcal{Y}_{xy} - C_\kappa^2 \mathcal{X}_{x^2}] y_{\kappa-1} + \\ \quad + [C_\kappa^1 \mathcal{Y}_{y^2} - \kappa^2 \mathcal{X}_{xy}] y_1 y_{\kappa-1} + [-C_\kappa^2 \mathcal{X}_y] y_2 y_{\kappa-1} + \\ \quad + [\mathcal{Y}_y - C_\kappa^1 \mathcal{X}_x] + [-C_{\kappa+1}^1 \mathcal{X}_y] y_1 y_\kappa + \text{remainder,} \end{array} \right.$$

where the term **remainder** denotes all the remaining monomials, and where $C_\kappa^\lambda := \frac{\kappa!}{(\kappa-\lambda)! \lambda!}$ is a notation for the binomial coefficient which occupies less space in LaTeX “equation mode” than the classical notation

$$(2.23) \quad \binom{\kappa}{\lambda}.$$

Now, we state directly the final theorem, without further inductive or intuitive information.

Theorem 2.24. For $\kappa \geq 1$, we have:

$$(2.25) \quad \boxed{\begin{array}{l} \mathbf{Y}_\kappa = \mathcal{Y}_{x^\kappa} + \sum_{d=1}^{\kappa+1} \sum_{1 \leq \lambda_1 < \dots < \lambda_d \leq \kappa} \sum_{\mu_1 \geq 1, \dots, \mu_d \geq 1} \sum_{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d \leq \kappa+1} \\ \quad \left[\frac{\kappa \cdots (\kappa - \mu_1 \lambda_1 - \dots - \mu_d \lambda_d + 1)}{(\lambda_1!)^{\mu_1} \mu_1! \cdots (\lambda_d!)^{\mu_d} \mu_d!} \cdot \mathcal{Y}_{x^{\kappa-\mu_1 \lambda_1 - \dots - \mu_d \lambda_d} y^{\mu_1 + \dots + \mu_d}} - \right. \\ \quad \left. \frac{\kappa \cdots (\kappa - \mu_1 \lambda_1 - \dots - \mu_d \lambda_d + 2)(\mu_1 \lambda_1 + \dots + \mu_d \lambda_d)}{(\lambda_1!)^{\mu_1} \mu_1! \cdots (\lambda_d!)^{\mu_d} \mu_d!} \cdot \mathcal{X}_{x^{\kappa-\mu_1 \lambda_1 - \dots - \mu_d \lambda_d + 1} y^{\mu_1 + \dots + \mu_d - 1}} \right] (y_{\lambda_1})^{\mu_1} \cdots (y_{\lambda_d})^{\mu_d}. \end{array}}$$

Once the correct theorem is formulated, its proof follows by accessible induction arguments which will not be developed here. It is better to continue through and to examine thoroughly the case of several variables, since it will help us considerably to explain how we discovered the exact values of the integer coefficients $A_{\kappa}^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)}$ and $B_{\kappa}^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)}$.

2.26. Verification and application. Before proceeding further, let us rapidly verify that the above general formula (2.25) is correct by inspecting two instances extracted from \mathbf{Y}_5 .

Firstly, the coefficient of $(y_1)^3 y_3$ in \mathbf{Y}_5 is obtained by putting $\kappa = 5$, $d = 2$, $\lambda_1 = 1$, $\mu_1 = 3$, $\lambda_2 = 3$ and $\mu_2 = 1$ in the general formula (2.25), which yields:

$$(2.27) \quad \left[0 - \frac{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 \cdot 6}{(1!)^3 3! (3!)^1 1!} \mathcal{X}_{y^3} \right] = [-20 \mathcal{X}_{y^3}].$$

This value is the same as in the original formula (2.10): confirmation.

Secondly, the coefficient of $y_1 (y_2)^2$ in \mathbf{Y}_5 is obtained by $\kappa = 5$, $d = 2$, $\lambda_1 = 1$, $\mu_1 = 1$, $\lambda_2 = 2$ and $\mu_2 = 2$ in the general formula (2.25), which yields:

$$(2.28) \quad \left[\frac{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}{(1!)^1 1! (2!)^2 2!} \mathcal{Y}_{y^3} - \frac{5 \cdot 4 \cdot 3 \cdot 2 \cdot 5}{(1!)^1 1! (2!)^2 2!} \mathcal{X}_{xy^2} \right] = [15 \mathcal{Y}_{y^3} - 75 \mathcal{X}_{xy^2}].$$

This value is the same as in the original formula (2.10); again: confirmation.

Finally, applying our general formula (2.25), we deduce the value of \mathbf{Y}_6 *without having to use \mathbf{Y}_5 and the induction formulas (2.4)*, which shortens substantially the computations.

For the pleasure, we obtain:

$$(2.29) \quad \left\{ \begin{aligned} \mathbf{Y}_6 = & \mathcal{Y}_{x^6} + [6 \mathcal{Y}_{x^5 y} - \mathcal{X}_{x^6}] y_1 + [15 \mathcal{Y}_{x^4 y^2} - 6 \mathcal{X}_{x^5 y}] (y_1)^2 + \\ & + [20 \mathcal{Y}_{x^3 y^3} - 15 \mathcal{X}_{x^4 y^2}] (y_1)^3 + [15 \mathcal{Y}_{x^2 y^4} - 20 \mathcal{X}_{x^3 y^3}] (y_1)^4 + \\ & + [6 \mathcal{Y}_{xy^5} - 15 \mathcal{X}_{x^2 y^4}] (y_1)^5 + [\mathcal{Y}_{y^6} - 6 \mathcal{X}_{xy^5}] (y_1)^6 + [-\mathcal{X}_{y^6}] (y_1)^7 + \\ & + [15 \mathcal{Y}_{x^4 y} - 6 \mathcal{X}_{x^5}] y_2 + [60 \mathcal{Y}_{x^3 y^2} - 45 \mathcal{X}_{x^4 y}] y_1 y_2 + \\ & + [90 \mathcal{Y}_{x^2 y^3} - 120 \mathcal{X}_{x^3 y^2}] (y_1)^2 y_2 + [60 \mathcal{Y}_{xy^4} - 150 \mathcal{X}_{x^2 y^3}] (y_1)^3 y_2 + \\ & + [15 \mathcal{Y}_{y^5} - 90 \mathcal{X}_{xy^4}] (y_1)^4 y_2 + [-21 \mathcal{X}_{y^5}] (y_1)^5 y_2 + \\ & + [45 \mathcal{Y}_{x^2 y^2} - 60 \mathcal{X}_{x^3 y}] (y_2)^2 + [90 \mathcal{Y}_{xy^3} - 225 \mathcal{X}_{x^2 y^2}] y_1 (y_2)^2 + \\ & + [45 \mathcal{Y}_{y^4} - 270 \mathcal{X}_{xy^3}] (y_1)^2 (y_2)^2 + [-210 \mathcal{X}_{y^4}] (y_1)^3 (y_2)^2 + \\ & + [15 \mathcal{Y}_{y^3} - 90 \mathcal{X}_{xy^2}] (y_2)^3 + [-105 \mathcal{X}_{y^3}] y_1 (y_2)^3 + \\ & + [20 \mathcal{Y}_{x^3 y} - 15 \mathcal{X}_{x^4}] y_3 + [60 \mathcal{Y}_{x^2 y^2} - 80 \mathcal{X}_{x^3 y}] y_1 y_3 + \\ & + [60 \mathcal{Y}_{xy^3} - 150 \mathcal{X}_{x^2 y^2}] (y_1)^2 y_3 + [20 \mathcal{Y}_{y^4} - 120 \mathcal{X}_{xy^3}] (y_1)^3 y_3 + \\ & + [-35 \mathcal{X}_{y^4}] (y_1)^4 y_3 + [60 \mathcal{Y}_{xy^2} - 150 \mathcal{X}_{x^2 y}] y_2 y_3 + \\ & + [60 \mathcal{Y}_{y^3} - 360 \mathcal{X}_{xy^2}] y_1 y_2 y_3 + [-210 \mathcal{X}_{y^3}] (y_1)^2 y_2 y_3 + \\ & + [-105 \mathcal{X}_{y^2}] (y_2)^2 y_3 + [10 \mathcal{Y}_{y^2} - 60 \mathcal{X}_{xy}] (y_3)^2 + \\ & + [-70 \mathcal{X}_{y^2}] y_1 (y_3)^2 + [15 \mathcal{Y}_{x^2 y} - 20 \mathcal{X}_{x^3}] y_4 + \\ & + [30 \mathcal{Y}_{xy^2} - 75 \mathcal{X}_{x^2 y}] y_1 y_4 + [15 \mathcal{Y}_{y^3} - 90 \mathcal{X}_{xy^2}] (y_1)^2 y_4 + \\ & + [-35 \mathcal{X}_{y^3}] (y_1)^3 y_4 + [15 \mathcal{Y}_{y^2} - 90 \mathcal{X}_{xy}] y_2 y_4 + \\ & + [-105 \mathcal{X}_{y^2}] y_1 y_2 y_4 + [-35 \mathcal{X}_{y^2}] y_3 y_4 + [6 \mathcal{Y}_{xy} - 15 \mathcal{X}_{x^2}] y_5 + \\ & + [6 \mathcal{Y}_{y^2} - 36 \mathcal{X}_{xy}] y_1 y_5 + [-21 \mathcal{X}_{y^2}] (y_1)^2 y_5 + [-21 \mathcal{X}_{y^2}] y_2 y_5 + \\ & + [\mathcal{Y}_y - 6 \mathcal{X}_y] y_6 + [-7 \mathcal{X}_y] y_1 y_6. \end{aligned} \right.$$

2.30. Deduction of the classical Faà di Bruno formula. Let $x, y \in \mathbb{K}$ and let $g = g(x)$, $f = f(y)$ be two C^∞ -smooth functions $\mathbb{K} \rightarrow \mathbb{K}$. Consider the composition $h := f \circ g$, namely $h(x) = f(g(x))$. For $\lambda \in \mathbb{N}$ with $\lambda \geq 1$, simply denote by g_λ the λ -th derivative $\frac{d^\lambda g}{dx^\lambda}$ and similarly for h_λ . Also, abbreviate $f_\lambda := \frac{d^\lambda f}{dy^\lambda}$.

By the classical formula for the derivative of a composite function, we have $h_1 = f_1 g_1$. Further computations provide the following list of subsequent derivatives of h :

$$(2.31) \quad \left\{ \begin{array}{l} h_1 = f_1 g_1, \\ h_2 = f_2 (g_1)^2 + f_1 g_2, \\ h_3 = f_3 (g_1)^3 + 3 f_2 g_1 g_2 + f_1 g_3, \\ h_4 = f_4 (g_1)^4 + 6 f_3 (g_1)^2 g_2 + 3 f_2 (g_2)^2 + 4 f_2 g_1 g_3 + f_1 g_4, \\ h_5 = f_5 (g_1)^5 + 10 f_4 (g_1)^3 g_2 + 15 f_3 (g_1)^2 g_3 + 10 f_3 g_1 (g_2)^2 + \\ \quad + 10 f_2 g_2 g_3 + 5 f_2 g_1 g_4 + f_1 g_5, \\ h_6 = f_6 (g_1)^6 + 15 f_5 (g_1)^4 g_2 + 45 f_4 (g_1)^2 (g_2)^2 + 15 f_3 (g_2)^3 + \\ \quad + 20 f_4 (g_1)^3 g_3 + 60 f_3 g_1 g_2 g_3 + 10 f_2 (g_3)^2 + 15 f_3 (g_1)^2 g_4 + \\ \quad + 15 f_2 g_2 g_4 + 6 f_2 g_1 g_5 + f_1 g_6. \end{array} \right.$$

Theorem 2.32. For every integer $\kappa \geq 1$, the κ -th derivative of the composite function $h = f \circ g$ may be expressed as an explicit polynomial in the partial derivatives of f and of g having integer coefficients:

$$(2.33) \quad \boxed{\frac{d^\kappa h}{dx^\kappa} = \sum_{d=1}^{\kappa} \sum_{1 \leq \lambda_1 < \dots < \lambda_d \leq \kappa} \sum_{\mu_1 \geq 1, \dots, \mu_d \geq 1} \sum_{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d = \kappa} \frac{\kappa!}{(\lambda_1!)^{\mu_1} \mu_1! \dots (\lambda_d!)^{\mu_d} \mu_d!} \frac{d^{\mu_1 + \dots + \mu_d} f}{dy^{\mu_1 + \dots + \mu_d}} \left(\frac{d^{\lambda_1} g}{dx^{\lambda_1}} \right)^{\mu_1} \dots \left(\frac{d^{\lambda_d} g}{dx^{\lambda_d}} \right)^{\mu_d} .}$$

This is the classical *Faà di Bruno formula*. Interestingly, we observe that this formula is included as a subpart of the general formula for Y_κ , after a suitable translation. Indeed, in the formulas for $Y_1, Y_2, Y_3, Y_4, Y_5, Y_6$ and in the general sum for Y_κ , pick only the terms for which $\mu_1 \lambda_1 + \dots + \mu_d \lambda_d = \kappa$ and drop \mathcal{X} , which yields:

$$(2.34) \quad \sum_{d=1}^{\kappa} \sum_{1 \leq \lambda_1 < \dots < \lambda_d \leq \kappa} \sum_{\mu_1 \geq 1, \dots, \mu_d \geq 1} \sum_{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d = \kappa} \left[\frac{\kappa!}{\mu_1! (\lambda_1!)^{\mu_1} \dots \mu_d! (\lambda_d!)^{\mu_d}} \mathcal{Y}_{y^{\mu_1 + \dots + \mu_d}} \right] (y_{\lambda_1})^{\mu_1} \dots (y_{\lambda_d})^{\mu_d} .$$

The similarity between the two formulas (2.33) and (2.34) is now clearly visible.

The Faà di Bruno formula may be established by means of substitutions of power series ([F1969], p. 222), by means of the umbral calculus ([CS1996]), or by means of some

induction formulas, which we write for completeness. Define the differential operators

$$\begin{aligned}
 (2.35) \quad & F_2 := g_2 \frac{\partial}{\partial g_1} + g_1 \left(f_2 \frac{\partial}{\partial f_1} \right), \\
 & F_3 := g_2 \frac{\partial}{\partial g_1} + g_3 \frac{\partial}{\partial g_2} + g_1 \left(f_2 \frac{\partial}{\partial f_1} + f_3 \frac{\partial}{\partial f_2} \right), \\
 & \dots\dots\dots \\
 & F_\lambda := g_2 \frac{\partial}{\partial g_1} + g_3 \frac{\partial}{\partial g_2} + \dots + g_\lambda \frac{\partial}{\partial g_{\lambda-1}} + g_1 \left(f_2 \frac{\partial}{\partial f_1} + f_3 \frac{\partial}{\partial f_2} + \dots + f_\lambda \frac{\partial}{\partial f_{\lambda-1}} \right).
 \end{aligned}$$

Then we have

$$\begin{aligned}
 (2.36) \quad & h_2 = F^2(h_1), \\
 & h_3 = F^3(h_2), \\
 & \dots\dots\dots \\
 & h_\lambda = F^\lambda(h_{\lambda-1}).
 \end{aligned}$$

§3. SEVERAL INDEPENDENT VARIABLES AND ONE DEPENDENT VARIABLE

3.1. Simplified adapted notations. As announced after the statement of Theorem 2.24, it is only after we have treated the case of several independent variables that we will understand perfectly the general formula (2.25), valid in the case of one independent variable and one dependent variable. We will discover massive formal computations, exciting our computational intuition.

Thus, assume $n \geq 1$ and $m = 1$, let $\kappa \in \mathbb{N}$ with $\kappa \geq 1$ and simply denote (instead of (1.2)) the jet variables by:

$$(3.2) \quad (x^i, y, y_{i_1}, y_{i_1, i_2}, \dots, y_{i_1, i_2, \dots, i_\kappa}).$$

Also, instead of (1.30), denote the κ -th prolongation of a vector field by:

$$(3.3) \quad \left\{ \begin{aligned}
 \mathcal{L}^{(\kappa)} = & \sum_{i=1}^n \mathcal{X}^i \frac{\partial}{\partial x^i} + \mathcal{Y} \frac{\partial}{\partial y} + \sum_{i_1=1}^n \mathbf{Y}_{i_1} \frac{\partial}{\partial y_{i_1}} + \sum_{i_1, i_2=1}^n \mathbf{Y}_{i_1, i_2} \frac{\partial}{\partial y_{i_1, i_2}} + \\
 & + \dots + \sum_{i_1, i_2, \dots, i_\kappa=1}^n \mathbf{Y}_{i_1, i_2, \dots, i_\kappa} \frac{\partial}{\partial y_{i_1, i_2, \dots, i_\kappa}}.
 \end{aligned} \right.$$

The induction formulas are

$$(3.4) \quad \left\{ \begin{aligned}
 \mathbf{Y}_{i_1} & := D_{i_1}^1(\mathcal{Y}) - \sum_{k=1}^n D_{i_1}^1(\mathcal{X}^k) y_k, \\
 \mathbf{Y}_{i_1, i_2} & := D_{i_2}^2(\mathbf{Y}_{i_1}) - \sum_{k=1}^n D_{i_2}^1(\mathcal{X}^k) y_{i_1, k}, \\
 & \dots\dots\dots \\
 \mathbf{Y}_{i_1, i_2, \dots, i_\kappa} & := D_{i_\kappa}^\kappa(\mathbf{Y}_{i_1, i_2, \dots, i_{\kappa-1}}) - \sum_{k=1}^n D_{i_\kappa}^1(\mathcal{X}^k) y_{i_1, i_2, \dots, i_{\kappa-1}, k},
 \end{aligned} \right.$$

where the total differentiation operators $D_{j'}^\lambda$ are defined as in (1.22), dropping the sums $\sum_{j'=1}^m$ and the indices j' .

3.5. Two instructing explicit computations. To begin with, let us compute \mathbf{Y}_{i_1} . With $D_{i_1}^1 = \frac{\partial}{\partial x^{i_1}} + y_{i_1} \frac{\partial}{\partial y}$, we have:

$$(3.6) \quad \begin{aligned} \mathbf{Y}_{i_1} &= D_{i_1}(\mathcal{Y}) - \sum_{k_1=1}^n D_{i_1}^1(\mathcal{X}^{k_1}) y_{k_1} \\ &= \mathcal{Y}_{x^{i_1}} + \mathcal{Y}_y y_{i_1} - \sum_{k_1=1}^n \mathcal{X}_{x^{i_1}}^{k_1} y_{k_1} - \sum_{k_1=1}^n \mathcal{X}_y^{k_1} y_{i_1} y_{k_1}. \end{aligned}$$

Searching for formal harmony and for coherence with the formula (2.6)₁, we must include the term $\mathcal{Y}_y y_{i_1}$ inside the sum $\sum_{k_1=1}^n [\cdot] y_{k_1}$. Using the Kronecker symbol, we may write:

$$(3.7) \quad \mathcal{Y}_y y_{i_1} \equiv \sum_{k_1=1}^n [\delta_{i_1}^{k_1} \mathcal{Y}_y] y_{k_1}.$$

Also, we may rewrite the last term of (3.6) with a double sum:

$$(3.8) \quad - \sum_{k_1=1}^n \mathcal{X}_y^{k_1} y_{i_1} y_{k_1} \equiv \sum_{k_1, k_2=1}^n [-\delta_{i_1}^{k_1} \mathcal{X}_y^{k_2}] y_{k_1} y_{k_2}.$$

From now on and up to equation (3.39), we shall abbreviate any sum $\sum_{k=1}^n$ from 1 to n as \sum_k . Putting everything together, we get the final desired perfect expression of \mathbf{Y}_{i_1} :

$$(3.9) \quad \mathbf{Y}_{i_1} = \mathcal{Y}_{x^{i_1}} + \sum_{k_1} [\delta_{i_1}^{k_1} \mathcal{Y}_y - \mathcal{X}_{x^{i_1}}^{k_1}] y_{k_1} + \sum_{k_1, k_2} [-\delta_{i_1}^{k_1} \mathcal{X}_y^{k_2}] y_{k_1} y_{k_2}.$$

This completes the first explicit computation.

The second one is about \mathbf{Y}_{i_1, i_2} . It becomes more delicate, because several algebraic transformations must be achieved until the final satisfying formula is obtained. Our goal is to present each step very carefully, explaining every tiny detail. Without such a care, it would be impossible to claim that some of our subsequent computations, for which we will not provide the intermediate steps, may be redone and verified. Consequently, we will expose our rules of formal computation thoroughly.

Replacing the value of \mathbf{Y}_1 just obtained in the induction formula (3.4)₂ and developing, we may conduct the very first steps of the computation:

$$\begin{aligned} \mathbf{Y}_{i_1, i_2} &= D_{i_2}^2(\mathbf{Y}_{i_1}) - \sum_{k_1} D_{i_2}^1(\mathcal{X}^{k_1}) y_{i_1, k_1} \\ &= \left(\frac{\partial}{\partial x^{i_2}} + y_{i_2} \frac{\partial}{\partial y} + \sum_{k_1} y_{i_2, k_1} \frac{\partial}{\partial y_{k_1}} \right) \left(\mathcal{Y}_{x^{i_1}} + \sum_{k_1} [\delta_{i_1}^{k_1} \mathcal{Y}_y - \mathcal{X}_{x^{i_1}}^{k_1}] y_{k_1} + \right. \\ &\quad \left. + \sum_{k_1, k_2} [-\delta_{i_1}^{k_1} \mathcal{X}_y^{k_2}] y_{k_1} y_{k_2} \right) - \sum_{k_1} [\mathcal{X}_{x^{i_2}}^{k_1} + y_{i_2} \mathcal{X}_y^{k_1}] y_{i_1, k_1} \end{aligned}$$

(3.10)

$$\begin{aligned}
&= \left(\frac{\partial}{\partial x^{i_2}} \right) \left(\mathcal{Y}_{x^{i_1}} + \sum_{k_1} [\delta_{i_1}^{k_1} \mathcal{Y}_y - \mathcal{X}_{x^{i_1}}^{k_1}] y_{k_1} + \sum_{k_1, k_2} [-\delta_{i_1}^{k_1} \mathcal{X}_y^{k_2}] y_{k_1} y_{k_2} \right) + \\
&+ \left(y_{i_2} \frac{\partial}{\partial y} \right) \left(\mathcal{Y}_{x^{i_1}} + \sum_{k_1} [\delta_{i_1}^{k_1} \mathcal{Y}_y - \mathcal{X}_{x^{i_1}}^{k_1}] y_{k_1} + \sum_{k_1, k_2} [-\delta_{i_1}^{k_1} \mathcal{X}_y^{k_2}] y_{k_1} y_{k_2} \right) + \\
&+ \left(\sum_{k_1} y_{i_2, k_1} \frac{\partial}{\partial y_{k_1}} \right) \left(\mathcal{Y}_{x^{i_1}} + \sum_{k_1} [\delta_{i_1}^{k_1} \mathcal{Y}_y - \mathcal{X}_{x^{i_1}}^{k_1}] y_{k_1} + \sum_{k_1, k_2} [-\delta_{i_1}^{k_1} \mathcal{X}_y^{k_2}] y_{k_1} y_{k_2} \right) + \\
&+ \sum_{k_1} [-\mathcal{X}_{x^{i_2}}^{k_1}] y_{k_1, i_1} + \sum_{k_1} [-\mathcal{X}_y^{k_1}] y_{i_2} y_{i_1, k_1} \\
&= \mathcal{Y}_{x^{i_1} x^{i_2}} + \sum_{k_1} [\delta_{i_1}^{k_1} \mathcal{Y}_{x^{i_2} y} - \mathcal{X}_{x^{i_1} x^{i_2}}^{k_1}] y_{k_1} + \sum_{k_1, k_2} [-\delta_{i_1}^{k_1} \mathcal{X}_{x^{i_2} y}^{k_2}] y_{k_1} y_{k_2} + \\
&+ \mathcal{Y}_{x^{i_1} y} y_{i_2} + \sum_{k_1} [\delta_{i_1}^{k_1} \mathcal{Y}_{yy} - \mathcal{X}_{x^{i_1} y}^{k_1}] y_{k_1} y_{i_2} + \sum_{k_1, k_2} [-\delta_{i_1}^{k_1} \mathcal{X}_{yy}^{k_2}] y_{k_1} y_{k_2} y_{i_2} + \\
&+ \sum_{k_1} [\delta_{i_1}^{k_1} \mathcal{Y}_y - \mathcal{X}_{x^{i_1}}^{k_1}] y_{i_2, k_1} + \sum_{k_1, k_2} [-\delta_{i_1}^{k_1} \mathcal{X}_y^{k_2}] y_{k_2} y_{i_2, k_1} + \sum_{k_1, k_2} [-\delta_{i_1}^{k_1} \mathcal{X}_y^{k_2}] y_{k_1} y_{i_2, k_2} + \\
&+ \sum_{k_1} [-\mathcal{X}_{x^{i_2}}^{k_1}] y_{k_1, i_1} + \sum_{k_1} [-\mathcal{X}_y^{k_1}] y_{i_2} y_{i_1, k_1}.
\end{aligned}$$

Some explanations are needed about the computation of the last two terms of line 11, *i.e.* about the passage from line 7 of (3.10) just above to line 11. We have to compute:

$$(3.11) \quad \left(\sum_{k_1} y_{i_2, k_1} \frac{\partial}{\partial y_{k_1}} \right) \left(\sum_{k_1, k_2} [-\delta_{i_1}^{k_1} \mathcal{X}_y^{k_2}] y_{k_1} y_{k_2} \right).$$

This term is of the form

$$(3.12) \quad \left(\sum_{k_1} A_{k_1} \frac{\partial}{\partial y_{k_1}} \right) \left(\sum_{k_1, k_2} [B_{k_1, k_2}] y_{k_1} y_{k_2} \right),$$

where the terms B_{k_1, k_2} are independent of the pure first jet variables y_{x^k} . By the rule of Leibniz for the differentiation of a product, we may write

$$\begin{aligned}
&\left(\sum_{k_1} A_{k_1} \frac{\partial}{\partial y_{k_1}} \right) \left(\sum_{k_1, k_2} [B_{k_1, k_2}] y_{k_1} y_{k_2} \right) = \\
(3.13) \quad &= \sum_{k_1, k_2} [B_{k_1, k_2}] y_{k_2} \left(\sum_{k'_1} A_{k'_1} \frac{\partial}{\partial y_{k'_1}} (y_{k_1}) \right) + \sum_{k_1, k_2} [B_{k_1, k_2}] y_{k_1} \left(\sum_{k'_2} A_{k'_2} \frac{\partial}{\partial y_{k'_2}} (y_{k_2}) \right) \\
&= \sum_{k_1, k_2} [B_{k_1, k_2}] y_{k_2} A_{k_1} + \sum_{k_1, k_2} [B_{k_1, k_2}] y_{k_1} A_{k_2}.
\end{aligned}$$

This is how we have written line 11 of (3.10).

Next, the first term $\mathcal{Y}_{x^{i_1} y} y_{i_2}$ in line 10 of (3.10) is not in a suitable shape. For reasons of harmony and coherence, we must insert it inside a sum of the form $\sum_{k_1} [\cdot] y_{k_1}$. Hence, using the Kronecker symbol, we transform:

$$(3.14) \quad \mathcal{Y}_{x^{i_1} y} y_{i_2} \equiv \sum_{k_1} [\delta_{i_2}^{k_1} \mathcal{Y}_{x^{i_1} y}] y_{k_1}.$$

Also, we must “summify” the seven other terms, remaining in lines 10, 11 and 12 of (3.10). Sometimes, we use the symmetry $y_{i_2, k_1} \equiv y_{k_1, i_2}$ without mention. Similarly, we get:

$$\begin{aligned}
& \sum_{k_1} [\delta_{i_1}^{k_1} \mathcal{Y}_{yy} - \mathcal{X}_{x^{i_1}y}^{k_1}] y_{k_1} y_{i_2} \equiv \sum_{k_1, k_2} [\delta_{i_1}^{k_1} \delta_{i_2}^{k_2} \mathcal{Y}_{yy} - \delta_{i_2}^{k_2} \mathcal{X}_{x^{i_1}y}^{k_1}] y_{k_1} y_{k_2}, \\
& \sum_{k_1, k_2} [-\delta_{i_1}^{k_1} \mathcal{X}_{yy}^{k_2}] y_{k_1} y_{k_2} y_{i_2} \equiv \sum_{k_1, k_2, k_3} [-\delta_{i_1}^{k_1} \delta_{i_2}^{k_3} \mathcal{X}_{yy}^{k_2}] y_{k_1} y_{k_2} y_{k_3}, \\
& \sum_{k_1} [\delta_{i_1}^{k_1} \mathcal{Y}_y - \mathcal{X}_{x^{i_1}}^{k_1}] y_{k_1, i_2} \equiv \sum_{k_1, k_2} [\delta_{i_1}^{k_1} \delta_{i_2}^{k_2} \mathcal{Y}_y - \delta_{i_2}^{k_2} \mathcal{X}_{x^{i_1}}^{k_1}] y_{k_1, k_2}, \\
& \sum_{k_1, k_2} [-\delta_{i_1}^{k_1} \mathcal{X}_y^{k_2}] y_{k_2} y_{k_1, i_2} = \sum_{k_1, k_2} [-\delta_{i_1}^{k_2} \mathcal{X}_y^{k_1}] y_{k_1} y_{k_2, i_2} \\
(3.15) \quad & \equiv \sum_{k_1, k_2, k_3} [-\delta_{i_1}^{k_2} \delta_{i_2}^{k_3} \mathcal{X}_y^{k_1}] y_{k_1} y_{k_2, k_3}, \\
& \sum_{k_1, k_2} [-\delta_{i_1}^{k_1} \mathcal{X}_y^{k_2}] y_{k_1} y_{k_2, i_2} \equiv \sum_{k_1, k_2, k_3} [-\delta_{i_1}^{k_1} \delta_{i_2}^{k_3} \mathcal{X}_y^{k_2}] y_{k_1} y_{k_2, k_3}, \\
& \sum_{k_1} [-\mathcal{X}_{x^{i_2}}^{k_1}] y_{k_1, i_1} \equiv \sum_{k_1, k_2} [-\delta_{i_1}^{k_2} \mathcal{X}_{x^{i_2}}^{k_1}] y_{k_1, k_2}, \\
& \sum_{k_1} [-\mathcal{X}_y^{k_1}] y_{i_2} y_{k_1, i_1} = \sum_{k_2} [-\mathcal{X}_y^{k_2}] y_{i_2} y_{k_2, i_1} \\
& \equiv \sum_{k_1, k_2, k_3} [-\delta_{i_2}^{k_1} \delta_{i_1}^{k_3} \mathcal{X}_y^{k_2}] y_{k_1} y_{k_2, k_3}.
\end{aligned}$$

In the sequel, for products of Kronecker symbols, it will be convenient to adopt the following self-evident contracted notation:

$$(3.16) \quad \delta_{i_1}^{k_1} \delta_{i_2}^{k_2} \equiv \delta_{i_1, i_2}^{k_1, k_2}; \quad \text{generally : } \delta_{i_1}^{k_1} \delta_{i_2}^{k_2} \cdots \delta_{i_\lambda}^{k_\lambda} \equiv \delta_{i_1, i_2, \dots, i_\lambda}^{k_1, k_2, \dots, k_\lambda}.$$

Re-inserting plainly these eight summified terms (3.14), (3.15) in the last expression (3.10) of \mathbf{Y}_{i_1, i_2} (lines 10, 11 and 12), we get:

$$\begin{aligned}
(3.17) \quad \mathbf{Y}_{i_1, i_2} = & \underbrace{\mathcal{Y}_{x^{i_1}x^{i_2}}}_{\boxed{1}} + \underbrace{\sum_{k_1} [\delta_{i_1}^{k_1} \mathcal{Y}_{x^{i_2}y} - \mathcal{X}_{x^{i_1}x^{i_2}}^{k_1}] y_{k_1}}_{\boxed{2}} + \underbrace{\sum_{k_1, k_2} [-\delta_{i_1}^{k_1} \mathcal{X}_{x^{i_2}y}^{k_2}] y_{k_1} y_{k_2}}_{\boxed{3}} + \\
& + \underbrace{\sum_{k_1} [\delta_{i_2}^{k_1} \mathcal{Y}_{x^{i_1}y}] y_{k_1}}_{\boxed{2}} + \underbrace{\sum_{k_1, k_2} [\delta_{i_1, i_2}^{k_1, k_2} \mathcal{Y}_{yy} - \delta_{i_2}^{k_2} \mathcal{X}_{x^{i_1}y}^{k_1}] y_{k_1} y_{k_2}}_{\boxed{3}} + \\
& + \underbrace{\sum_{k_1, k_2, k_3} [-\delta_{i_1, i_2}^{k_1, k_3} \mathcal{X}_{yy}^{k_2}] y_{k_1} y_{k_2} y_{k_3}}_{\boxed{4}} + \underbrace{\sum_{k_1, k_2} [\delta_{i_1, i_2}^{k_1, k_2} \mathcal{Y}_y - \delta_{i_2}^{k_2} \mathcal{X}_{x^{i_1}}^{k_1}] y_{k_1, k_2}}_{\boxed{5}} + \\
& + \underbrace{\sum_{k_1, k_2, k_3} [-\delta_{i_1, i_2}^{k_2, k_3} \mathcal{X}_y^{k_1}] y_{k_1} y_{k_2, k_3}}_{\boxed{6}} + \underbrace{\sum_{k_1, k_2, k_3} [-\delta_{i_1, i_2}^{k_1, k_3} \mathcal{X}_y^{k_2}] y_{k_1} y_{k_2, k_3}}_{\boxed{6}} + \\
& + \underbrace{\sum_{k_1, k_2} [-\delta_{i_1}^{k_2} \mathcal{X}_{x^{i_2}}^{k_1}] y_{k_1, k_2}}_{\boxed{5}} + \underbrace{\sum_{k_1, k_2, k_3} [-\delta_{i_2, i_1}^{k_1, k_3} \mathcal{X}_y^{k_2}] y_{k_1} y_{k_2, k_3}}_{\boxed{6}}.
\end{aligned}$$

Next, we gather the underlined terms, ordering them according to their number. This yields 6 collections of sums of monomials in the pure jet variables:

$$\begin{aligned}
(3.18) \quad \mathbf{Y}_{i_1, i_2} &= \mathcal{Y}_{x^{i_1} x^{i_2}} + \sum_{k_1} \left[\delta_{i_1}^{k_1} \mathcal{Y}_{x^{i_2} y} + \delta_{i_2}^{k_1} \mathcal{Y}_{x^{i_1} y} - \mathcal{X}_{x^{i_1} x^{i_2}}^{k_1} \right] y_{k_1} + \\
&+ \sum_{k_1, k_2} \left[\delta_{i_1, i_2}^{k_1, k_2} \mathcal{Y}_{yy} - \delta_{i_1}^{k_1} \mathcal{X}_{x^{i_2} y}^{k_2} - \delta_{i_2}^{k_2} \mathcal{X}_{x^{i_1} y}^{k_1} \right] y_{k_1} y_{k_2} + \\
&+ \sum_{k_1, k_2, k_3} \left[-\delta_{i_1, i_2}^{k_1, k_3} \mathcal{X}_{yy}^{k_2} \right] y_{k_1} y_{k_2} y_{k_3} + \\
&+ \sum_{k_1, k_2} \left[\delta_{i_1, i_2}^{k_1, k_2} \mathcal{Y}_y - \delta_{i_2}^{k_2} \mathcal{X}_{x^{i_1}}^{k_1} - \delta_{i_1}^{k_2} \mathcal{X}_{x^{i_2}}^{k_1} \right] y_{k_1, k_2} + \\
&+ \sum_{k_1, k_2, k_3} \left[-\delta_{i_1, i_2}^{k_2, k_3} \mathcal{X}_y^{k_1} - \delta_{i_1, i_2}^{k_1, k_3} \mathcal{X}_y^{k_2} - \delta_{i_2, i_1}^{k_1, k_3} \mathcal{X}_y^{k_2} \right] y_{k_1} y_{k_2, k_3}.
\end{aligned}$$

To attain the real perfect harmony, this last expression has still to be worked out a little bit.

Lemma 3.19. *The final expression of \mathbf{Y}_{i_1, i_2} is as follows:*

$$(3.20) \quad \left\{ \begin{aligned}
\mathbf{Y}_{i_1, i_2} &= \mathcal{Y}_{x^{i_1} x^{i_2}} + \sum_{k_1} \left[\delta_{i_1}^{k_1} \mathcal{Y}_{x^{i_2} y} + \delta_{i_2}^{k_1} \mathcal{Y}_{x^{i_1} y} - \mathcal{X}_{x^{i_1} x^{i_2}}^{k_1} \right] y_{k_1} + \\
&+ \sum_{k_1, k_2} \left[\delta_{i_1, i_2}^{k_1, k_2} \mathcal{Y}_{yy} - \delta_{i_1}^{k_1} \mathcal{X}_{x^{i_2} y}^{k_2} - \delta_{i_2}^{k_1} \mathcal{X}_{x^{i_1} y}^{k_2} \right] y_{k_1} y_{k_2} + \\
&+ \sum_{k_1, k_2, k_3} \left[-\delta_{i_1, i_2}^{k_1, k_2} \mathcal{X}_{yy}^{k_3} \right] y_{k_1} y_{k_2} y_{k_3} + \\
&+ \sum_{k_1, k_2} \left[\delta_{i_1, i_2}^{k_1, k_2} \mathcal{Y}_y - \delta_{i_1}^{k_1} \mathcal{X}_{x^{i_2}}^{k_2} - \delta_{i_2}^{k_1} \mathcal{X}_{x^{i_1}}^{k_2} \right] y_{k_1, k_2} + \\
&+ \sum_{k_1, k_2, k_3} \left[-\delta_{i_1, i_2}^{k_1, k_2} \mathcal{X}_y^{k_3} - \delta_{i_1, i_2}^{k_3, k_1} \mathcal{X}_y^{k_2} - \delta_{i_1, i_2}^{k_2, k_3} \mathcal{X}_y^{k_1} \right] y_{k_1} y_{k_2, k_3}.
\end{aligned} \right.$$

Proof. As promised, we explain every tiny detail.

The first lines of (3.18) and of (3.20) are exactly the same. For the transformations of terms in the second, in the third and in the fourth lines, we use the following device. Let Υ_{k_1, k_2} be an indexed quantity which is symmetric: $\Upsilon_{k_1, k_2} = \Upsilon_{k_2, k_1}$. Let A_{k_1, k_2} be an arbitrary indexed quantity. Then obviously:

$$(3.21) \quad \sum_{k_1, k_2} A_{k_1, k_2} \Upsilon_{k_1, k_2} = \sum_{k_1, k_2} A_{k_2, k_1} \Upsilon_{k_1, k_2}.$$

Similar relations hold with a quantity $\Upsilon_{i_1, i_2, \dots, i_\lambda}$ which is symmetric with respect to its λ indices. Consequently, in the second, in the third and in the fourth lines of (3.18), we may permute freely certain indices in some of the terms inside the brackets. This yields the passage from lines 2, 3 and 4 of (3.18) to lines 2, 3 and 4 of (3.20).

It remains to explain how we pass from the fifth (last) line of (3.18) to the fifth (last) line of (3.20). The bracket in the fifth line of (3.18) contains three terms: $[-T_1 - T_2 - T_3]$. The term T_3 involves the product $\delta_{i_2, i_1}^{k_1, k_3}$, which we rewrite as $\delta_{i_1, i_2}^{k_3, k_1}$, in order that i_1 appears before i_2 . Then, we rewrite the three terms in the new order $[-T_2 - T_3 - T_1]$, which yields:

$$(3.22) \quad \sum_{k_1, k_2, k_3} \left[-\delta_{i_1, i_2}^{k_1, k_3} \mathcal{X}_y^{k_2} - \delta_{i_1, i_2}^{k_3, k_1} \mathcal{X}_y^{k_2} - \delta_{i_1, i_2}^{k_2, k_3} \mathcal{X}_y^{k_1} \right] y_{k_1} y_{k_2, k_3}.$$

It remains to observe that we can permute k_2 and k_3 in the first term $-T_2$, which yields the last line of (3.20). The detailed proof is complete. \square

3.23. Final perfect expression of $\mathbf{Y}_{i_1, i_2, i_3}$. Thanks to similar (longer) computations, we have obtained an expression of $\mathbf{Y}_{i_1, i_2, i_3}$ which we consider to be in final harmonious shape. Without copying the intermediate steps, let us write down the result. The comments which are necessary to read it and to interpret it start just below.

$$\begin{aligned}
(3.24) \quad \mathbf{Y}_{i_1, i_2, i_3} = & \mathcal{Y}_{x^{i_1} x^{i_2} x^{i_3}} + \sum_{k_1} \left[\delta_{i_1}^{k_1} \mathcal{Y}_{x^{i_2} x^{i_3} y} + \delta_{i_2}^{k_1} \mathcal{Y}_{x^{i_1} x^{i_3} y} + \delta_{i_3}^{k_1} \mathcal{Y}_{x^{i_1} x^{i_2} y} - \mathcal{X}_{x^{i_1} x^{i_2} x^{i_3}}^{k_1} \right] y_{k_1} + \\
& + \sum_{k_1, k_2} \left[\delta_{i_1, i_2}^{k_1, k_2} \mathcal{Y}_{x^{i_3} y^2} + \delta_{i_3, i_1}^{k_1, k_2} \mathcal{Y}_{x^{i_2} y^2} + \delta_{i_2, i_3}^{k_1, k_2} \mathcal{Y}_{x^{i_1} y^2} - \right. \\
& \quad \left. - \delta_{i_1}^{k_1} \mathcal{X}_{x^{i_2} x^{i_3} y}^{k_2} - \delta_{i_2}^{k_1} \mathcal{X}_{x^{i_1} x^{i_3} y}^{k_2} - \delta_{i_3}^{k_1} \mathcal{X}_{x^{i_1} x^{i_2} y}^{k_2} \right] y_{k_1} y_{k_2} + \\
& + \sum_{k_1, k_2, k_3} \left[\delta_{i_1, i_2, i_3}^{k_1, k_2, k_3} \mathcal{Y}_{y^3} - \delta_{i_1, i_2}^{k_1, k_2} \mathcal{X}_{x^{i_3} y^2}^{k_3} - \delta_{i_1, i_3}^{k_1, k_2} \mathcal{X}_{x^{i_2} y^2}^{k_3} - \delta_{i_2, i_3}^{k_1, k_2} \mathcal{X}_{x^{i_1} y^2}^{k_3} \right] y_{k_1} y_{k_2} y_{k_3} + \\
& + \sum_{k_1, k_2, k_3, k_4} \left[-\delta_{i_1, i_2, i_3}^{k_1, k_2, k_3} \mathcal{X}_{y^3}^{k_4} \right] y_{k_1} y_{k_2} y_{k_3} y_{k_4} + \\
& + \sum_{k_1, k_2} \left[\delta_{i_1, i_2}^{k_1, k_2} \mathcal{Y}_{x^{i_3} y} + \delta_{i_3, i_1}^{k_1, k_2} \mathcal{Y}_{x^{i_2} y} + \delta_{i_2, i_3}^{k_1, k_2} \mathcal{Y}_{x^{i_1} y} - \right. \\
& \quad \left. - \delta_{i_1}^{k_1} \mathcal{X}_{x^{i_2} x^{i_3}}^{k_2} - \delta_{i_2}^{k_1} \mathcal{X}_{x^{i_1} x^{i_3}}^{k_2} - \delta_{i_3}^{k_1} \mathcal{X}_{x^{i_1} x^{i_2}}^{k_2} \right] y_{k_1, k_2} + \\
& + \sum_{k_1, k_2, k_3} \left[\delta_{i_1, i_2, i_3}^{k_1, k_2, k_3} \mathcal{Y}_{y^2} + \delta_{i_1, i_2, i_3}^{k_3, k_1, k_2} \mathcal{Y}_{y^2} + \delta_{i_1, i_2, i_3}^{k_2, k_3, k_1} \mathcal{Y}_{y^2} - \right. \\
& \quad - \delta_{i_1, i_2}^{k_1, k_2} \mathcal{X}_{x^{i_3} y}^{k_3} - \delta_{i_1, i_2}^{k_3, k_1} \mathcal{X}_{x^{i_3} y}^{k_2} - \delta_{i_1, i_2}^{k_2, k_3} \mathcal{X}_{x^{i_3} y}^{k_1} - \\
& \quad - \delta_{i_1, i_3}^{k_1, k_2} \mathcal{X}_{x^{i_2} y}^{k_3} - \delta_{i_1, i_3}^{k_3, k_1} \mathcal{X}_{x^{i_2} y}^{k_2} - \delta_{i_1, i_3}^{k_2, k_3} \mathcal{X}_{x^{i_2} y}^{k_1} - \\
& \quad \left. - \delta_{i_2, i_3}^{k_1, k_2} \mathcal{X}_{x^{i_1} y}^{k_3} - \delta_{i_2, i_3}^{k_3, k_1} \mathcal{X}_{x^{i_1} y}^{k_2} - \delta_{i_2, i_3}^{k_2, k_3} \mathcal{X}_{x^{i_1} y}^{k_1} \right] y_{k_1} y_{k_2, k_3} + \\
& + \sum_{k_1, k_2, k_3, k_4} \left[-\delta_{i_1, i_2, i_3}^{k_1, k_2, k_3} \mathcal{X}_{y^2}^{k_4} - \delta_{i_1, i_2, i_3}^{k_2, k_3, k_1} \mathcal{X}_{y^2}^{k_4} - \delta_{i_1, i_2, i_3}^{k_3, k_2, k_1} \mathcal{X}_{y^2}^{k_4} - \right. \\
& \quad \left. - \delta_{i_1, i_2, i_3}^{k_3, k_4, k_1} \mathcal{X}_{y^2}^{k_2} - \delta_{i_1, i_2, i_3}^{k_3, k_1, k_4} \mathcal{X}_{y^2}^{k_2} - \delta_{i_1, i_2, i_3}^{k_1, k_3, k_4} \mathcal{X}_{y^2}^{k_2} \right] y_{k_1} y_{k_2} y_{k_3, k_4} + \\
& + \sum_{k_1, k_2, k_3, k_4} \left[-\delta_{i_1, i_2, i_3}^{k_1, k_2, k_3} \mathcal{X}_y^{k_4} - \delta_{i_1, i_2, i_3}^{k_2, k_3, k_1} \mathcal{X}_y^{k_4} - \delta_{i_1, i_2, i_3}^{k_3, k_1, k_2} \mathcal{X}_y^{k_4} \right] y_{k_1, k_2} y_{k_3, k_4} + \\
& + \sum_{k_1, k_2, k_3} \left[\delta_{i_1, i_2, i_3}^{k_1, k_2, k_3} \mathcal{Y}_y - \delta_{i_1, i_2}^{k_1, k_2} \mathcal{X}_{x^{i_3}}^{k_3} - \delta_{i_1, i_3}^{k_1, k_2} \mathcal{X}_{x^{i_2}}^{k_3} - \delta_{i_2, i_3}^{k_1, k_2} \mathcal{X}_{x^{i_1}}^{k_3} \right] y_{k_1, k_2, k_3} + \\
& + \sum_{k_1, k_2, k_3, k_4} \left[-\delta_{i_1, i_2, i_3}^{k_1, k_2, k_3} \mathcal{X}_y^{k_4} - \delta_{i_1, i_2, i_3}^{k_4, k_1, k_2} \mathcal{X}_y^{k_3} - \delta_{i_1, i_2, i_3}^{k_3, k_4, k_1} \mathcal{X}_y^{k_2} - \delta_{i_1, i_2, i_3}^{k_2, k_3, k_4} \mathcal{X}_y^{k_1} \right] y_{k_1} y_{k_2, k_3, k_4}.
\end{aligned}$$

3.25. Comments, analysis and induction. First of all, by comparing this expression of $\mathbf{Y}_{i_1, i_2, i_3}$ with the expression (2.8) of \mathbf{Y}_3 , we easily guess a part of the (inductional) dictionary between the cases $n = 1$ and the case $n \geq 1$. For instance, the three monomials $[\cdot](y_1)^3$, $[\cdot]y_1 y_2$ and $[\cdot](y_1)^2 y_2$ in \mathbf{Y}_3 are replaced in $\mathbf{Y}_{i_1, i_2, i_3}$ by the following three sums:

$$(3.26) \quad \sum_{k_1, k_2, k_3} [\cdot] y_{k_1} y_{k_2} y_{k_3}, \quad \sum_{k_1, k_2, k_3} [\cdot] y_{k_1} y_{k_2, k_3}, \quad \text{and} \quad \sum_{k_1, k_2, k_3, k_4} [\cdot] y_{k_1} y_{k_2} y_{k_3, k_4}.$$

Similar formal correspondences may be observed for all the monomials of Y_1 , Y_{i_1} , of Y_2 , Y_{i_1, i_2} and of Y_3 , Y_{i_1, i_2, i_3} . Generally and inductively speaking, the monomial

$$(3.27) \quad [\cdot] (y_{\lambda_1})^{\mu_1} \cdots (y_{\lambda_d})^{\mu_d}$$

appearing in the expression (2.25) of Y_κ should be replaced by a certain multiple sum generalizing (3.26). However, it is necessary to think, to pause and to search for an appropriate formalism before writing down the desired multiple sum.

The jet variable y_{λ_1} should be replaced by a jet variable corresponding to a λ_1 -th partial derivative, say $y_{k_1, \dots, k_{\lambda_1}}$, where $k_1, \dots, k_{\lambda_1} = 1, \dots, n$. For the moment, to simplify the discussion, we leave out the presence of a sum of the form $\sum_{k_1, \dots, k_{\lambda_1}}$. The μ_1 -th power $(y_{\lambda_1})^{\mu_1}$ should be replaced *not* by $(y_{k_1, \dots, k_{\lambda_1}})^{\mu_1}$, but by a product of μ_1 different jet variables $y_{k_1, \dots, k_{\lambda_1}}$ of length λ , *with all indices $k_\alpha = 1, \dots, n$ being distinct*. This rule may be confirmed by inspecting the expressions of Y_{i_1} , of Y_{i_1, i_2} and of Y_{i_1, i_2, i_3} . So $y_{k_1, \dots, k_{\lambda_1}}$ should be developed as a product of the form

$$(3.28) \quad y_{k_1, \dots, k_{\lambda_1}} y_{k_{\lambda_1+1}, \dots, k_{2\lambda_1}} \cdots y_{k_{(\mu_1-1)\lambda_1+1}, \dots, k_{\mu_1\lambda_1}},$$

where

$$(3.29) \quad k_1, \dots, k_{\lambda_1}, \dots, k_{\mu_1\lambda_1} = 1, \dots, n.$$

Consider now the product $(y_{\lambda_1})^{\mu_1} (y_{\lambda_2})^{\mu_2}$. How should it develop in the case of several independent variables? For instance, in the expression of Y_{i_1, i_2, i_3} , we have developed the product $(y_1)^2 y_2$ as $y_{k_1} y_{k_2} y_{k_3, k_4}$. Thus, a reasonable proposal of formalism would be that the product $(y_{\lambda_1})^{\mu_1} (y_{\lambda_2})^{\mu_2}$ should be developed as a product of the form

$$(3.30) \quad y_{k_1, \dots, k_{\lambda_1}} y_{k_{\lambda_1+1}, \dots, k_{2\lambda_1}} \cdots y_{k_{(\mu_1-1)\lambda_1+1}, \dots, k_{\mu_1\lambda_1}} \\ y_{k_{\mu_1\lambda_1+1}, \dots, k_{\mu_1\lambda_1+\lambda_2}} \cdots y_{k_{\mu_1\lambda_1+(\mu_2-1)\lambda_2+1}, \dots, k_{\mu_1\lambda_1+\mu_2\lambda_2}},$$

where

$$(3.31) \quad k_1, \dots, k_{\lambda_1}, \dots, k_{\mu_1\lambda_1}, \dots, k_{\mu_1\lambda_1+\mu_2\lambda_2} = 1, \dots, n.$$

However, when trying to write down the development of the general monomial $(y_{\lambda_1})^{\mu_1} (y_{\lambda_2})^{\mu_2} \cdots (y_{\lambda_d})^{\mu_d}$, we would obtain the complicated product

$$(3.32) \quad y_{k_1, \dots, k_{\lambda_1}} y_{k_{\lambda_1+1}, \dots, k_{2\lambda_1}} \cdots y_{k_{(\mu_1-1)\lambda_1+1}, \dots, k_{\mu_1\lambda_1}} \\ y_{k_{\mu_1\lambda_1+1}, \dots, k_{\mu_1\lambda_1+\lambda_2}} \cdots y_{k_{\mu_1\lambda_1+(\mu_2-1)\lambda_2+1}, \dots, k_{\mu_1\lambda_1+\mu_2\lambda_2}} \\ \dots \dots \dots \\ y_{k_{\mu_1\lambda_1+\dots+\mu_{d-1}\lambda_{d-1}+1}, \dots, k_{\mu_1\lambda_1+\dots+\mu_{d-1}\lambda_{d-1}+\lambda_d}} \cdots \\ \cdots y_{k_{\mu_1\lambda_1+\dots+\mu_{d-1}\lambda_{d-1}+(\mu_d-1)\lambda_d+1}, \dots, k_{\mu_1\lambda_1+\dots+\mu_d\lambda_d}}.$$

Essentially, this product is still readable. However, in it, some of the integers k_α have a too long index α , often involving a sum. Such a length of α would be very inconvenient in writing down and in reading the general Kronecker symbols $\delta_{i_1, \dots, i_\lambda}^{k_{\alpha_1}, \dots, k_{\alpha_\lambda}}$ which should appear in the final expression of Y_{i_1, \dots, i_κ} . One should read in advance Theorem 3.73 below to observe the presence of such multiple Kronecker symbols. Consequently, for $\alpha = 1, \dots, \mu_1\lambda_1, \dots, \mu_1\lambda_1 + \dots + \mu_d\lambda_d$, we have to denote the indices k_α differently.

Notational Convention 3.33. We denote d collection of μ_d groups of λ_d (a priori distinct) integers $k_\alpha = 1, \dots, n$ by

$$(3.34) \quad \underbrace{\underbrace{k_{1:1:1}, \dots, k_{1:1:\lambda_1}}_{\lambda_1}, \dots, \underbrace{k_{1:\mu_1:1}, \dots, k_{1:\mu_1:\lambda_1}}_{\lambda_1}}_{\mu_1}$$

$$\underbrace{\underbrace{k_{2:1:1}, \dots, k_{2:1:\lambda_2}}_{\lambda_2}, \dots, \underbrace{k_{2:\mu_2:1}, \dots, k_{2:\mu_2:\lambda_2}}_{\lambda_2}}_{\mu_2}$$

.....

$$\underbrace{\underbrace{k_{d:1:1}, \dots, k_{d:1:\lambda_d}}_{\lambda_d}, \dots, \underbrace{k_{d:\mu_d:1}, \dots, k_{d:\mu_d:\lambda_d}}_{\lambda_d}}_{\mu_d}$$

Correspondingly, we identify the set

$$(3.35) \quad \{1, \dots, \lambda_1, \dots, \mu_1 \lambda_1, \dots, \mu_1 \lambda_1 + \mu_2 \lambda_2, \dots, \mu_1 \lambda_1 + \mu_2 \lambda_2 + \dots + \mu_d \lambda_d\}$$

of all integers α from 1 to $\mu_1 \lambda_1 + \mu_2 \lambda_2 + \dots + \mu_d \lambda_d$ with the following specific set

$$(3.36) \quad \underbrace{\underbrace{\{1:1:1, \dots, 1:1:\lambda_1, \dots, 1:\mu_1:\lambda_1, \dots, 2:\mu_2:\lambda_2, \dots, d:\mu_d:\lambda_d\}}_{\lambda_1}}_{\mu_1 \lambda_1}$$

$$\underbrace{\hspace{10em}}_{\mu_1 \lambda_1 + \mu_2 \lambda_2}$$

$$\underbrace{\hspace{15em}}_{\mu_1 \lambda_1 + \mu_2 \lambda_2 + \dots + \mu_d \lambda_d}$$

written in a lexicographic way which emphasizes clearly the subdivision in d collections of μ_d groups of λ_d integers.

With this notation at hand, we see that the development, in several independent variables, of the general monomial $(y_{\lambda_1})^{\mu_1} \dots (y_{\lambda_d})^{\mu_d}$, may be written as follows:

$$(3.37) \quad y_{k_{1:1:1}, \dots, k_{1:1:\lambda_1}} \dots y_{k_{1:\mu_1:1}, \dots, k_{1:\mu_1:\lambda_1}} \dots y_{k_{d:1:1}, \dots, k_{d:1:\lambda_d}} \dots y_{k_{d:\mu_d:1}, \dots, k_{d:\mu_d:\lambda_d}}$$

Formally speaking, this expression is better than (3.32). Using product symbols, we may even write it under the slightly more compact form

$$(3.38) \quad \prod_{1 \leq \nu_1 \leq \mu_1} y_{k_{1:\nu_1:1}, \dots, k_{1:\nu_1:\lambda_1}} \dots \prod_{1 \leq \nu_d \leq \mu_d} y_{k_{d:\nu_d:1}, \dots, k_{d:\nu_d:\lambda_d}}$$

Now that we have translated the monomial, we may add all the summation symbols: the general expression of Y_κ (which generalizes our three previous examples (3.26)) will be of the form:

$$(3.39) \quad Y_\kappa = \mathcal{Y}_{x^{i_1} \dots x^{i_\kappa}} + \sum_{d=1}^{\kappa+1} \sum_{1 \leq \lambda_1 < \dots < \lambda_d \leq \kappa} \sum_{\mu_1 \geq 1, \dots, \mu_d \geq 1} \sum_{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d \leq \kappa+1}$$

$$\sum_{k_{1:1:1}, \dots, k_{1:1:\lambda_1}=1}^n \dots \sum_{k_{1:\mu_1:1}, \dots, k_{1:\mu_1:\lambda_1}=1}^n \dots \sum_{k_{d:1:1}, \dots, k_{d:1:\lambda_d}=1}^n \dots \sum_{k_{d:\mu_d:1}, \dots, k_{d:\mu_d:\lambda_d}=1}^n$$

$$[?] \prod_{1 \leq \nu_1 \leq \mu_1} y_{k_{1:\nu_1:1}, \dots, k_{1:\nu_1:\lambda_1}} \dots \prod_{1 \leq \nu_d \leq \mu_d} y_{k_{d:\nu_d:1}, \dots, k_{d:\nu_d:\lambda_d}}$$

From now on, up to the end of the article, to be very precise, we will restate the bounds $\sum_{k=1}^n$ of all the previously abbreviated sums \sum_k . This is justified by the fact that, since we shall deal in Section 5 below simultaneously with several independent variables (x^1, \dots, x^n) and with several dependent variables (y^1, \dots, y^m) , we shall encounter sums $\sum_{l=1}^m$, not to be confused with sums $\sum_{k=1}^n$.

3.40. Combinatorics of the Kronecker symbols. Our next task is to determine what appears inside the brackets [?] of the above equation. We will treat this rather delicate question very progressively. Inductively, we have to guess how we may pass from the bracketed term of (2.25), namely from

$$(3.41) \quad \left[\frac{\kappa \cdots (\kappa - \mu_1 \lambda_1 - \cdots - \mu_d \lambda_d + 1)}{(\lambda_1!)^{\mu_1} \mu_1! \cdots (\lambda_d!)^{\mu_d} \mu_d!} \cdot \mathcal{Y}_{x^{\kappa - \mu_1 \lambda_1 - \cdots - \mu_d \lambda_d} y^{\mu_1 + \cdots + \mu_d}} - \frac{\kappa \cdots (\kappa - \mu_1 \lambda_1 - \cdots - \mu_d \lambda_d + 2)(\mu_1 \lambda_1 + \cdots + \mu_d \lambda_d)}{(\lambda_1!)^{\mu_1} \mu_1! \cdots (\lambda_d!)^{\mu_d} \mu_d!} \cdot \mathcal{X}_{x^{\kappa - \mu_1 \lambda_1 - \cdots - \mu_d \lambda_d + 1} y^{\mu_1 + \cdots + \mu_d - 1}} \right],$$

to the corresponding (still unknown) bracketed term [?].

First of all, we examine the following term, extracted from the complete expression of $\mathbf{Y}_{i_1, i_2, i_3}$ (first line of (3.24)):

$$(3.42) \quad \sum_{k_1=1}^n \left[\delta_{i_1}^{k_1} \mathcal{Y}_{x^{i_2} x^{i_3} y} + \delta_{i_2}^{k_1} \mathcal{Y}_{x^{i_1} x^{i_3} y} + \delta_{i_3}^{k_1} \mathcal{Y}_{x^{i_1} x^{i_2} y} - \mathcal{X}_{x^{i_1} x^{i_2} x^{i_3}} \right] y_{k_1}.$$

Here, the coefficient $[3 \mathcal{Y}_{x^2 y} - \mathcal{X}_{x^3}]$ of the monomial y_1 in \mathbf{Y}_3 is replaced by the above bracketed terms.

Let us precisely analyze the combinatorics. Here, \mathcal{X}_{x^3} is replaced by $\mathcal{X}_{x^{i_1} x^{i_2} x^{i_3}}$, where the lower indices i_1, i_2, i_3 come from $\mathbf{Y}_{i_1, i_2, i_3}$ and where the upper index k_1 is the summation index. Also, the integer 3 in $3 \mathcal{Y}_{x^2 y}$ is replaced by a sum of exactly three terms, each involving a single Kronecker symbol δ_i^k , in which the lower index is always an index $i = i_1, i_2, i_3$ and in which the upper index is always equal to the summation index k_1 . By the way, more generally, we immediately observe that all the successive positive integers

$$(3.43) \quad 1, 3, 1, 3, 3, 1, 3, 1, 3, 3, 3, 9, 6, 3, 1, 3, 4$$

appearing in the formula (2.8) for \mathbf{Y}_3 are replaced, in the formula (3.24) for $\mathbf{Y}_{i_1, i_2, i_3}$, by sums of exactly the same number of terms involving Kronecker symbols. This observation will be a precious guide. Finally, in the symbol $\delta_i^{k_1}$, if i is chosen among the set $\{i_1, i_2, i_3\}$, for instance if $i = i_1$, it follows that the development of $\mathcal{Y}_{x^2 y}$ necessarily involves the remaining indices, for instance $\mathcal{Y}_{x^{i_2} x^{i_3} y}$. Since there are three choices for $i = i_1, i_2, i_3$, we recover the number 3.

Next, comparing $[\mathcal{Y}_{yy} - 2 \mathcal{X}_{xy}] (y_1)^2$ with the term

$$(3.44) \quad \sum_{k_1, k_2=1}^n \left[\delta_{i_1, i_2}^{k_1, k_2} \mathcal{Y}_{yy} - \delta_{i_1}^{k_1} \mathcal{X}_{x^{i_2} y} - \delta_{i_2}^{k_1} \mathcal{X}_{x^{i_1} y} \right] y_{k_1} y_{k_2},$$

extracted from the complete expression of \mathbf{Y}_{i_1, i_2} (second line of (3.18)), we learn and we guess that the number of Kronecker symbols before $\mathcal{Y}_{x^\gamma y^\delta}$ must be equal to the number of indices k_α minus γ . This rule is confirmed by examining the term (second and third line

of (3.24))

$$(3.45) \quad \sum_{k_1, k_2} \left[\delta_{i_1, i_2}^{k_1, k_2} \mathcal{Y}_{x^{i_3} y^2} + \delta_{i_3, i_1}^{k_1, k_2} \mathcal{Y}_{x^{i_2} y^2} + \delta_{i_2, i_3}^{k_1, k_2} \mathcal{Y}_{x^{i_1} y^2} - \right. \\ \left. - \delta_{i_1}^{k_1} \mathcal{X}_{x^{i_2} x^{i_3} y}^{k_2} - \delta_{i_2}^{k_1} \mathcal{X}_{x^{i_1} x^{i_3} y}^{k_2} - \delta_{i_3}^{k_1} \mathcal{X}_{x^{i_1} x^{i_2} y}^{k_2} \right] y_{k_1} y_{k_2},$$

developing $[3 \mathcal{Y}_{xy^2} - 3 \mathcal{X}_{x^2 y}] (y_1)^2$.

Also, we may examine the following term

$$(3.46) \quad \sum_{k_1, k_2=1}^n \left[\delta_{i_1, i_2}^{k_1, k_2} \mathcal{Y}_{x^{i_3} x^{i_4} y^2} + \delta_{i_1, i_3}^{k_1, k_2} \mathcal{Y}_{x^{i_2} x^{i_4} y^2} + \delta_{i_1, i_4}^{k_1, k_2} \mathcal{Y}_{x^{i_2} x^{i_3} y^2} + \right. \\ \left. + \delta_{i_2, i_3}^{k_1, k_2} \mathcal{Y}_{x^{i_1} x^{i_4} y^2} + \delta_{i_2, i_4}^{k_1, k_2} \mathcal{Y}_{x^{i_1} x^{i_3} y^2} + \delta_{i_3, i_4}^{k_1, k_2} \mathcal{Y}_{x^{i_1} x^{i_2} y^2} - \right. \\ \left. - \delta_{i_1}^{k_1} \mathcal{X}_{x^{i_2} x^{i_3} x^{i_4} y}^{k_1} - \delta_{i_2}^{k_1} \mathcal{X}_{x^{i_1} x^{i_2} x^{i_3} y}^{k_1} - \delta_{i_3}^{k_1} \mathcal{X}_{x^{i_1} x^{i_2} x^{i_4} y}^{k_1} - \right. \\ \left. - \delta_{i_4}^{k_1} \mathcal{X}_{x^{i_1} x^{i_2} x^{i_3} y}^{k_1} \right] y_{k_1} y_{k_2},$$

extracted from $\mathbf{Y}_{i_1, i_2, i_3, i_4}$ and developing $[6 \mathcal{Y}_{x^2 y^2} - 4 \mathcal{X}_{x^3 y}] (y_1)^2$. We would like to mention that we have not written the complete expression of $\mathbf{Y}_{i_1, i_2, i_3, i_4}$, because it would cover two and a half printed pages.

By inspecting the way how the indices are permuted in the multiple Kronecker symbols of the first two lines of this expression (3.46), we observe that the six terms correspond exactly to the six possible choices of two complementary ordered couples of integers in the set $\{1, 2, 3, 4\}$, namely

$$(3.47) \quad \begin{array}{lll} \{1, 2\} \cup \{3, 4\}, & \{1, 3\} \cup \{2, 4\}, & \{1, 4\} \cup \{2, 3\}, \\ \{2, 3\} \cup \{1, 4\}, & \{2, 4\} \cup \{1, 3\}, & \{3, 4\} \cup \{1, 2\}. \end{array}$$

At this point, we start to devise the general combinatorics. Before proceeding further, we need some notation.

3.48. Permutation groups. For every $p \in \mathbb{N}$ with $p \geq 1$, we denote by \mathfrak{S}_p the full permutation group of the set $\{1, 2, \dots, p-1, p\}$. Its cardinal equals $p!$. The letters σ and τ will be used to denote an element of \mathfrak{S}_p . If $p \geq 2$, and if $q \in \mathbb{N}$ satisfies $1 \leq q \leq p-1$, we denote by \mathfrak{S}_p^q the subset of permutations $\sigma \in \mathfrak{S}_p$ satisfying the two collections of inequalities

$$(3.49) \quad \sigma(1) < \sigma(2) < \dots < \sigma(q) \quad \text{and} \quad \sigma(q+1) < \sigma(q+2) < \dots < \sigma(p).$$

The cardinal of \mathfrak{S}_p^q equals $C_p^q = \frac{p!}{q! (p-q)!}$.

Lemma 3.50. For $\kappa \geq 1$, the development of (2.20) to several independent variables (x^1, \dots, x^n) is:

$$\begin{aligned}
\mathbf{Y}_{i_1, i_2, \dots, i_\kappa} &= \mathcal{Y}_{x^{i_1} x^{i_2} \dots x^{i_\kappa}} + \sum_{k_1=1}^n \left[\sum_{\tau \in \mathfrak{S}_\kappa^1} \delta_{i_\tau(1)}^{k_1} \mathcal{Y}_{x^{i_\tau(2)} \dots x^{i_\tau(\kappa)} y} - \mathcal{X}_{x^{i_1} x^{i_2} \dots x^{i_\kappa}}^{k_1} \right] y_{k_1} + \\
&+ \sum_{k_1, k_2=1}^n \left[\sum_{\tau \in \mathfrak{S}_\kappa^2} \delta_{i_\tau(1), i_\tau(2)}^{k_1, k_2} \mathcal{Y}_{x^{i_\tau(3)} \dots x^{i_\tau(\kappa)} y^2} - \sum_{\tau \in \mathfrak{S}_\kappa^1} \delta_{i_\tau(1)}^{k_1} \mathcal{X}_{x^{i_\tau(2)} \dots x^{i_\tau(\kappa)} y}^{k_2} \right] y_{k_1} y_{k_2} + \\
&+ \sum_{k_1, k_2, k_3=1}^n \left[\sum_{\tau \in \mathfrak{S}_\kappa^3} \delta_{i_\tau(1), i_\tau(2), i_\tau(3)}^{k_1, k_2, k_3} \mathcal{Y}_{x^{i_\tau(4)} \dots x^{i_\tau(\kappa)} y^3} - \right. \\
(3.51) \quad &\left. - \sum_{\tau \in \mathfrak{S}_\kappa^2} \delta_{i_\tau(1), i_\tau(2)}^{k_1, k_2} \mathcal{X}_{x^{i_\tau(3)} \dots x^{i_\tau(\kappa)} y^2}^{k_3} \right] y_{k_1} y_{k_2} y_{k_3} + \\
&+ \dots + \\
&+ \sum_{k_1, \dots, k_\kappa=1}^n \left[\delta_{i_1, \dots, i_\kappa}^{k_1, \dots, k_\kappa} \mathcal{Y}_{y^\kappa} - \sum_{\tau \in \mathfrak{S}_\kappa^{\kappa-1}} \delta_{i_\tau(1), \dots, i_\tau(\kappa-1)}^{k_1, \dots, k_{\kappa-1}} \mathcal{X}_{x^{i_\tau(\kappa)} y^{\kappa-1}}^{k_\kappa} \right] y_{k_1} \dots y_{k_\kappa} + \\
&+ \sum_{k_1, \dots, k_\kappa, k_{\kappa+1}=1}^n \left[-\delta_{i_1, \dots, i_\kappa}^{k_1, \dots, k_\kappa} \mathcal{X}_{y^\kappa}^{k_{\kappa+1}} \right] y_{k_1} \dots y_{k_\kappa} y_{k_{\kappa+1}} + \text{remainder.}
\end{aligned}$$

Here, the term remainder collects all remaining monomials in the pure jet variables $y_{k_1, \dots, k_\lambda}$.

3.52. Continuation. Thus, we have devised how the part of $\mathbf{Y}_{i_1, \dots, i_\kappa}$ which involves only the jet variables y_{k_α} must be written. To proceed further, we shall examine the following term, extracted from $\mathbf{Y}_{i_1, i_2, i_3}$ (lines 12 and 13 of (3.24))

$$\begin{aligned}
(3.53) \quad &\sum_{k_1, k_2, k_3, k_4} \left[-\delta_{i_1, i_2, i_3}^{k_1, k_2, k_3} \mathcal{X}_{y^2}^{k_4} - \delta_{i_1, i_2, i_3}^{k_2, k_3, k_1} \mathcal{X}_{y^2}^{k_4} - \delta_{i_1, i_2, i_3}^{k_3, k_2, k_1} \mathcal{X}_{y^2}^{k_4} - \right. \\
&\left. - \delta_{i_1, i_2, i_3}^{k_3, k_4, k_1} \mathcal{X}_{y^2}^{k_2} - \delta_{i_1, i_2, i_3}^{k_3, k_1, k_4} \mathcal{X}_{y^2}^{k_2} - \delta_{i_1, i_2, i_3}^{k_1, k_3, k_4} \mathcal{X}_{y^2}^{k_2} \right] y_{k_1} y_{k_2} y_{k_3, k_4},
\end{aligned}$$

which develops the term $[-6 \mathcal{X}_{y^2}] (y_1)^2 y_2$ of \mathbf{Y}_3 (third line of (2.8)). During the computation which led us to the final expression (3.24), we organized the formula in order that, in the six Kronecker symbols, the lower indices i_1, i_2, i_3 are all written in the same order. But then, *what is the rule for the appearance of the four upper indices k_1, k_2, k_3, k_4 ?*

In April 2001, we discovered the rule by inspecting both (3.53) and the following complicated term, extracted from the complete expression of $\mathbf{Y}_{i_1, i_2, i_3, i_4}$ written in one of our manuscripts:

$$\begin{aligned}
(3.54) \quad &\sum_{k_1, k_2, k_3} \left[\delta_{i_1, i_2, i_3}^{k_1, k_2, k_3} \mathcal{Y}_{x^{i_4} y^2} + \delta_{i_1, i_2, i_3}^{k_2, k_1, k_3} \mathcal{Y}_{x^{i_4} y^2} + \delta_{i_1, i_2, i_3}^{k_2, k_3, k_1} \mathcal{Y}_{x^{i_4} y^2} + \right. \\
&+ \delta_{i_1, i_2, i_4}^{k_1, k_2, k_3} \mathcal{Y}_{x^{i_3} y^2} + \delta_{i_1, i_2, i_4}^{k_2, k_1, k_3} \mathcal{Y}_{x^{i_3} y^2} + \delta_{i_1, i_2, i_4}^{k_2, k_3, k_1} \mathcal{Y}_{x^{i_3} y^2} + \\
&+ \delta_{i_1, i_3, i_4}^{k_1, k_2, k_3} \mathcal{Y}_{x^{i_2} y^2} + \delta_{i_1, i_3, i_4}^{k_2, k_1, k_3} \mathcal{Y}_{x^{i_2} y^2} + \delta_{i_1, i_3, i_4}^{k_2, k_3, k_1} \mathcal{Y}_{x^{i_2} y^2} + \\
&+ \delta_{i_2, i_3, i_4}^{k_1, k_2, k_3} \mathcal{Y}_{x^{i_1} y^2} + \delta_{i_2, i_3, i_4}^{k_2, k_1, k_3} \mathcal{Y}_{x^{i_1} y^2} + \delta_{i_2, i_3, i_4}^{k_2, k_3, k_1} \mathcal{Y}_{x^{i_1} y^2} - \\
&\left. - \delta_{i_1, i_2}^{k_1, k_2} \mathcal{X}_{x^{i_3} x^{i_4} y}^{k_3} - \delta_{i_1, i_2}^{k_2, k_1} \mathcal{X}_{x^{i_3} x^{i_4} y}^{k_3} - \delta_{i_1, i_2}^{k_2, k_3} \mathcal{X}_{x^{i_3} x^{i_4} y}^{k_1} - \right.
\end{aligned}$$

$$\begin{aligned}
& -\delta_{i_1, i_3}^{k_1, k_2} \mathcal{X}_{x^{i_2} x^{i_4} y}^{k_3} - \delta_{i_1, i_3}^{k_2, k_1} \mathcal{X}_{x^{i_2} x^{i_4} y}^{k_3} - \delta_{i_1, i_3}^{k_2, k_3} \mathcal{X}_{x^{i_2} x^{i_4} y}^{k_1} - \\
& -\delta_{i_1, i_4}^{k_1, k_2} \mathcal{X}_{x^{i_2} x^{i_3} y}^{k_3} - \delta_{i_1, i_4}^{k_2, k_1} \mathcal{X}_{x^{i_2} x^{i_3} y}^{k_3} - \delta_{i_1, i_4}^{k_2, k_3} \mathcal{X}_{x^{i_2} x^{i_3} y}^{k_1} - \\
& -\delta_{i_2, i_3}^{k_1, k_2} \mathcal{X}_{x^{i_1} x^{i_4} y}^{k_3} - \delta_{i_2, i_3}^{k_2, k_1} \mathcal{X}_{x^{i_1} x^{i_4} y}^{k_3} - \delta_{i_2, i_3}^{k_2, k_3} \mathcal{X}_{x^{i_1} x^{i_4} y}^{k_1} - \\
& -\delta_{i_2, i_4}^{k_1, k_2} \mathcal{X}_{x^{i_1} x^{i_3} y}^{k_3} - \delta_{i_2, i_4}^{k_2, k_1} \mathcal{X}_{x^{i_1} x^{i_3} y}^{k_3} - \delta_{i_2, i_4}^{k_2, k_3} \mathcal{X}_{x^{i_1} x^{i_3} y}^{k_1} - \\
& -\delta_{i_3, i_4}^{k_1, k_2} \mathcal{X}_{x^{i_1} x^{i_2} y}^{k_3} - \delta_{i_3, i_4}^{k_2, k_1} \mathcal{X}_{x^{i_1} x^{i_2} y}^{k_3} - \delta_{i_3, i_4}^{k_2, k_3} \mathcal{X}_{x^{i_1} x^{i_2} y}^{k_1} \Big] y_{k_1} y_{k_2, k_3}.
\end{aligned}$$

This sum develops the term $[12 \mathcal{Y}_{xy^2} - 18 \mathcal{X}_{x^2y}] y_1 y_2$ of \mathbf{Y}_3 (third line of (2.9)). Let us explain what are the formal rules.

In the bracketed terms of (3.53), there are no permutation of the indices i_1, i_2, i_3 , but there is a certain unknown subset of all the permutations of the four indices k_1, k_2, k_3, k_4 . In the bracketed terms of (3.54), two combinatorics are present:

- there are some permutations of the indices i_1, i_2, i_3, i_4 and
- there are some permutations of the indices k_1, k_2, k_3 .

Here, the permutations of the indices i_1, i_2, i_3, i_4 are easily guessed, since they are the same as the permutations which were introduced in §3.48 above. Indeed, in the first four lines of (3.54), we see the four decompositions

$$(3.55) \quad \{i_1, i_2, i_3\} \cup \{i_4\}, \quad \{i_1, i_2, i_4\} \cup \{i_3\}, \quad \{i_1, i_3, i_4\} \cup \{i_2\}, \quad \{i_2, i_3, i_4\} \cup \{i_1\},$$

of the set $\{i_1, i_2, i_3, i_4\}$, and in the last six lines of (3.54), we see the six decompositions

$$(3.56) \quad \begin{array}{ccc} \{i_1, i_2\} \cup \{i_3, i_4\}, & \{i_1, i_3\} \cup \{i_2, i_4\}, & \{i_1, i_4\} \cup \{i_2, i_3\}, \\ \{i_2, i_3\} \cup \{i_1, i_4\}, & \{i_2, i_4\} \cup \{i_1, i_3\}, & \{i_3, i_4\} \cup \{i_1, i_2\}, \end{array}$$

so that (3.54) may be written under the form

$$(3.57) \quad \sum_{k_1, k_2, k_3} \left[\sum_{\tau \in \mathfrak{S}_4^?} \sum_{\sigma \in ?} \delta_{i_{\tau(1)}, i_{\tau(2)}, i_{\tau(3)}}^{k_{\tau(1)}, k_{\tau(2)}, k_{\tau(3)}} \mathcal{Y}_{x^{i_{\tau(4)} y^2}} - \sum_{\tau \in \mathfrak{S}_4^?} \sum_{\sigma \in ?} \delta_{i_{\tau(1)}, i_{\tau(2)}}^{k_{\tau(1)}, k_{\tau(2)}} \mathcal{X}_{x^{i_{\tau(3)} x^{i_{\tau(4)} y}}^{k_{\tau(3)}} \right] y_{k_1} y_{k_2, k_3},$$

where in the two above sums $\sum_{\sigma \in ?}$, the letter σ denotes a permutation of the set $\{1, 2, 3\}$ and where the sign $?$ refers to two (still unknown) subset of the full permutation group \mathfrak{S}_3 . *The only remaining question is to determine how the indices k_α are permuted in (3.53) and in (3.54).*

The answer may be guessed by looking at the permutations of the set $\{k_1, k_2, k_3, k_4\}$ which stabilize the monomial $y_{k_1} y_{k_2} y_{k_3, k_4}$ in (3.53): we clearly have the following four symmetry relations between monomials:

$$(3.58) \quad y_{k_1} y_{k_2} y_{k_3, k_4} \equiv y_{k_2} y_{k_1} y_{k_3, k_4} \equiv y_{k_1} y_{k_2} y_{k_4, k_3} \equiv y_{k_2} y_{k_1} y_{k_4, k_3},$$

and nothing more. Then the number 6 of bracketed terms in (3.53) is exactly equal to the cardinal $24 = 4!$ of the full permutation group of the set $\{k_1, k_2, k_3, k_4\}$ divided by the number 4 of these symmetry relations. The set of permutations σ of $\{1, 2, 3, 4\}$ satisfying these symmetry relations

$$(3.59) \quad y_{k_{\sigma(1)}} y_{k_{\sigma(2)}} y_{k_{\sigma(3)}, k_{\sigma(4)}} \equiv y_{k_1} y_{k_2} y_{k_3, k_4}$$

constitutes a subgroup of \mathfrak{S}_4 which we will denote by $\mathfrak{H}_4^{(2,1), (1,2)}$. Furthermore, the coset

$$(3.60) \quad \mathfrak{F}_4^{(2,1), (1,2)} := \mathfrak{S}_4 / \mathfrak{H}_4^{(2,1), (1,2)}$$

possesses the six representatives

$$(3.61) \quad \begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & 2 & 3 & 4 \end{pmatrix}, \quad \begin{pmatrix} 1 & 2 & 3 & 4 \\ 2 & 3 & 1 & 4 \end{pmatrix}, \quad \begin{pmatrix} 1 & 2 & 3 & 4 \\ 3 & 2 & 1 & 4 \end{pmatrix}, \\ \begin{pmatrix} 1 & 2 & 3 & 4 \\ 3 & 4 & 1 & 2 \end{pmatrix}, \quad \begin{pmatrix} 1 & 2 & 3 & 4 \\ 3 & 1 & 4 & 2 \end{pmatrix}, \quad \begin{pmatrix} 1 & 2 & 3 & 4 \\ 1 & 3 & 4 & 2 \end{pmatrix},$$

which exactly appear as the permutations of the upper indices of our example (3.53). Of course, the question arises whether the choice of such six representatives in the quotient $\mathfrak{S}_4/\mathfrak{H}_4^{(2,1),(1,2)}$ is legitimate.

Fortunately, we observe that after conjugation by any permutation $\sigma \in \mathfrak{H}_4^{(2,1),(1,2)}$, we do not perturb any of the six terms of (3.53), for instance the third term of (3.53) is not perturbed, as shown by the following computation

$$(3.62) \quad \sum_{k_1, k_2, k_3, k_4} \left[-\delta_{i_1, i_2, i_3}^{k_{\sigma(3)}, k_{\sigma(2)}, k_{\sigma(1)}} \mathcal{X}_{y^2}^{k_{\sigma(4)}} \right] y_{k_1} y_{k_2} y_{k_3, k_4} = \\ = \sum_{k_1, k_2, k_3, k_4} \left[-\delta_{i_1, i_2, i_3}^{k_3, k_2, k_1} \mathcal{X}_{y^2}^{k_{\sigma(4)}} \right] y_{k_{\sigma^{-1}(1)}} y_{k_{\sigma^{-1}(2)}} y_{k_{\sigma^{-1}(3)}, k_{\sigma^{-1}(4)}} \\ = \sum_{k_1, k_2, k_3, k_4} \left[-\delta_{i_1, i_2, i_3}^{k_3, k_2, k_1} \mathcal{X}_{y^2}^{k_{\sigma(4)}} \right] y_{k_1} y_{k_2} y_{k_3, k_4}$$

thanks to the symmetry (3.59). Thus, as expected, the choice of 6 arbitrary representatives $\sigma \in \mathfrak{H}_4^{(2,1),(1,2)}$ in the bracketed terms of (3.53) is free. In conclusion, we have shown that (3.53) may be written under the form:

$$(3.63) \quad \sum_{k_1, k_2, k_3, k_4} \left[- \sum_{\sigma \in \mathfrak{H}_4^{(2,1),(1,2)}} \delta_{i_1, i_2, i_3}^{k_{\sigma(1)}, k_{\sigma(2)}, k_{\sigma(3)}} \mathcal{X}_{y^2}^{k_{\sigma(4)}} \right] y_{k_1} y_{k_2} y_{k_3, k_4},$$

This rule is confirmed by inspecting (3.54) (as well as all the other terms of $\mathbf{Y}_{i_1, i_2, i_3}$ and of $\mathbf{Y}_{i_1, i_2, i_3, i_4}$). Indeed, the permutations σ of the set $\{k_1, k_2, k_3\}$ which stabilize the monomial $y_{k_1} y_{k_2, k_3}$ consist just of the identity permutation and the transposition of k_2 and k_3 . The coset $\mathfrak{S}_3/\mathfrak{H}_3^{(1,1),(1,2)}$ has the three representatives

$$(3.64) \quad \begin{pmatrix} 1 & 2 & 3 \\ 1 & 2 & 3 \end{pmatrix}, \quad \begin{pmatrix} 1 & 2 & 3 \\ 2 & 1 & 3 \end{pmatrix}, \quad \begin{pmatrix} 1 & 2 & 3 \\ 2 & 3 & 1 \end{pmatrix},$$

which appear in the upper index position of each of the ten lines of (3.54). It follows that (3.54) may be written under the form

$$(3.65) \quad \sum_{k_1, k_2, k_3} \left[\sum_{\tau \in \mathfrak{S}_4^3} \sum_{\sigma \in \mathfrak{H}_3^{(1,1),(1,2)}} \delta_{i_{\tau(1)}, i_{\tau(2)}, i_{\tau(3)}}^{k_{\sigma(1)}, k_{\sigma(2)}, k_{\sigma(3)}} \mathcal{Y}_{x^{i_{\tau(4)}} y^2} - \right. \\ \left. - \sum_{\sigma \in \mathfrak{S}_4^2} \sum_{\tau \in \mathfrak{H}_3^{(1,1),(1,2)}} \delta_{i_{\tau(1)}, i_{\tau(2)}}^{k_{\sigma(1)}, k_{\sigma(2)}} \mathcal{X}_{x^{i_{\tau(3)}} x^{i_{\tau(4)}} y} \right] y_{k_1} y_{k_2, k_3}.$$

3.66. General complete expression of $\mathbf{Y}_{i_1, \dots, i_\kappa}$. As in the incomplete expression (3.39) of $\mathbf{Y}_{i_1, \dots, i_\kappa}$, consider integers $1 \leq \lambda_1 < \dots < \lambda_d \leq \kappa$ and $\mu_1 \geq 1, \dots, \mu_d \geq 1$ satisfying $\mu_1 \lambda_1 + \dots + \mu_d \lambda_d \leq \kappa + 1$. By $\mathfrak{H}_{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d}$, we denote the subgroup of permutations $\tau \in$

$\mathfrak{S}_{\mu_1\lambda_1+\dots+\mu_d\lambda_d}$ that leave unchanged the general monomial (3.38), namely that satisfy

$$(3.67) \quad \prod_{1 \leq \nu_1 \leq \mu_1} y_{k_{\sigma(1:\nu_1:1)}, \dots, k_{\sigma(1:\nu_1:\lambda_1)}} \cdots \prod_{1 \leq \nu_d \leq \mu_d} y_{k_{\sigma(d:\nu_d:1)}, \dots, k_{\sigma(d:\nu_d:\lambda_d)}} = \\ = \prod_{1 \leq \nu_1 \leq \mu_1} y_{k_{1:\nu_1:1}, \dots, k_{1:\nu_1:\lambda_1}} \cdots \prod_{1 \leq \nu_d \leq \mu_d} y_{k_{d:\nu_d:1}, \dots, k_{d:\nu_d:\lambda_d}}.$$

The structure of this group may be described as follows. For every $e = 1, \dots, d$, an arbitrary permutation σ of the set

$$(3.68) \quad \underbrace{\{e: 1:1, \dots, e: 1:\lambda_e\}}_{\lambda_e} \underbrace{\{e: 2:1, \dots, e: 2:\lambda_e, \dots, e: \mu_e:1, \dots, e: \mu_e:\lambda_e\}}_{\lambda_e} \underbrace{\hspace{10em}}_{\mu_e}$$

which leaves unchanged the monomial

$$(3.69) \quad \prod_{1 \leq \nu_e \leq \mu_e} y_{k_{\sigma(e:\nu_e:1)}, \dots, k_{\sigma(e:\nu_e:\lambda_e)}} = \prod_{1 \leq \nu_e \leq \mu_e} y_{k_{e:\nu_e:1}, \dots, k_{e:\nu_e:\lambda_e}}.$$

uniquely decomposes as the composition of

- μ_e arbitrary permutations of the μ_e groups of λ_e integers $\{e:\nu_e:1, \dots, e:\nu_e:\lambda_e\}$, of total cardinal $(\lambda_e!)^{\mu_e}$;
- an arbitrary permutation between these μ_e groups, of total cardinal $\mu_e!$.

Consequently

$$(3.70) \quad \text{Card} \left(\mathfrak{H}_{\mu_1\lambda_1+\dots+\mu_d\lambda_d}^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)} \right) = \mu_1!(\lambda_1!)^{\mu_1} \cdots \mu_d!(\lambda_d!)^{\mu_d}.$$

Finally, define the coset

$$(3.71) \quad \mathfrak{F}_{\mu_1\lambda_1+\dots+\mu_d\lambda_d}^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)} := \mathfrak{S}_{\mu_1\lambda_1+\dots+\mu_d\lambda_d} / \mathfrak{H}_{\mu_1\lambda_1+\dots+\mu_d\lambda_d}^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)}$$

with

$$(3.72) \quad \text{Card} \left(\mathfrak{F}_{\mu_1\lambda_1+\dots+\mu_d\lambda_d}^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)} \right) = \frac{\text{Card} \left(\mathfrak{S}_{\mu_1\lambda_1+\dots+\mu_d\lambda_d} \right)}{\text{Card} \left(\mathfrak{H}_{\mu_1\lambda_1+\dots+\mu_d\lambda_d}^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)} \right)} \\ = \frac{(\mu_1\lambda_1 + \dots + \mu_d\lambda_d)!}{\mu_1!(\lambda_1!)^{\mu_1} \cdots \mu_d!(\lambda_d!)^{\mu_d}}.$$

In conclusion, by means of this formalism, we may write down the complete generalization of Theorem 2.24 to several independent variables.

Theorem 3.73. For every $\kappa \geq 1$ and for every choice of κ indices i_1, \dots, i_κ in the set $\{1, 2, \dots, n\}$, the general expression of $\mathbf{Y}_{i_1, \dots, i_\kappa}$ is as follows:

(3.74)

$$\begin{aligned}
 \mathbf{Y}_{i_1, \dots, i_\kappa} = & \mathcal{Y}_{x^{i_1} \dots x^{i_\kappa}} + \sum_{d=1}^{\kappa+1} \sum_{1 \leq \lambda_1 < \dots < \lambda_d \leq \kappa} \sum_{\mu_1 \geq 1, \dots, \mu_d \geq 1} \sum_{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d \leq \kappa+1} \\
 & \sum_{k_1:1:1, \dots, k_{1:1}:\lambda_1=1}^n \dots \sum_{k_1:\mu_1:1, \dots, k_{1:\mu_1}:\lambda_1=1}^n \dots \dots \dots \sum_{k_d:1:1, \dots, k_{d:1}:\lambda_d=1}^n \dots \sum_{k_d:\mu_d:1, \dots, k_{d:\mu_d}:\lambda_d=1}^n \\
 & \left[\sum_{\sigma \in \mathfrak{S}_{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d}}^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)} \sum_{\tau \in \mathfrak{S}_\kappa}^{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d} \right. \\
 & \quad \delta_{i_{\tau(1)}, \dots, i_{\tau(\mu_1 \lambda_1)}, \dots, i_{\tau(\mu_1 \lambda_1 + \dots + \mu_d \lambda_d)}}^{k_{\sigma(1:1:1)}, \dots, k_{\sigma(1:\mu_1:\lambda_1)}, \dots, k_{\sigma(d:\mu_d:\lambda_d)}} \frac{\partial^{\kappa - \mu_1 \lambda_1 - \dots - \mu_d \lambda_d + \mu_1 + \dots + \mu_d} \mathcal{Y}}{\partial x^{i_{\tau(\mu_1 \lambda_1 + \dots + \mu_d \lambda_d + 1)}} \dots \partial x^{i_{\tau(\kappa)}} (\partial y)^{\mu_1 + \dots + \mu_d}} \\
 & \quad - \sum_{\sigma \in \mathfrak{S}_{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d}}^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)} \sum_{\tau \in \mathfrak{S}_\kappa}^{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d - 1} \\
 & \quad \delta_{i_{\tau(1)}, \dots, i_{\tau(\mu_1 \lambda_1)}, \dots, i_{\tau(\mu_1 \lambda_1 + \dots + \mu_d \lambda_d - 1)}}^{k_{\sigma(1:1:1)}, \dots, k_{\sigma(1:\mu_1:\lambda_1)}, \dots, k_{\sigma(d:\mu_d:\lambda_d - 1)}} \frac{\partial^{\kappa - \mu_1 \lambda_1 - \dots - \mu_d \lambda_d + \mu_1 + \dots + \mu_d} \mathcal{X}^{k_{\sigma(d:\mu_d:\lambda_d)}}}{\partial x^{i_{\tau(\mu_1 \lambda_1 + \dots + \mu_d \lambda_d)}} \dots \partial x^{i_{\tau(\kappa)}} (\partial y)^{\mu_1 + \dots + \mu_d - 1}} \\
 & \quad \cdot \prod_{1 \leq \nu_1 \leq \mu_1} y_{k_{1:\nu_1:1}, \dots, k_{1:\nu_1}:\lambda_1} \dots \prod_{1 \leq \nu_d \leq \mu_d} y_{k_{d:\nu_d:1}, \dots, k_{d:\nu_d}:\lambda_d}.
 \end{aligned}$$

Since the fundamental monomials appearing in the last line of (3.74) just above are not independent of each other, this formula has still to be modified a little bit. We refer to Section 6 for details.

3.75. Deduction of a multivariate Faà di Bruno formula. Let $n \in \mathbb{N}$ with $n \geq 1$, let $x = (x^1, \dots, x^n) \in \mathbb{K}^n$, let $g = g(x^1, \dots, x^n)$ be a \mathcal{C}^∞ -smooth function from \mathbb{K}^n to \mathbb{K} , let $y \in \mathbb{K}$ and let $f = f(y)$ be a \mathcal{C}^∞ function from \mathbb{K} to \mathbb{K} . The goal is to obtain an explicit formula for the partial derivatives of the composition $h := f \circ g$, namely $h(x^1, \dots, x^n) := f(g(x^1, \dots, x^n))$. For $\lambda \in \mathbb{N}$ with $\lambda \geq 1$ and for arbitrary indices $i_1, \dots, i_\lambda = 1, \dots, n$, we shall abbreviate the partial derivative $\frac{\partial^\lambda g}{\partial x^{i_1} \dots \partial x^{i_\lambda}}$ by $g_{i_1, \dots, i_\lambda}$ and similarly for $h_{i_1, \dots, i_\lambda}$. The derivative $\frac{d^\lambda f}{dy^\lambda}$ will be abbreviated by f_λ .

Applying the chain rule, we may compute:

$$\begin{aligned}
 h_{i_1} &= f_1 [g_{i_1}], \\
 h_{i_1, i_2} &= f_2 [g_{i_1} g_{i_2}] + f_1 [g_{i_1, i_2}], \\
 h_{i_1, i_2, i_3} &= f_3 [g_{i_1} g_{i_2} g_{i_3}] + f_2 [g_{i_1} g_{i_2, i_3} + g_{i_2} g_{i_1, i_3} + g_{i_3} g_{i_1, i_2}] + f_1 [g_{i_1, i_2, i_3}], \\
 h_{i_1, i_2, i_3, i_4} &= f_4 [g_{i_1} g_{i_2} g_{i_3} g_{i_4}] + f_3 [g_{i_2} g_{i_3} g_{i_1, i_4} + g_{i_3} g_{i_1} g_{i_2, i_4} + g_{i_1} g_{i_2} g_{i_3, i_4} + \\
 & \quad + g_{i_1} g_{i_4} g_{i_2, i_3} + g_{i_2} g_{i_4} g_{i_1, i_3} + g_{i_3} g_{i_4} g_{i_1, i_2}] + \\
 & \quad + f_2 [g_{i_1, i_2} g_{i_3, i_4} + g_{i_1, i_3} g_{i_2, i_4} + g_{i_1, i_4} g_{i_2, i_3}] + \\
 & \quad + f_1 [g_{i_1} g_{i_2, i_3, i_4} + g_{i_2} g_{i_1, i_3, i_4} + g_{i_3} g_{i_1, i_2, i_4} + g_{i_4} g_{i_1, i_2, i_3}] + \\
 & \quad + f_1 [g_{i_1, i_2, i_3, i_4}].
 \end{aligned}
 \tag{3.76}$$

Introducing the derivations

$$\begin{aligned}
 (3.77) \quad F_i^2 &:= \sum_{k_1=1}^n g_{k_1,i} \frac{\partial}{\partial g_{k_1}} + g_i \left(f_2 \frac{\partial}{\partial f_1} \right), \\
 F_i^3 &:= \sum_{k_1=1}^n g_{k_1,i} \frac{\partial}{\partial g_{k_1}} + \sum_{k_1,k_2=1}^n g_{k_1,k_2,i} \frac{\partial}{\partial g_{k_1,k_2}} + g_i \left(f_2 \frac{\partial}{\partial f_1} + f_3 \frac{\partial}{\partial f_2} \right), \\
 &\dots\dots\dots \\
 F_i^\lambda &:= \sum_{k_1=1}^n g_{k_1,i} \frac{\partial}{\partial g_{k_1}} + \sum_{k_1,k_2=1}^n g_{k_1,k_2,i} \frac{\partial}{\partial g_{k_1,k_2}} + \dots + \\
 &\quad + \sum_{k_1,\dots,k_{\lambda-1}=1}^n g_{k_1,\dots,k_{\lambda-1},i} \frac{\partial}{\partial g_{k_1,\dots,k_{\lambda-1}}} + \\
 &\quad + g_i \left(f_2 \frac{\partial}{\partial f_1} + f_3 \frac{\partial}{\partial f_2} + \dots + f_\lambda \frac{\partial}{\partial f_{\lambda-1}} \right),
 \end{aligned}$$

we observe that the following induction relations hold:

$$\begin{aligned}
 (3.78) \quad h_{i_1,i_2} &= F_{i_2}^2(h_{i_1}), \\
 h_{i_1,i_2,i_3} &= F_{i_3}^3(h_{i_1,i_2}), \\
 &\dots\dots\dots \\
 h_{i_1,i_2,\dots,i_\lambda} &= F_{i_\lambda}^\lambda(h_{i_1,i_2,\dots,i_{\lambda-1}}).
 \end{aligned}$$

To obtain the explicit version of the Faà di Bruno in the case of several variables (x^1, \dots, x^n) and one variable y , it suffices to extract from the expression of Y_{i_1,\dots,i_κ} provided by Theorem 3.73 only the terms corresponding to $\mu_1\lambda_1 + \dots + \mu_d\lambda_d = \kappa$, dropping all the \mathcal{X} terms. After some simplifications and after a translation by means of an elementary dictionary, we obtain a statement.

Theorem 3.79. *For every integer $\kappa \geq 1$ and for every choice of indices i_1, \dots, i_κ in the set $\{1, 2, \dots, n\}$, the κ -th partial derivative of the composite function $h = h(x^1, \dots, x^n) = f(g(x^1, \dots, x^n))$ with respect to the variables $x^{i_1}, \dots, x^{i_\kappa}$ may be expressed as an explicit polynomial depending on the derivatives of f , on the partial derivatives of g and having integer coefficients:*

$$(3.80) \quad \boxed{\frac{\partial^\kappa h}{\partial x^{i_1} \dots \partial x^{i_\kappa}} = \sum_{d=1}^\kappa \sum_{1 \leq \lambda_1 < \dots < \lambda_d \leq \kappa} \sum_{\mu_1 \geq 1, \dots, \mu_d \geq 1} \sum_{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d = \kappa} \frac{d^{\mu_1 + \dots + \mu_d} f}{dy^{\mu_1 + \dots + \mu_d}} \left[\sum_{\sigma \in \mathfrak{F}_\kappa^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)}} \prod_{1 \leq \nu_1 \leq \mu_1} \frac{\partial^{\lambda_1} g}{\partial x^{i_\sigma(1:\nu_1:1)} \dots \partial x^{i_\sigma(1:\nu_1:\lambda_1)}} \dots \dots \prod_{1 \leq \nu_d \leq \mu_d} \frac{\partial^{\lambda_d} g}{\partial x^{i_\sigma(d:\nu_d:1)} \dots \partial x^{i_\sigma(d:\nu_d:\lambda_d)}} \right]}.$$

In this formula, the coset $\mathfrak{F}_\kappa^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)}$ was defined in equation (3.71); we have made the identification:

$$(3.81) \quad \{1, \dots, \kappa\} \equiv \{1:1:1, \dots, 1:\mu_1:\lambda_1, \dots, d:1:1, \dots, d:\mu_d:\lambda_d\};$$

and also, for the sake of clarity, we have restituted the complete (not abbreviated) notation for the (partial) derivatives of f and of g .

We refer to Section 6 for the final writing of the above formula (3.80).

§4. ONE INDEPENDENT VARIABLE AND SEVERAL DEPENDENT VARIABLES

4.1. Simplified adapted notations. Assume $n = 1$ and $m \geq 1$, let $\kappa \in \mathbb{N}$ with $\kappa \geq 1$ and simply denote the jet variables by (instead of (1.2)):

$$(4.2) \quad (x, y^j, y_1^j, y_2^j, \dots, y_\kappa^j) \in \mathcal{J}_{1,m}^\kappa.$$

Instead of (1.30), denote the κ -th prolongation of a vector field by:

$$(4.3) \quad \left\{ \begin{array}{l} \mathcal{L}^{(\kappa)} = \mathcal{X} \frac{\partial}{\partial x} + \sum_{j=1}^m \mathcal{Y}^j \frac{\partial}{\partial y^j} + \sum_{j=1}^m \mathbf{Y}_1^j \frac{\partial}{\partial y_1^j} + \sum_{j=1}^m \mathbf{Y}_2^j \frac{\partial}{\partial y_2^j} + \\ \dots + \sum_{j=1}^m \mathbf{Y}_\kappa^j \frac{\partial}{\partial y_\kappa^j}. \end{array} \right.$$

The induction formulas are:

$$(4.4) \quad \left\{ \begin{array}{l} \mathbf{Y}_1^j := D^1(\mathcal{Y}^j) - D^1(\mathcal{X})y_1^j, \\ \mathbf{Y}_2^j := D^2(\mathbf{Y}_1^j) - D^1(\mathcal{X})y_2^j, \\ \dots \dots \dots \\ \mathbf{Y}_\lambda^j := D^\lambda(\mathbf{Y}_{\lambda-1}^j) - D^1(\mathcal{X})y_\lambda^j, \end{array} \right.$$

where the total differentiation operators D^λ are denoted by (instead of (1.22)):

$$(4.5) \quad D^\lambda := \frac{\partial}{\partial x} + \sum_{l=1}^m y_1^l \frac{\partial}{\partial y^l} + \sum_{l=1}^m y_2^l \frac{\partial}{\partial y_1^l} + \dots + \sum_{l=1}^m y_\lambda^l \frac{\partial}{\partial y_{\lambda-1}^l}.$$

Applying the definitions in the first two lines of (4.4), we compute, we simplify and we organize the results in a harmonious way, using in an essential way the Kronecker symbol. Here, the computations are more elementary than the computations of \mathbf{Y}_{i_1} and of \mathbf{Y}_{i_1, i_2} achieved thoroughly in the previous Section 3, so that we do not provide a Latex track of the details. Firstly and secondly:

$$(4.6) \quad \left\{ \begin{array}{l} \mathbf{Y}_1^j = \mathcal{Y}_x^j + \sum_{l_1=1}^m [\mathcal{Y}_{y^{l_1}}^j - \delta_{l_1}^j \mathcal{X}_x] y_1^{l_1} + \sum_{l_1, l_2=1}^m [-\delta_{l_1}^j \mathcal{X}_{y^{l_2}}] y_1^{l_1} y_1^{l_2}, \\ \mathbf{Y}_2^j = \mathcal{Y}_{x^2}^j + \sum_{l_1=1}^m [2\mathcal{Y}_{xy^{l_1}}^j - \delta_{l_1}^j \mathcal{X}_{x^2}] y_1^{l_1} + \sum_{l_1, l_2=1}^m [\mathcal{Y}_{y^{l_1} y^{l_2}}^j - \delta_{l_1}^j 2\mathcal{X}_{xy^{l_2}}] y_1^{l_1} y_1^{l_2} + \\ + \sum_{l_1, l_2, l_3} [-\delta_{l_1}^j \mathcal{X}_{y^{l_2} y^{l_3}}] y_1^{l_1} y_1^{l_2} y_1^{l_3} + \sum_{l_1} [\mathcal{Y}_{y^{l_1}}^j - \delta_{l_1}^j 2\mathcal{X}_x] y_2^{l_1} + \\ + \sum_{l_1, l_2=1}^m [-\delta_{l_1}^j \mathcal{X}_{y^{l_2}} - \delta_{l_2}^j 2\mathcal{X}_{y^{l_1}}] y_1^{l_1} y_2^{l_2}. \end{array} \right.$$

Thirdly:

$$(4.7) \quad \begin{aligned} \mathbf{Y}_3^j &= \mathcal{Y}_{x^3}^j + \sum_{l_1=1}^m [3\mathcal{Y}_{x^2 y^{l_1}}^j - \delta_{l_1}^j \mathcal{X}_{x^3}] y_1^{l_1} + \sum_{l_1, l_2=1}^m [3\mathcal{Y}_{xy^{l_1} y^{l_2}}^j - \delta_{l_1}^j 3\mathcal{X}_{x^2 y^{l_2}}] y_1^{l_1} y_1^{l_2} + \\ &+ \sum_{l_1, l_2, l_3} [\mathcal{Y}_{y^{l_1} y^{l_2} y^{l_3}}^j - \delta_{l_1}^j 3\mathcal{X}_{xy^{l_2} y^{l_3}}] y_1^{l_1} y_1^{l_2} y_1^{l_3} + \\ &+ \sum_{l_1, l_2, l_3, l_4} [-\delta_{l_1}^j \mathcal{X}_{y^{l_2} y^{l_3} y^{l_4}}] y_1^{l_1} y_1^{l_2} y_1^{l_3} y_1^{l_4} + \sum_{l_1=1}^m [3\mathcal{Y}_{xy^{l_1}}^j - \delta_{l_1}^j 3\mathcal{X}_{x^2}] y_2^{l_1} + \end{aligned}$$

$$\begin{aligned}
& + \sum_{l_1, l_2=1}^m \left[3\mathcal{Y}_{y^l_1 y^l_2}^j - \delta_{l_1}^j 3\mathcal{X}_{xy^l_2} - \delta_{l_2}^j 6\mathcal{X}_{xy^l_1} \right] y_1^{l_1} y_2^{l_2} + \\
& + \sum_{l_1, l_2, l_3=1}^m \left[-\delta_{l_1}^j 3\mathcal{X}_{y^l_2 y^l_3} - \delta_{l_3}^j 3\mathcal{X}_{y^l_1 y^l_2} \right] y_1^{l_1} y_1^{l_2} y_2^{l_3} + \sum_{l_1, l_2=1}^m \left[-\delta_{l_3}^j 3\mathcal{X}_{y^l_2} \right] y_2^{l_1} y_2^{l_2} + \\
& + \sum_{l_1=1}^m \left[\mathcal{Y}_{y^l_1}^j - \delta_{l_1}^j 3\mathcal{X}_x \right] y_3^{l_1} + \sum_{l_1, l_2=1}^m \left[-\delta_{l_1}^j \mathcal{X}_{y^l_2} - \delta_{l_2}^j 3\mathcal{X}_{y^l_1} \right] y_1^{l_1} y_3^{l_2}.
\end{aligned}$$

Fourthly:

$$\begin{aligned}
\mathbf{Y}_4^j & = \mathcal{Y}_{x^4}^j + \sum_{l_1=1}^m \left[4\mathcal{Y}_{x^3 y^l_1}^j - \delta_{l_1}^j \mathcal{X}_{x^4} \right] y_1^{l_1} + \sum_{l_1, l_2=1}^m \left[6\mathcal{Y}_{x^2 y^l_1 y^l_2}^j - \delta_{l_1}^j 4\mathcal{X}_{x^3 y^l_2} \right] y_1^{l_1} y_1^{l_2} + \\
& + \sum_{l_1, l_2, l_3=1}^m \left[4\mathcal{Y}_{xy^l_1 y^l_2 y^l_3}^j - \delta_{l_1}^j 6\mathcal{X}_{x^2 y^l_2 y^l_3} \right] y_1^{l_1} y_1^{l_2} y_1^{l_3} + \\
& + \sum_{l_1, l_2, l_3, l_4=1}^m \left[\mathcal{Y}_{xy^l_1 y^l_2 y^l_3 y^l_4}^j - \delta_{l_1}^j 4\mathcal{X}_{xy^l_2 y^l_3 y^l_4} \right] y_1^{l_1} y_1^{l_2} y_1^{l_3} y_1^{l_4} + \\
& + \sum_{l_1, l_2, l_3, l_4, l_5=1}^m \left[-\delta_{l_1}^j \mathcal{X}_{y^l_2 y^l_3 y^l_4 y^l_5} \right] y_1^{l_1} y_1^{l_2} y_1^{l_3} y_1^{l_4} y_1^{l_5} + \sum_{l_1=1}^m \left[6\mathcal{Y}_{x^2 y^l_1}^j - \delta_{l_1}^j 4\mathcal{X}_{x^3} \right] y_2^{l_1} + \\
& + \sum_{l_1, l_2=1}^m \left[12\mathcal{Y}_{xy^l_1 y^l_2}^j - \delta_{l_1}^j 6\mathcal{X}_{x^2 y^l_2} - \delta_{l_2}^j 12\mathcal{X}_{x^2 y^l_1} \right] y_1^{l_1} y_2^{l_2} + \\
(4.8) \quad & + \sum_{l_1, l_2, l_3=1}^m \left[6\mathcal{Y}_{y^l_1 y^l_2 y^l_3}^j - \delta_{l_1}^j 12\mathcal{X}_{xy^l_2 y^l_3} - \delta_{l_3}^j 12\mathcal{X}_{xy^l_1 y^l_2} \right] y_1^{l_1} y_1^{l_2} y_2^{l_3} + \\
& + \sum_{l_1, l_2, l_3, l_4=1}^m \left[-\delta_{l_1}^j 6\mathcal{X}_{y^l_2 y^l_3 y^l_4} - \delta_{l_4}^j 4\mathcal{X}_{y^l_1 y^l_2 y^l_3} \right] y_1^{l_1} y_1^{l_2} y_1^{l_3} y_2^{l_4} + \\
& + \sum_{l_1, l_2=1}^m \left[3\mathcal{Y}_{y^l_1 y^l_2}^j - \delta_{l_1}^j 12\mathcal{X}_{xy^l_2} \right] y_2^{l_1} y_2^{l_2} + \\
& + \sum_{l_1, l_2, l_3=1}^m \left[-\delta_{l_1}^j 3\mathcal{X}_{y^l_2 y^l_3} - \delta_{l_2}^j 12\mathcal{X}_{y^l_1 y^l_3} \right] y_1^{l_1} y_2^{l_2} y_2^{l_3} + \sum_{l_1=1}^m \left[4\mathcal{Y}_{xy^l_1}^j - \delta_{l_1}^j 6\mathcal{X}_{x^2} \right] y_3^{l_1} + \\
& + \sum_{l_1, l_2=1}^m \left[4\mathcal{Y}_{y^l_1 y^l_2}^j - \delta_{l_1}^j 4\mathcal{X}_{xy^l_2} - \delta_{l_2}^j 12\mathcal{X}_{xy^l_1} \right] y_1^{l_1} y_3^{l_2} + \\
& + \sum_{l_1, l_2, l_3=1}^m \left[-\delta_{l_1}^j 4\mathcal{X}_{y^l_2 y^l_3} - \delta_{l_3}^j 6\mathcal{X}_{y^l_1 y^l_2} \right] y_1^{l_1} y_1^{l_2} y_3^{l_3} + \\
& + \sum_{l_1, l_2=1}^m \left[-\delta_{l_1}^j 4\mathcal{X}_{y^l_2} - \delta_{l_2}^j 6\mathcal{X}_{y^l_1} \right] y_2^{l_1} y_3^{l_2} + \\
& + \sum_{l_1=1}^m \left[\mathcal{Y}_{y^l_1}^j - \delta_{l_1}^j 4\mathcal{X}_x \right] y_4^{l_1} + \sum_{l_1, l_2=1}^m \left[-\delta_{l_1}^j \mathcal{X}_{y^l_2} - \delta_{l_2}^j 4\mathcal{X}_{y^l_1} \right] y_1^{l_1} y_4^{l_2}.
\end{aligned}$$

4.9. Inductive elaboration of the general formula. Now we compare the formula (2.9) for \mathbf{Y}_4 with the above formula (4.8) for \mathbf{Y}_4^j . The goal is to find the rules of transformation

and of development by inspecting several instances, in order to devise how to transform and to develop the formula (2.25) to several dependent variables.

First of all, we have to develop the general monomial $(y_{\lambda_1})^{\mu_1} \cdots (y_{\lambda_d})^{\mu_d}$. In every monomial present in the expressions of \mathbf{Y}_1^j , of \mathbf{Y}_2^j , of \mathbf{Y}_3^j and of \mathbf{Y}_4^j above, we see that the number α of indices l_β appearing in all the sums $\sum_{l_1, \dots, l_\alpha=1}^m$ is exactly equal to $\mu_1 + \cdots + \mu_d$. To denote these $\mu_1 + \cdots + \mu_d$ indices l_β , we shall use the notation:

$$(4.10) \quad \underbrace{\underbrace{l_{1:1}, \dots, l_{1:\mu_1}}_{\mu_1}, \dots, \underbrace{l_{d:1}, \dots, l_{d:\mu_d}}_{\mu_d}}_{\mu_1 + \cdots + \mu_d}$$

inspired by Convention 3.33. With such a choice of notation, we may avoid long subscripts in the indices l_β , like $l_{\mu_1 + \cdots + \mu_d}$. It follows that the development of the general monomial $(y_{\lambda_1})^{\mu_1} \cdots (y_{\lambda_d})^{\mu_d}$ to several dependent variables yields $m^{\mu_1 + \cdots + \mu_d}$ possible choices:

$$(4.11) \quad \prod_{1 \leq \nu_1 \leq \mu_1} y_{\lambda_1}^{l_{1:\nu_1}} \cdots \prod_{1 \leq \nu_d \leq \mu_d} y_{\lambda_d}^{l_{d:\nu_d}},$$

where the indices $l_{1:1}, \dots, l_{1:\mu_1}, \dots, l_{d:1}, \dots, l_{d:\mu_d}$ take their values in the set $\{1, 2, \dots, m\}$. Consequently, the general expression of \mathbf{Y}_κ^j must be of the form:

$$(4.12) \quad \mathbf{Y}_\kappa^j = \mathcal{Y}_{x^\kappa}^j + \sum_{d=1}^{\kappa+1} \sum_{1 \leq \lambda_1 < \cdots < \lambda_d \leq \kappa} \sum_{\mu_1 \geq 1, \dots, \mu_d \geq 1} \sum_{\mu_1 \lambda_1 + \cdots + \mu_d \lambda_d \leq \kappa+1} \sum_{l_{1:1}=1}^m \cdots \sum_{l_{1:\mu_1}=1}^m \cdots \sum_{l_{d:1}=1}^m \cdots \sum_{l_{d:\mu_d}=1}^m \prod_{1 \leq \nu_1 \leq \mu_1} y_{\lambda_1}^{l_{1:\nu_1}} \cdots \prod_{1 \leq \nu_d \leq \mu_d} y_{\lambda_d}^{l_{d:\nu_d}}, \quad [?]$$

where the term in brackets [?] is still unknown. To determine it, let us examine a few instances.

According to (4.8) (fourth line), the term $[6 \mathcal{Y}_{x^2 y} - 4 \mathcal{X}_{x^3}] y_2$ of \mathbf{Y}_4 develops as $\sum_{l_1=1}^m [6 \mathcal{Y}_{x^2 y^{l_1}} - \delta_{l_1}^j 4 \mathcal{X}_{x^3}] y_2^{l_1}$ in \mathbf{Y}_4^j . Here, $6 \mathcal{Y}_{x^2 y}$ just becomes $6 \mathcal{Y}_{x^2 y^{l_1}}$. Thus, we suspect that the term $\frac{\kappa \cdots (\kappa - \mu_1 \lambda_1 - \cdots - \mu_d \lambda_d + 1)}{(\lambda_1!)^{\mu_1} \mu_1! \cdots (\lambda_d!)^{\mu_d} \mu_d!} \cdot \mathcal{Y}_{x^{\kappa - \mu_1 \lambda_1 - \cdots - \mu_d \lambda_d} y^{\mu_1 + \cdots + \mu_d}}$ of the second line of (2.25) should simply be developed as

$$(4.13) \quad \frac{\kappa(\kappa - 1) \cdots (\kappa - \mu_1 \lambda_1 - \cdots - \mu_d \lambda_d + 1)}{(\lambda_1!)^{\mu_1} \mu_1! \cdots (\lambda_d!)^{\mu_d} \mu_d!} \cdot \frac{\partial^{\kappa - \mu_1 \lambda_1 - \cdots - \mu_d \lambda_d + \mu_1 + \cdots + \mu_d} \mathcal{Y}^j}{(\partial x)^{\kappa - \mu_1 \lambda_1 - \cdots - \mu_d \lambda_d} \partial y^{l_{1:1}} \cdots \partial y^{l_{1:\mu_1}} \cdots \partial y^{l_{d:1}} \cdots \partial y^{l_{d:\mu_d}}}.$$

This rule is confirmed by inspecting all the other monomials of \mathbf{Y}_1^j , of \mathbf{Y}_2^j , of \mathbf{Y}_3^j and of \mathbf{Y}_4^j .

It remains to determine how we must develop the term in \mathcal{X} appearing in the last two lines of (2.25). To begin with, let us rewrite in advance this term in the slightly different shape, emphasizing a factorization:

$$(4.14) \quad \frac{\kappa \cdots (\kappa - \mu_1 \lambda_1 - \cdots - \mu_d \lambda_d + 2)}{(\lambda_1!)^{\mu_1} \mu_1! \cdots (\lambda_d!)^{\mu_d} \mu_d!} \left[(\mu_1 \lambda_1 + \cdots + \mu_d \lambda_d) \mathcal{X}_{x^{\kappa - \mu_1 \lambda_1 - \cdots - \mu_d \lambda_d + 1} y^{\mu_1 + \cdots + \mu_d - 1}} \right].$$

Then we examine four instances extracted from the complete expression of \mathbf{Y}_4^j :

$$(4.15) \quad \left\{ \begin{array}{l} \sum_{l_1, l_2, l_3=1}^m \left[4 \mathcal{Y}_{xy^{l_1}y^{l_2}y^{l_3}}^j - \delta_{l_1}^j 6 \mathcal{X}_{x^2y^{l_2}y^{l_3}} \right] y_1^{l_1} y_1^{l_2} y_1^{l_3}, \\ \sum_{l_1, l_2=1}^m \left[12 \mathcal{Y}_{xy^{l_1}y^{l_2}}^j - \delta_{l_1}^j 6 \mathcal{X}_{x^2y^{l_2}} - \delta_{l_2}^j 12 \mathcal{X}_{x^2y^{l_1}} \right] y_1^{l_1} y_2^{l_2}, \\ \sum_{l_1, l_2, l_3, l_4=1}^m \left[-\delta_{l_1}^j 6 \mathcal{X}_{y^{l_2}y^{l_3}y^{l_4}} - \delta_{l_4}^j 4 \mathcal{X}_{y^{l_1}y^{l_2}y^{l_3}} \right] y_1^{l_1} y_1^{l_2} y_1^{l_3} y_2^{l_4}, \\ \sum_{l_1, l_2, l_3=1}^m \left[-\delta_{l_1}^j 4 \mathcal{X}_{y^{l_2}y^{l_3}} - \delta_{l_3}^j 6 \mathcal{X}_{y^{l_1}y^{l_2}} \right] y_1^{l_1} y_1^{l_2} y_3^{l_3}, \end{array} \right.$$

and we compare them to the corresponding terms of \mathbf{Y}_4 :

$$(4.16) \quad \left\{ \begin{array}{l} [4 \mathcal{Y}_{xy^3} - 6 \mathcal{X}_{x^2y^2}] (y_1)^3, \\ [12 \mathcal{Y}_{xy^2} - 18 \mathcal{X}_{x^2y}] y_1 y_2, \\ [-10 \mathcal{X}_{y^3}] (y_1)^3 y_2, \\ [-10 \mathcal{X}_{y^2}] (y_1)^2 y_3. \end{array} \right.$$

In the development from (4.16) to (4.15), we see that the four integers just before \mathcal{X} , namely $6 = 6$, $18 = 6 + 12$, $10 = 6 + 4$ and $10 = 4 + 6$, are split in a certain manner. Also, a single Kronecker symbol $\delta_{l_\alpha}^j$ is added as a factor. *What are the rules?*

In the second splitting $18 = 6 + 12$, we see that the relative weight of 6 and of 12 is the same as the relative weight of 1 and 2 in the splitting $3 = 1 + 2$ issued from the lower indices of the corresponding monomial $y_1^{l_1} y_2^{l_2}$. Similarly, in the third splitting $10 = 6 + 4$, the relative weight of 6 and of 4 is the same as the relative weight of $1 + 1 + 1$ and of 2 issued from the lower indices of the corresponding monomial $y_1^{l_1} y_1^{l_2} y_1^{l_3} y_2^{l_4}$. This rule may be confirmed by inspecting all the other monomials of \mathbf{Y}_2 , \mathbf{Y}_2^j , of \mathbf{Y}_3 , \mathbf{Y}_3^j and of \mathbf{Y}_4 , \mathbf{Y}_4^j . For a general $\kappa \geq 1$, the splitting of integers just amounts to decompose the sum appearing inside the brackets of (4.14) as $\mu_1 \lambda_1, \mu_2 \lambda_2, \dots, \mu_d \lambda_d$. In fact, when we wrote (4.14), we emphasized in advance the decomposable factor $(\mu_1 \lambda_1 + \dots + \mu_d \lambda_d)$.

Next, we have to determine what is the subscript α in the Kronecker symbol $\delta_{l_\alpha}^j$. We claim that in the four instances (4.15), the subscript α is intrinsically related to weight splitting. Indeed, recall that in the second line of (4.15), the number 6 of the splitting $18 = 6 + 12$ is related to the number 1 in the splitting $3 = 1 + 2$ of the lower indices of the monomial $y_1^{l_1} y_2^{l_2}$. It follows that the index l_α *must be* the index l_1 of the monomial $y_1^{l_1}$. Similarly, also in the second line of (4.15), the number 12 of the splitting $18 = 6 + 12$ being related to the number 2 in the splitting $3 = 1 + 2$ of the lower indices of the monomial $y_1^{l_1} y_2^{l_2}$, it follows that the index l_α attached to the second \mathcal{X} term must be the index l_2 of the monomial $y_2^{l_2}$.

This rule is still ambiguous. Indeed, let us examine the third line of (4.15). We have the splitting $10 = 6 + 4$, homologous to the splitting of relative weights $5 = (1 + 1 + 1) + 2$ in the monomial $y_1^{l_1} y_1^{l_2} y_1^{l_3} y_2^{l_4}$. Of course, it is clear that we must choose the index l_4 for the Kronecker symbol associated to the second \mathcal{X} term $-4 \mathcal{X}_{y^3}$, thus obtaining $-\delta_{l_4}^j 4 \mathcal{X}_{y^{l_1}y^{l_2}y^{l_3}}$. However, since the monomial $y_1^{l_1} y_1^{l_2} y_1^{l_3}$ has three indices l_1, l_2 and l_3 , there arises a question: *what index l_α must we choose for the Kronecker symbol $\delta_{l_\alpha}^j$ attached to the first \mathcal{X} term $6 \mathcal{X}_{y^3}$: the index l_1 , the index l_2 or the index l_3 ?*

The answer is simple: *any of the three indices l_1, l_2 or l_3 works*. Indeed, since the monomial $y_1^{l_1} y_1^{l_2} y_1^{l_3}$ is symmetric with respect to all permutations of the set of three indices $\{l_1, l_2, l_3\}$, we have

(4.17)

$$\begin{aligned} \sum_{l_1, l_2, l_3, l_4=1}^m \left[-\delta_{l_1}^j 6 \mathcal{X}_{y^{l_2} y^{l_3} y^{l_4}} \right] y_1^{l_1} y_1^{l_2} y_1^{l_3} y_2^{l_4} &= \sum_{l_1, l_2, l_3, l_4=1}^m \left[-\delta_{l_2}^j 6 \mathcal{X}_{y^{l_1} y^{l_3} y^{l_4}} \right] y_1^{l_1} y_1^{l_2} y_1^{l_3} y_2^{l_4} = \\ &= \sum_{l_1, l_2, l_3, l_4=1}^m \left[-\delta_{l_3}^j 6 \mathcal{X}_{y^{l_1} y^{l_2} y^{l_4}} \right] y_1^{l_1} y_1^{l_2} y_1^{l_3} y_2^{l_4}. \end{aligned}$$

In fact, we have systematically used such symmetries during the intermediate computations (not exposed here) which we achieved manually to obtain the final expressions of \mathbf{Y}_1^j , of \mathbf{Y}_2^j , of \mathbf{Y}_3^j and of \mathbf{Y}_4^j . To fix ideas, we have always chosen the first index. Here, the first index is l_1 ; in the first sum of line 9 of (4.8), the first index l_α for the second weight 12 is l_2 .

This rule may be confirmed by inspecting all the monomials of \mathbf{Y}_2^j , of \mathbf{Y}_3^j , of \mathbf{Y}_4^j (and also of \mathbf{Y}_5^j , which we have computed in a manuscript, but not copied in this Latex file).

From these considerations, we deduce that for the general formula, the weight decomposition is simply $\mu_1 \lambda_1, \dots, \mu_d \lambda_d$ and that the Kronecker symbol δ_α^j is intrinsically associated to the weights. In conclusion, building on inductive reasonings, we have formulated the following statement.

Theorem 4.18. *For one independent variable x , for several dependent variables (y^1, \dots, y^m) and for $\kappa \geq 1$, the general expression of the coefficient \mathbf{Y}_κ^j of the prolongation (4.3) of a vector field is:*

(4.19)

$$\begin{aligned} \mathbf{Y}_\kappa^j &= \mathcal{Y}_{x^\kappa}^j + \sum_{d=1}^{\kappa+1} \sum_{1 \leq \lambda_1 < \dots < \lambda_d \leq \kappa} \sum_{\mu_1 \geq 1, \dots, \mu_d \geq 1} \sum_{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d \leq \kappa+1} \\ &\quad \sum_{l_{1:1}=1}^m \dots \sum_{l_{1:\mu_1}=1}^m \dots \sum_{l_{d:1}=1}^m \dots \sum_{l_{d:\mu_d}=1}^m \frac{\kappa(\kappa-1) \dots (\kappa - \mu_1 \lambda_1 + \dots + \mu_d \lambda_d + 2)}{(\lambda_1!)^{\mu_1} \mu_1! \dots (\lambda_d!)^{\mu_d} \mu_d!} \\ &\quad \left[\begin{aligned} &(\kappa - \mu_1 \lambda_1 - \dots - \mu_d \lambda_d + 1) \frac{\partial^{\kappa - \mu_1 \lambda_1 - \dots - \mu_d \lambda_d + \mu_1 + \dots + \mu_d} \mathcal{Y}^j}{(\partial x)^{\kappa - \mu_1 \lambda_1 - \dots - \mu_d \lambda_d} \partial y^{l_{1:1}} \dots \partial y^{l_{1:\mu_1}} \dots \partial y^{l_{d:1}} \dots \partial y^{l_{d:\mu_d}}} \\ &- \delta_{l_{1:1}}^j \mu_1 \lambda_1 \frac{\partial^{\kappa - \mu_1 \lambda_1 - \dots - \mu_d \lambda_d + \mu_1 + \dots + \mu_d} \mathcal{X}}{(\partial x)^{\kappa - \mu_1 \lambda_1 - \dots - \mu_d \lambda_d + 1} \widehat{\partial y^{l_{1:1}}} \dots \partial y^{l_{1:\mu_1}} \dots \partial y^{l_{d:1}} \dots \partial y^{l_{d:\mu_d}}} \\ &\dots \dots \dots \\ &- \delta_{l_{d:1}}^j \mu_d \lambda_d \frac{\partial^{\kappa - \mu_1 \lambda_1 - \dots - \mu_d \lambda_d + \mu_1 + \dots + \mu_d} \mathcal{X}}{(\partial x)^{\kappa - \mu_1 \lambda_1 - \dots - \mu_d \lambda_d + 1} \partial y^{l_{1:1}} \dots \partial y^{l_{1:\mu_1}} \dots \widehat{\partial y^{l_{d:1}}} \dots \partial y^{l_{d:\mu_d}}} \end{aligned} \right] \\ &\quad \cdot \prod_{1 \leq \nu_1 \leq \mu_1} y_{\lambda_1}^{l_{1:\nu_1}} \dots \prod_{1 \leq \nu_d \leq \mu_d} y_{\lambda_d}^{l_{d:\nu_d}}. \end{aligned}$$

Here, the notation $\widehat{\partial y^l}$ means that the partial derivative is dropped.

Since the fundamental monomials appearing in the last line of (4.19) just above are not independent of each other, this formula has still to be modified a little bit. We refer to Section 6 for details.

4.20. Deduction of a multivariate Faà di Bruno formula. Let $m \in \mathbb{N}$ with $m \geq 1$, let $y = (y^1, \dots, y^m) \in \mathbb{K}^m$, let $f = f(y^1, \dots, y^m)$ be a C^∞ -smooth function from \mathbb{K}^m to \mathbb{K} ,

let $x \in \mathbb{K}$ and let $g^1 = g^1(x), \dots, g^m = g^m(x)$ be C^∞ functions from \mathbb{K} to \mathbb{K} . The goal is to obtain an explicit formula for the derivatives, with respect to x , of the composition $h := f \circ g$, namely $h(x) := f(g^1(x), \dots, g^m(x))$. For $\lambda \in \mathbb{N}$ with $\lambda \geq 1$, and for $j = 1, \dots, m$, we shall abbreviate the derivative $\frac{d^\lambda g^j}{dx^\lambda}$ by g_λ^j and similarly for h_λ . The partial derivatives $\frac{\partial^\lambda f}{\partial y^{l_1} \dots \partial y^{l_\lambda}}$ will be abbreviated by $f_{l_1, \dots, l_\lambda}$.

Applying the chain rule, we may compute:

(4.21)

$$\begin{aligned}
 h_1 &= \sum_{l_1=1}^m f_{l_1} g_1^{l_1}, \\
 h_2 &= \sum_{l_1, l_2=1}^m f_{l_1, l_2} g_1^{l_1} g_1^{l_2} + \sum_{l_1=1}^m f_{l_1} g_2^{l_1}, \\
 h_3 &= \sum_{l_1, l_2, l_3=1}^m f_{l_1, l_2, l_3} g_1^{l_1} g_1^{l_2} g_1^{l_3} + 3 \sum_{l_1, l_2=1}^m f_{l_1, l_2} g_1^{l_1} g_2^{l_2} + \sum_{l_1=1}^m f_{l_1} g_3^{l_1}, \\
 h_4 &= \sum_{l_1, l_2, l_3, l_4=1}^m f_{l_1, l_2, l_3, l_4} g_1^{l_1} g_1^{l_2} g_1^{l_3} g_1^{l_4} + 6 \sum_{l_1, l_2, l_3=1}^m f_{l_1, l_2, l_3} g_1^{l_1} g_1^{l_2} g_2^{l_3} + \\
 &\quad + 3 \sum_{l_1, l_2=1}^m f_{l_1, l_2} g_2^{l_1} g_2^{l_2} + 4 \sum_{l_1, l_2=1}^m f_{l_1, l_2} g_1^{l_1} g_3^{l_2} + \sum_{l_1=1}^m f_{l_1} g_4^{l_1}, \\
 h_5 &= \sum_{l_1, l_2, l_3, l_4, l_5=1}^m f_{l_1, l_2, l_3, l_4, l_5} g_1^{l_1} g_1^{l_2} g_1^{l_3} g_1^{l_4} g_1^{l_5} + 10 \sum_{l_1, l_2, l_3, l_4=1}^m f_{l_1, l_2, l_3, l_4} g_1^{l_1} g_1^{l_2} g_1^{l_3} g_2^{l_4} + \\
 &\quad + 15 \sum_{l_1, l_2, l_3=1}^m f_{l_1, l_2, l_3} g_1^{l_1} g_2^{l_2} g_2^{l_3} + 10 \sum_{l_1, l_2, l_3=1}^m f_{l_1, l_2, l_3} g_1^{l_1} g_1^{l_2} g_3^{l_3} + \\
 &\quad + 10 \sum_{l_1, l_2=1}^m f_{l_1, l_2} g_2^{l_1} g_3^{l_2} + 5 \sum_{l_1, l_2=1}^m f_{l_1, l_2} g_1^{l_1} g_4^{l_2} + \sum_{l_1=1}^m f_{l_1} g_5^{l_1}.
 \end{aligned}$$

Introducing the derivations

(4.22)

$$\begin{aligned}
 F^2 &:= \sum_{l_1=1}^m g_2^{l_1} \frac{\partial}{\partial g_1^{l_1}} + \sum_{l_1=1}^m g_1^{l_1} \left(\sum_{l_2=1}^m f_{l_1, l_2} \frac{\partial}{\partial f_{l_2}} \right), \\
 F^3 &:= \sum_{l_1=1}^m g_2^{l_1} \frac{\partial}{\partial g_1^{l_1}} + \sum_{l_1=1}^m g_3^{l_1} \frac{\partial}{\partial g_2^{l_1}} + \sum_{l_1=1}^m g_1^{l_1} \left(\sum_{l_2=1}^m f_{l_1, l_2} \frac{\partial}{\partial f_{l_2}} + \sum_{l_2, l_3=1}^m f_{l_1, l_2, l_3} \frac{\partial}{\partial f_{l_2, l_3}} \right), \\
 &\dots\dots\dots \\
 F^\lambda &:= \sum_{l_1=1}^m g_2^{l_1} \frac{\partial}{\partial g_1^{l_1}} + \sum_{l_1=1}^m g_3^{l_1} \frac{\partial}{\partial g_2^{l_1}} + \dots + \sum_{l_1=1}^m g_\lambda^{l_1} \frac{\partial}{\partial g_{\lambda-1}^{l_1}} + \\
 &\quad + \sum_{l_1=1}^m g_1^{l_1} \left(\sum_{l_2=1}^m f_{l_1, l_2} \frac{\partial}{\partial f_{l_2}} + \sum_{l_2, l_3=1}^m f_{l_1, l_2, l_3} \frac{\partial}{\partial f_{l_2, l_3}} + \dots + \sum_{l_2, \dots, l_\lambda=1}^m f_{l_1, l_2, \dots, l_\lambda} \frac{\partial}{\partial f_{l_2, \dots, l_\lambda}} \right),
 \end{aligned}$$

we observe that the following induction relations hold:

$$\begin{aligned}
 (4.23) \quad & h_2 = F^2(h_1), \\
 & h_3 = F^3(h_2), \\
 & \dots\dots\dots \\
 & h_\lambda = F^\lambda(h_{\lambda-1}).
 \end{aligned}$$

To obtain the explicit version of the Faà di Bruno in the case of one variable x and several variables (y^1, \dots, y^m) , it suffices to extract from the expression of \mathbf{Y}_κ^j provided by Theorem 4.18 only the terms corresponding to $\mu_1\lambda_1 + \dots + \mu_d\lambda_d = \kappa$, dropping all the \mathcal{X} terms. After some simplifications and after a translation by means of an elementary dictionary, we may formulate a statement.

Theorem 4.24. *For every integer $\kappa \geq 1$, the κ -th partial derivative of the composite function $h = h(x) = f(g^1(x), \dots, g^m(x))$ with respect to x may be expressed as an explicit polynomial depending on the partial derivatives of f , on the derivatives of g and having integer coefficients:*

$$(4.25) \quad \frac{d^\kappa h}{dx^\kappa} = \sum_{d=1}^{\kappa} \sum_{1 \leq \lambda_1 < \dots < \lambda_d \leq \kappa} \sum_{\mu_1 \geq 1, \dots, \mu_d \geq 1} \sum_{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d = \kappa} \frac{\kappa!}{(\lambda_1!)^{\mu_1} \mu_1! \dots (\lambda_d!)^{\mu_d} \mu_d!}$$

$$\sum_{l_{1:1}, \dots, l_{1:\mu_1}=1}^m \dots \sum_{l_{d:1}, \dots, l_{d:\mu_d}=1}^m$$

$$\frac{\partial^{\mu_1 + \dots + \mu_d} f}{\partial y^{l_{1:1}} \dots \partial y^{l_{1:\mu_1}} \dots \partial y^{l_{d:1}} \dots \partial y^{l_{d:\mu_d}}} \prod_{1 \leq \nu_1 \leq \mu_1} \frac{d^{\lambda_1} g^{l_{1:\nu_1}}}{dx^{\lambda_1}} \dots \prod_{1 \leq \nu_d \leq \mu_d} \frac{d^{\lambda_d} g^{l_{d:\nu_d}}}{dx^{\lambda_d}}.$$

We refer to Section 6 for the final writing of the above formula (4.25).

§5. SEVERAL INDEPENDENT VARIABLES AND SEVERAL DEPENDENT VARIABLES

5.1. Expression of $\mathbf{Y}_{i_1}^j$, of \mathbf{Y}_{i_1, i_2}^j and of $\mathbf{Y}_{i_1, i_2, i_3}^j$. Applying the induction (1.31) and working out the obtained formulas until they take a perfect shape, we obtain firstly:

$$(5.2) \quad \mathbf{Y}_{i_1}^j = \mathcal{Y}_{x^{i_1}}^j + \sum_{l_1=1}^m \sum_{k_1=1}^n \left[\delta_{i_1}^{k_1} \mathcal{Y}_{y^{l_1}}^j - \delta_{l_1}^j \mathcal{X}_{x^{i_1}}^{k_1} \right] y_{k_1}^{l_1} + \sum_{l_1, l_2=1}^m \sum_{k_1, k_2=1}^n \left[-\delta_{l_2}^j \delta_{i_1}^{k_1} \mathcal{X}_{y^{l_1}}^{k_2} \right] y_{k_1}^{l_1} y_{k_2}^{l_2}.$$

Secondly:

$$(5.3) \quad \begin{aligned} \mathbf{Y}_{i_1, i_2}^j &= \mathcal{Y}_{x^{i_1} x^{i_2}}^j + \sum_{l_1=1}^m \sum_{k_1=1}^n \left[\delta_{i_1}^{k_1} \mathcal{Y}_{x^{i_2} y^{l_1}}^j + \delta_{i_2}^{k_1} \mathcal{Y}_{x^{i_1} y^{l_1}}^j - \delta_{l_1}^j \mathcal{X}_{x^{i_1} x^{i_2}}^{k_1} \right] y_{k_1}^{l_1} + \\ &+ \sum_{l_1, l_2=1}^m \sum_{k_1, k_2=1}^n \left[\delta_{i_1, i_2}^{k_1, k_2} \mathcal{Y}_{y^{l_1} y^{l_2}}^j - \delta_{l_2}^j \delta_{i_1}^{k_1} \mathcal{X}_{x^{i_2} y^{l_1}}^{k_2} - \delta_{l_2}^j \delta_{i_2}^{k_1} \mathcal{X}_{x^{i_1} y^{l_1}}^{k_2} \right] y_{k_1}^{l_1} y_{k_2}^{l_2} + \\ &+ \sum_{l_1, l_2, l_3=1}^m \sum_{k_1, k_2, k_3=1}^n \left[-\delta_{l_3}^j \delta_{i_1, i_2}^{k_1, k_2} \mathcal{X}_{y^{l_1} y^{l_2}}^{k_3} \right] y_{k_1}^{l_1} y_{k_2}^{l_2} y_{k_3}^{l_3} + \\ &+ \sum_{l_1=1}^m \sum_{k_1, k_2=1}^n \left[\delta_{i_1, i_2}^{k_1, k_2} \mathcal{Y}_{y^{l_1}}^j - \delta_{l_1}^j \delta_{i_1}^{k_1} \mathcal{X}_{x^{i_2}}^{k_2} - \delta_{l_1}^j \delta_{i_2}^{k_1} \mathcal{X}_{x^{i_1}}^{k_2} \right] y_{k_1, k_2}^{l_1} + \\ &+ \sum_{l_1, l_2=1}^m \sum_{k_1, k_2, k_3=1}^n \left[-\delta_{l_1}^j \delta_{i_1, i_2}^{k_2, k_3} \mathcal{X}_{y^{l_2}}^{k_1} - \delta_{l_2}^j \delta_{i_1, i_2}^{k_3, k_1} \mathcal{X}_{y^{l_1}}^{k_2} - \delta_{l_2}^j \delta_{i_1, i_2}^{k_1, k_2} \mathcal{X}_{y^{l_1}}^{k_3} \right] y_{k_1}^{l_1} y_{k_2}^{l_2} y_{k_3}^{l_3}. \end{aligned}$$

Thirdly:

$$(5.4) \quad \begin{aligned} \mathbf{Y}_{i_1, i_2, i_3}^j &= \mathcal{Y}_{x^{i_1} x^{i_2} x^{i_3}}^j + \sum_{l_1=1}^m \sum_{k_1=1}^n \left[\delta_{i_1}^{k_1} \mathcal{Y}_{x^{i_2} x^{i_3} y^{l_1}}^j + \delta_{i_2}^{k_1} \mathcal{Y}_{x^{i_1} x^{i_3} y^{l_1}}^j + \delta_{i_3}^{k_1} \mathcal{Y}_{x^{i_1} x^{i_2} y^{l_1}}^j - \delta_{l_1}^j \mathcal{X}_{x^{i_1} x^{i_2} x^{i_3}}^{k_1} \right] y_{k_1}^{l_1} + \\ &+ \sum_{l_1, l_2=1}^m \sum_{k_1, k_2=1}^n \left[\delta_{i_1, i_2}^{k_1, k_2} \mathcal{Y}_{x^{i_3} y^{l_1} y^{l_2}}^j + \delta_{i_3, i_1}^{k_1, k_2} \mathcal{Y}_{x^{i_2} y^{l_1} y^{l_2}}^j + \delta_{i_2, i_3}^{k_1, k_2} \mathcal{Y}_{x^{i_1} y^{l_1} y^{l_2}}^j - \right. \\ &\quad \left. - \delta_{l_2}^j \delta_{i_1}^{k_1} \mathcal{X}_{x^{i_2} x^{i_3} y^{l_1}}^{k_2} - \delta_{l_2}^j \delta_{i_2}^{k_1} \mathcal{X}_{x^{i_1} x^{i_3} y^{l_1}}^{k_2} - \delta_{l_2}^j \delta_{i_3}^{k_1} \mathcal{X}_{x^{i_1} x^{i_2} y^{l_1}}^{k_2} \right] y_{k_1}^{l_1} y_{k_2}^{l_2} + \\ &+ \sum_{l_1, l_2, l_3=1}^m \sum_{k_1, k_2, k_3=1}^n \left[\delta_{i_1, i_2, i_3}^{k_1, k_2, k_3} \mathcal{Y}_{y^{l_1} y^{l_2} y^{l_3}}^j - \delta_{l_3}^j \delta_{i_1, i_2}^{k_1, k_2} \mathcal{X}_{x^{i_3} y^{l_1} y^{l_2}}^{k_3} - \right. \\ &\quad \left. - \delta_{l_3}^j \delta_{i_1, i_3}^{k_1, k_2} \mathcal{X}_{x^{i_2} y^{l_1} y^{l_2}}^{k_3} - \delta_{l_3}^j \delta_{i_2, i_3}^{k_1, k_2} \mathcal{X}_{x^{i_1} y^{l_1} y^{l_2}}^{k_3} \right] y_{k_1}^{l_1} y_{k_2}^{l_2} y_{k_3}^{l_3} + \\ &+ \sum_{l_1, l_2, l_3, l_4=1}^m \sum_{k_1, k_2, k_3, k_4=1}^n \left[-\delta_{l_4}^j \delta_{i_1, i_2, i_3}^{k_1, k_2, k_3} \mathcal{X}_{y^{l_1} y^{l_2} y^{l_3}}^{k_4} \right] y_{k_1}^{l_1} y_{k_2}^{l_2} y_{k_3}^{l_3} y_{k_4}^{l_4} + \\ &+ \sum_{l_1=1}^m \sum_{k_1, k_2=1}^n \left[\delta_{i_1, i_2}^{k_1, k_2} \mathcal{Y}_{x^{i_3} y^{l_1}}^j + \delta_{i_3, i_1}^{k_1, k_2} \mathcal{Y}_{x^{i_2} y^{l_1}}^j + \delta_{i_2, i_3}^{k_1, k_2} \mathcal{Y}_{x^{i_1} y^{l_1}}^j - \right. \\ &\quad \left. - \delta_{l_1}^j \delta_{i_1}^{k_1} \mathcal{X}_{x^{i_2} x^{i_3}}^{k_2} - \delta_{l_1}^j \delta_{i_2}^{k_1} \mathcal{X}_{x^{i_1} x^{i_3}}^{k_2} - \delta_{l_1}^j \delta_{i_3}^{k_1} \mathcal{X}_{x^{i_1} x^{i_2}}^{k_2} \right] y_{k_1, k_2}^{l_1} + \end{aligned}$$

$$(5.4) \quad \begin{aligned} &+ \sum_{l_1, l_2=1}^m \sum_{k_1, k_2, k_3=1}^n \left[\delta_{i_1, i_2, i_3}^{k_1, k_2, k_3} \mathcal{Y}_{y^{l_1} y^{l_2}}^j + \delta_{i_1, i_2, i_3}^{k_3, k_1, k_2} \mathcal{Y}_{y^{l_1} y^{l_2}}^j + \delta_{i_1, i_2, i_3}^{k_2, k_3, k_1} \mathcal{Y}_{y^{l_1} y^{l_2}}^j - \right. \\ &\quad - \delta_{l_1}^j \delta_{i_1, i_2}^{k_2, k_3} \mathcal{X}_{x^{i_3} y^{l_2}}^{k_1} - \delta_{l_1}^j \delta_{i_1, i_3}^{k_2, k_3} \mathcal{X}_{x^{i_2} y^{l_2}}^{k_1} - \delta_{l_1}^j \delta_{i_2, i_3}^{k_2, k_3} \mathcal{X}_{x^{i_1} y^{l_2}}^{k_1} - \\ &\quad - \delta_{l_2}^j \delta_{i_1, i_2}^{k_3, k_1} \mathcal{X}_{x^{i_3} y^{l_1}}^{k_2} - \delta_{l_2}^j \delta_{i_1, i_3}^{k_3, k_1} \mathcal{X}_{x^{i_2} y^{l_1}}^{k_2} - \delta_{l_2}^j \delta_{i_2, i_3}^{k_3, k_1} \mathcal{X}_{x^{i_1} y^{l_1}}^{k_2} - \\ &\quad \left. - \delta_{l_2}^j \delta_{i_1, i_2}^{k_1, k_2} \mathcal{X}_{x^{i_3} y^{l_1}}^{k_3} - \delta_{l_2}^j \delta_{i_1, i_3}^{k_1, k_2} \mathcal{X}_{x^{i_2} y^{l_1}}^{k_3} - \delta_{l_2}^j \delta_{i_2, i_3}^{k_1, k_2} \mathcal{X}_{x^{i_1} y^{l_1}}^{k_3} \right] y_{k_1}^{l_1} y_{k_2}^{l_2} y_{k_3}^{l_3} + \end{aligned}$$

$$\begin{aligned}
& + \sum_{l_1, l_2, l_3=1}^m \sum_{k_1, k_2, k_3, k_4=1}^n \left[-\delta_{l_3}^j \delta_{i_1, i_2, i_3}^{k_1, k_2, k_3} \mathcal{X}_{y^{l_1} y^{l_2}}^{k_4} - \delta_{l_3}^j \delta_{i_1, i_2, i_3}^{k_2, k_3, k_1} \mathcal{X}_{y^{l_1} y^{l_2}}^{k_4} - \delta_{l_3}^j \delta_{i_1, i_2, i_3}^{k_3, k_2, k_1} \mathcal{X}_{y^{l_1} y^{l_2}}^{k_4} - \right. \\
& \quad \left. - \delta_{l_2}^j \delta_{i_1, i_2, i_3}^{k_3, k_4, k_1} \mathcal{X}_{y^{l_1} y^{l_3}}^{k_2} - \delta_{l_2}^j \delta_{i_1, i_2, i_3}^{k_3, k_1, k_4} \mathcal{X}_{y^{l_1} y^{l_3}}^{k_2} - \delta_{l_2}^j \delta_{i_1, i_2, i_3}^{k_1, k_3, k_4} \mathcal{X}_{y^{l_1} y^{l_3}}^{k_2} \right] y_{k_1}^{l_1} y_{k_2}^{l_2} y_{k_3, k_4}^{l_3} + \\
& + \sum_{l_1, l_2=1}^m \sum_{k_1, k_2, k_3, k_4=1}^n \left[-\delta_{l_2}^j \delta_{i_1, i_2, i_3}^{k_1, k_2, k_3} \mathcal{X}_{y^{l_1}}^{k_3} - \delta_{l_2}^j \delta_{i_1, i_2, i_3}^{k_2, k_4, k_1} \mathcal{X}_{y^{l_1}}^{k_3} - \delta_{l_2}^j \delta_{i_1, i_2, i_3}^{k_4, k_1, k_2} \mathcal{X}_{y^{l_1}}^{k_3} \right] y_{k_1, k_2}^{l_1} y_{k_3, k_4}^{l_2} + \\
& + \sum_{l_1=1}^m \sum_{k_1, k_2, k_3=1}^n \left[\delta_{i_1, i_2, i_3}^{k_1, k_2, k_3} \mathcal{Y}_{y^{l_1}}^j - \delta_{l_1}^j \delta_{i_1, i_2}^{k_1, k_2} \mathcal{X}_{x^{i_3}}^{k_3} - \delta_{l_1}^j \delta_{i_1, i_3}^{k_1, k_2} \mathcal{X}_{x^{i_2}}^{k_3} - \delta_{l_1}^j \delta_{i_2, i_3}^{k_1, k_2} \mathcal{X}_{x^{i_1}}^{k_3} \right] y_{k_1, k_2, k_3}^{l_1} + \\
& + \sum_{l_1, l_2=1}^m \sum_{k_1, k_2, k_3, k_4=1}^n \left[-\delta_{l_2}^j \delta_{i_1, i_2, i_3}^{k_1, k_2, k_3} \mathcal{X}_{y^{l_1}}^{k_4} - \delta_{l_2}^j \delta_{i_1, i_2, i_3}^{k_4, k_1, k_2} \mathcal{X}_{y^{l_1}}^{k_3} - \delta_{l_2}^j \delta_{i_1, i_2, i_3}^{k_3, k_4, k_1} \mathcal{X}_{y^{l_1}}^{k_2} - \right. \\
& \quad \left. - \delta_{l_1}^j \delta_{i_1, i_2, i_3}^{k_2, k_3, k_4} \mathcal{X}_{y^{l_2}}^{k_1} \right] y_{k_1}^{l_1} y_{k_2, k_3, k_4}^{l_2}.
\end{aligned}$$

5.5. Final synthesis. To obtain the general formula for $\mathbf{Y}_{i_1, \dots, i_\kappa}^j$, we have to achieve the synthesis between the two formulas (3.74) and (4.19). We start with (3.74) and we modify it until we reach the final formula for $\mathbf{Y}_{i_1, \dots, i_\kappa}^j$.

We have to add the $\mu_1 + \dots + \mu_d$ sums $\sum_{l_{1:1}=1}^m \dots \sum_{l_{1:\mu_1}=1}^m \dots \sum_{l_{d:1}=1}^m \dots \sum_{l_{d:\mu_d}=1}^m$, together with various indices l_α . About these indices, the only point which is not obvious may be analyzed as follows.

A permutation $\sigma \in \mathfrak{S}_{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d}^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)}$ yields the list:

$$\begin{aligned}
(5.6) \quad & \sigma(1:1:1), \dots, \sigma(1:1:\lambda_1), \dots, \sigma(1:\mu_1:1), \dots, \sigma(1:\mu_1:\lambda_1), \dots \\
& \dots, \sigma(d:1:1), \dots, \sigma(1:1:\lambda_d), \dots, \sigma(d:\mu_d:1), \dots, \sigma(d:\mu_d:\lambda_d),
\end{aligned}$$

In the sixth line of (3.74), the last term $\sigma(d:\mu_d:\lambda_d)$ of the above list appears as the subscript of the upper index $k_{\sigma(d:\mu_d:\lambda_d)}$ of the term $\mathcal{X}^{k_{\sigma(d:\mu_d:\lambda_d)}}$. According to the formal rules of Theorem 4.19, we have to multiply the partial derivative of $\mathcal{X}^{k_{\sigma(d:\mu_d:\lambda_d)}}$ by a certain Kronecker symbol $\delta_{l_\alpha}^j$. The question is: *what is the subscript α and how to denote it?*

As explained before the statement of Theorem 4.19, the subscript α is obtained as follows. The term $\sigma(d:\mu_d:\lambda_d)$ is of the form $(e:\nu_d:\gamma_e)$, for some e with $1 \leq e \leq d$, for some ν_e with $1 \leq \nu_e \leq \mu_e$ and for some γ_e with $1 \leq \gamma_e \leq \lambda_e$. The single pure jet variable

$$(5.7) \quad y_{k_{e:\nu_e:1}, \dots, k_{e:\nu_e:\gamma_e}, \dots, k_{e:\nu_e:\lambda_e}}^{l_{e:\nu_e}}$$

appears inside the total monomial

$$(5.8) \quad \prod_{1 \leq \nu_1 \leq \mu_1} y_{k_{1:\nu_1:1}, \dots, k_{1:\nu_1:\lambda_1}}^{l_{1:\nu_1}} \cdots \prod_{1 \leq \nu_d \leq \mu_d} y_{k_{d:\nu_d:1}, \dots, k_{d:\nu_d:\lambda_d}}^{l_{d:\nu_d}},$$

placed at the end of the formula for $\mathbf{Y}_{i_1, \dots, i_\kappa}^j$ (see in advance formula (5.13) below; this total monomial generalizes to several dependent variables the total monomial appearing in the last line of (3.74)). According to the rule explained before the statement of Theorem 4.18, the index l_α must be equal to $l_{e:\nu_e}$, since $l_{e:\nu_e}$ is attached to the monomial (5.7). Coming back to the term $\sigma(d:\mu_d:\lambda_d)$, we shall denote this index by

$$(5.9) \quad l_{e:\nu_e} =: l_{\pi(e:\nu_e:\gamma_e)} =: l_{\pi\sigma(d:\mu_d:\lambda_d)},$$

where the symbol π denotes the projection from the set

$$(5.10) \quad \{1:1:1, \dots, 1:\mu_1:\lambda_1, \dots, d:1:1, \dots, d:\mu_d:\lambda_d\}$$

to the set

$$(5.11) \quad \{1:1, \dots, 1:\mu_1, \dots, d:1, \dots, d:\mu_d\}$$

simply defined by $\pi(e:\nu_e:\gamma_e) := (e:\nu_e)$.

In conclusion, by means of this formalism, we may write down the complete generalization of Theorems 2.24, 3.73 and 4.18 to several independent variables and several dependent variables

Theorem 5.12. *For $j = 1, \dots, m$, for every $\kappa \geq 1$ and for every choice of κ indices i_1, \dots, i_κ in the set $\{1, 2, \dots, n\}$, the general expression of $\mathbf{Y}_{i_1, \dots, i_\kappa}^j$ is as follows:*

(5.13)

$$\begin{aligned} \mathbf{Y}_{i_1, \dots, i_\kappa}^j = & \mathcal{Y}_{x^{i_1} \dots x^{i_\kappa}}^j + \sum_{d=1}^{\kappa+1} \sum_{1 \leq \lambda_1 < \dots < \lambda_d \leq \kappa} \sum_{\mu_1 \geq 1, \dots, \mu_d \geq 1} \sum_{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d \leq \kappa+1} \\ & \sum_{l_{1:1}=1}^m \dots \sum_{l_{1:\mu_1}=1}^m \dots \dots \sum_{l_{d:1}=1}^m \dots \sum_{l_{d:\mu_d}=1}^m \\ & \sum_{k_{1:1:1}, \dots, k_{1:1:\lambda_1}=1}^n \dots \sum_{k_{1:\mu_1:1}, \dots, k_{1:\mu_1:\lambda_1}=1}^n \dots \dots \sum_{k_{d:1:1}, \dots, k_{d:1:\lambda_d}=1}^n \dots \sum_{k_{d:\mu_d:1}, \dots, k_{d:\mu_d:\lambda_d}=1}^n \\ & \left[\sum_{\sigma \in \mathfrak{F}_{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d}^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)}} \sum_{\tau \in \mathfrak{S}_\kappa^{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d}} \delta_{i_\tau(1), \dots, i_\tau(\mu_1 \lambda_1), \dots, i_\tau(\mu_1 \lambda_1 + \dots + \mu_d \lambda_d)}^{k_\sigma(1:1:1), \dots, k_\sigma(1:\mu_1:\lambda_1), \dots, k_\sigma(d:\mu_d:\lambda_d)} \cdot \right. \\ & \quad \frac{\partial^{\kappa - \mu_1 \lambda_1 - \dots - \mu_d \lambda_d + \mu_1 + \dots + \mu_d} \mathcal{Y}^j}{\partial x^{i_\tau(\mu_1 \lambda_1 + \dots + \mu_d \lambda_d + 1)} \dots \partial x^{i_\tau(\kappa)} \partial y^{l_{1:1}} \dots \partial y^{l_{d:\mu_d}}} - \\ & \quad - \sum_{\sigma \in \mathfrak{F}_{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d}^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)}} \sum_{\tau \in \mathfrak{S}_\kappa^{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d - 1}} \delta_{i_\tau(1), \dots, i_\tau(\mu_1 \lambda_1), \dots, i_\tau(\mu_1 \lambda_1 + \dots + \mu_d \lambda_d - 1)}^{k_\sigma(1:1:1), \dots, k_\sigma(1:\mu_1:\lambda_1), \dots, k_\sigma(d:\mu_d:\lambda_d - 1)} \cdot \\ & \quad \cdot \delta_{l_{\pi\sigma}(d:\mu_d:\lambda_d)}^j \cdot \frac{\partial^{\kappa - \mu_1 \lambda_1 - \dots - \mu_d \lambda_d + \mu_1 + \dots + \mu_d} \mathcal{X}^{k_\sigma(d:\mu_d:\lambda_d)}}{\partial x^{i_\tau(\mu_1 \lambda_1 + \dots + \mu_d \lambda_d)} \dots \partial x^{i_\tau(\kappa)} \partial y^{l_{1:1}} \dots \partial y^{\widehat{l_{\pi\sigma}(d:\mu_d:\lambda_d)}} \dots \partial y^{l_{d:\mu_d}}} \left. \right] \\ & \cdot \prod_{1 \leq \nu_1 \leq \mu_1} y_{k_{1:\nu_1:1}, \dots, k_{1:\nu_1:\lambda_1}}^{l_{1:\nu_1}} \dots \prod_{1 \leq \nu_d \leq \mu_d} y_{k_{d:\nu_d:1}, \dots, k_{d:\nu_d:\lambda_d}}^{l_{d:\nu_d}} \end{aligned}$$

In this formula, the coset $\mathfrak{F}_{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d}^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)}$ was defined in equation (3.71); as in Theorem 3.73, we have made the identification:

$$(5.14) \quad \{1, \dots, \kappa\} \equiv \{1:1:1, \dots, 1:\mu_1:\lambda_1, \dots, d:1:1, \dots, d:\mu_d:\lambda_d\}.$$

Since the fundamental monomials appearing in the last line of (4.19) just above are not independent of each other, this formula has still to be modified a little bit. We refer to Section 6 for details.

5.15. Deduction of the most general multivariate Faà di Bruno formula. Let $n \in \mathbb{N}$ with $n \geq 1$, let $x = (x^1, \dots, x^n) \in \mathbb{K}^n$, let $m \in \mathbb{N}$ with $m \geq 1$, let $g^j = g^j(x^1, \dots, x^n)$, $j = 1, \dots, m$, be \mathcal{C}^∞ -smooth functions from \mathbb{K}^n to \mathbb{K}^m , let $y = (y^1, \dots, y^m) \in \mathbb{K}^m$ and let $f = f(y^1, \dots, y^m)$ be a \mathcal{C}^∞ function from \mathbb{K}^m to \mathbb{K} . The goal is to obtain an explicit formula for the partial derivatives of the composition $h := f \circ g$, namely

$$(5.16) \quad h(x^1, \dots, x^n) := f(g^1(x^1, \dots, x^n), \dots, g^m(x^1, \dots, x^n)).$$

For $j = 1, \dots, m$, for $\lambda \in \mathbb{N}$ with $\lambda \geq 1$ and for arbitrary indices $i_1, \dots, i_\lambda = 1, \dots, n$, we shall abbreviate the partial derivative $\frac{\partial^\lambda g^j}{\partial x^{i_1} \dots \partial x^{i_\lambda}}$ by $g_{i_1, \dots, i_\lambda}^j$ and similarly for $h_{i_1, \dots, i_\lambda}$. For

arbitrary indices $l_1, \dots, l_\lambda = 1, \dots, m$, the partial derivative $\frac{\partial^\lambda f}{\partial y^{l_1} \dots \partial y^{l_\lambda}}$ will be abbreviated by $f_{l_1, \dots, l_\lambda}$.

Applying the chain rule, we may compute:

(5.17)

$$\begin{aligned}
 h_{i_1} &= \sum_{l_1=1}^m f_{l_1} \left[g_{i_1}^{l_1} \right], \\
 h_{i_1, i_2} &= \sum_{l_1, l_2=1}^m f_{l_1, l_2} \left[g_{i_1}^{l_1} g_{i_2}^{l_2} \right] + \sum_{l_1=1}^m f_{l_1} \left[g_{i_1, i_2}^{l_1} \right], \\
 h_{i_1, i_2, i_3} &= \sum_{l_1, l_2, l_3=1}^m f_{l_1, l_2, l_3} \left[g_{i_1}^{l_1} g_{i_2}^{l_2} g_{i_3}^{l_3} \right] + \sum_{l_1, l_2=1}^m f_{l_1, l_2} \left[g_{i_1}^{l_1} g_{i_2, i_3}^{l_2} + g_{i_2}^{l_1} g_{i_1, i_3}^{l_2} + g_{i_3}^{l_1} g_{i_1, i_2}^{l_2} \right] + \\
 &\quad + \sum_{l_1=1}^m f_{l_1} \left[g_{i_1, i_2, i_3}^{l_1} \right],
 \end{aligned}$$

$$\begin{aligned}
 h_{i_1, i_2, i_3, i_4} &= \sum_{l_1, l_2, l_3, l_4=1}^m f_{l_1, l_2, l_3, l_4} \left[g_{i_1}^{l_1} g_{i_2}^{l_2} g_{i_3}^{l_3} g_{i_4}^{l_4} \right] + \\
 &\quad \sum_{l_1, l_2, l_3=1}^m f_{l_1, l_2, l_3} \left[g_{i_2}^{l_1} g_{i_3}^{l_2} g_{i_1, i_4}^{l_3} + g_{i_3}^{l_1} g_{i_1}^{l_2} g_{i_2, i_4}^{l_3} + g_{i_1}^{l_1} g_{i_2}^{l_2} g_{i_3, i_4}^{l_3} + \right. \\
 &\quad \left. + g_{i_1}^{l_1} g_{i_4}^{l_2} g_{i_2, i_3}^{l_3} + g_{i_2}^{l_1} g_{i_4}^{l_2} g_{i_3, i_1}^{l_3} + g_{i_3}^{l_1} g_{i_4}^{l_2} g_{i_1, i_2}^{l_3} \right] + \\
 &\quad + \sum_{l_1, l_2=1}^m f_{l_1, l_2} \left[g_{i_1, i_2}^{l_1} g_{i_3, i_4}^{l_2} + g_{i_1, i_3}^{l_1} g_{i_2, i_4}^{l_2} + g_{i_1, i_4}^{l_1} g_{i_2, i_3}^{l_2} \right] + \\
 &\quad + \sum_{l_1, l_2=1}^m f_{l_1, l_2} \left[g_{i_1}^{l_1} g_{i_2, i_3, i_4}^{l_2} + g_{i_2}^{l_1} g_{i_1, i_3, i_4}^{l_2} + g_{i_3}^{l_1} g_{i_1, i_2, i_4}^{l_2} + g_{i_4}^{l_1} g_{i_1, i_2, i_3}^{l_2} \right] + \\
 &\quad + \sum_{l_1=1}^m f_{l_1} \left[g_{i_1, i_2, i_3, i_4}^{l_1} \right].
 \end{aligned}$$

Introducing the derivations

$$\begin{aligned}
 F_i^2 &:= \sum_{k_1=1}^n \sum_{l_1=1}^m g_{k_1, i}^{l_1} \frac{\partial}{\partial g_{k_1}^{l_1}} + \sum_{l_1=1}^m g_i^{l_1} \left(\sum_{l_2=1}^m f_{l_1, l_2} \frac{\partial}{\partial f_{l_2}} \right), \\
 F_i^3 &:= \sum_{k_1=1}^n \sum_{l_1=1}^m g_{k_1, i}^{l_1} \frac{\partial}{\partial g_{k_1}^{l_1}} + \sum_{k_1, k_2=1}^n \sum_{l_1=1}^m g_{k_1, k_2, i}^{l_1} \frac{\partial}{\partial g_{k_1, k_2}^{l_1}} + \\
 &\quad + \sum_{l_1=1}^m g_i^{l_1} \left(\sum_{l_2=1}^m f_{l_1, l_2} \frac{\partial}{\partial f_{l_2}} + \sum_{l_2, l_3=1}^m f_{l_1, l_2, l_3} \frac{\partial}{\partial f_{l_2, l_3}} \right),
 \end{aligned}$$

(5.18)

.....

$$\begin{aligned}
 F_i^\lambda := & \sum_{k_1=1}^n \sum_{l_1=1}^m g_{k_1, i}^{l_1} \frac{\partial}{\partial g_{k_1}^{l_1}} + \sum_{k_1, k_2=1}^n \sum_{l_1=1}^m g_{k_1, k_2, i}^{l_1} \frac{\partial}{\partial g_{k_1, k_2}^{l_1}} + \dots + \\
 & + \sum_{k_1, k_2, \dots, k_{\lambda-1}=1}^n \sum_{l_1=1}^m g_{k_1, k_2, \dots, k_{\lambda-1}, i} \frac{\partial}{\partial g_{k_1, \dots, k_{\lambda-1}}^{l_1}} + \\
 & + \sum_{l_1=1}^m g_i^{l_1} \left(\sum_{l_2=1}^m f_{l_1, l_2} \frac{\partial}{\partial f_{l_2}} + \sum_{l_2, l_3=1}^m f_{l_1, l_2, l_3} \frac{\partial}{\partial f_{l_2, l_3}} + \right. \\
 & \left. + \dots + \sum_{l_2, l_3, \dots, l_\lambda} f_{l_1, l_2, l_3, \dots, l_\lambda} \frac{\partial}{\partial f_{l_2, l_3, \dots, l_\lambda}} \right),
 \end{aligned}$$

we observe that the following induction relations hold:

$$\begin{aligned}
 h_{i_1, i_2} &= F_{i_2}^2(h_{i_1}), \\
 h_{i_1, i_2, i_3} &= F_{i_3}^3(h_{i_1, i_2}), \\
 &\dots\dots\dots \\
 h_{i_1, i_2, \dots, i_\lambda} &= F_{i_\lambda}^\lambda(h_{i_1, i_2, \dots, i_{\lambda-1}}).
 \end{aligned}
 \tag{5.19}$$

To obtain the explicit version of the Faà di Bruno in the case of several variables (x^1, \dots, x^n) and several variables (y^1, \dots, y^m) , it suffices to extract from the expression of $\mathbf{Y}_{i_1, \dots, i_\kappa}^j$ provided by Theorem 5.12 only the terms corresponding to $\mu_1 \lambda_1 + \dots + \mu_d \lambda_d = \kappa$, dropping all the \mathcal{X} terms. After some simplifications and after a translation by means of an elementary dictionary, we obtain the fourth and the most general multivariate Faà di Bruno formula.

Theorem 5.20. *For every integer $\kappa \geq 1$ and for every choice of indices i_1, \dots, i_κ in the set $\{1, 2, \dots, n\}$, the κ -th partial derivative of the composite function*

$$h = h(x^1, \dots, x^n) = f(g^1(x^1, \dots, x^n), \dots, g^m(x^1, \dots, x^n))
 \tag{5.21}$$

with respect to the variables $x^{i_1}, \dots, x^{i_\kappa}$ may be expressed as an explicit polynomial depending on the partial derivatives of f , on the partial derivatives of the g^j and having integer coefficients:

$$\tag{5.22}$$

$$\begin{aligned}
 \frac{\partial^\kappa h}{\partial x^{i_1} \dots \partial x^{i_\kappa}} &= \sum_{d=1}^\kappa \sum_{1 \leq \lambda_1 < \dots < \lambda_d \leq \kappa} \sum_{\mu_1 \geq 1, \dots, \mu_d \geq 1} \sum_{\mu_1 \lambda_1 + \dots + \mu_d \lambda_d = \kappa} \\
 & \sum_{l_{1:1}, \dots, l_{1:\mu_1}=1}^m \dots \sum_{l_{d:1}, \dots, l_{d:\mu_d}=1}^m \frac{\partial^{\mu_1 + \dots + \mu_d} f}{\partial y^{l_{1:1}} \dots \partial y^{l_{1:\mu_1}} \dots \partial y^{l_{d:1}} \dots \partial y^{l_{d:\mu_d}}} \\
 & \left[\sum_{\sigma \in \mathfrak{S}_\kappa^{(\mu_1, \lambda_1), \dots, (\mu_d, \lambda_d)}} \prod_{1 \leq \nu_1 \leq \mu_1} \frac{\partial^{\lambda_1} g^{l_{1:\nu_1}}}{\partial x^{i_{\sigma(1:\nu_1:1)}} \dots \partial x^{i_{\sigma(1:\nu_1:\lambda_1)}}} \dots \right. \\
 & \left. \dots \prod_{1 \leq \nu_d \leq \mu_d} \frac{\partial^{\lambda_d} g^{l_{d:\nu_d}}}{\partial x^{i_{\sigma(d:\nu_d:1)}} \dots \partial x^{i_{\sigma(d:\nu_d:\lambda_d)}}} \right].
 \end{aligned}$$

III: Systems of second order

Table of contents

1. Explicit characterizations of flatness96.
 2. Completely integrable systems of second order ordinary differential equations ... 100.
 3. First and second auxiliary systems107.

§1. EXPLICIT CHARACTERIZATIONS OF FLATNESS

In 1883, S. Lie obtained the following explicit characterization of the local equivalence of a second order ordinary differential equation (\mathcal{E}_1): $y_{xx} = F(x, y, y_x)$ to the Newtonian free particle equation with one degree of freedom $Y_{XX} = 0$. All the functions are assumed to be analytic.

Theorem 1.1. ([Lie1883], pp. 362–365) *Let $\mathbb{K} = \mathbb{R}$ or \mathbb{C} . Let $x \in \mathbb{K}$ and $y \in \mathbb{K}$. A local second order ordinary differential equation $y_{xx} = F(x, y, y_x)$ is equivalent under an invertible point transformation $(x, y) \mapsto (X(x, y), Y(x, y))$ to the free particle equation $Y_{XX} = 0$ if and only if the following two conditions are satisfied:*

- (i) $F_{y_x y_x y_x y_x} = 0$, or equivalently F is a degree three polynomial in y_x , namely there exist four functions G, H, L and M of (x, y) such that F can be written as

$$(1.2) \quad F(x, y, y_x) = G(x, y) + y_x \cdot H(x, y) + (y_x)^2 \cdot L(x, y) + (y_x)^3 \cdot M(x, y);$$

- (ii) the four functions G, H, L and M satisfy the following system of two second order quasi-linear partial differential equations:

$$(1.3) \quad \left\{ \begin{array}{l} 0 = -2 G_{yy} + \frac{4}{3} H_{xy} - \frac{2}{3} L_{xx} + \\ \qquad \qquad \qquad + 2 (G L)_y - 2 G_x M - 4 G M_x + \frac{2}{3} H L_x - \frac{4}{3} H H_y, \\ 0 = -\frac{2}{3} H_{yy} + \frac{4}{3} L_{xy} - 2 M_{xx} + \\ \qquad \qquad \qquad + 2 G M_y + 4 G_y M - 2 (H M)_x - \frac{2}{3} H_y L + \frac{4}{3} L L_x. \end{array} \right.$$

Open question 1.4. *Deduce an explicit necessary and sufficient condition for the associated submanifold of solutions $y = \Pi(x, a, b)$ to be locally equivalent to $Y = B + X A$.*

Assuming $F = F(x, y_x)$ to be independent of y , or equivalently assuming $\mathcal{M}_{(\mathcal{E}_1)}$ to be:

$$(1.5) \quad y = b + \Pi(x, a),$$

the author has checked that equivalence to $Y = B + XA$ holds if and only if two differential rational expressions annihilate:

$$(1.6) \quad \begin{aligned} 0 &= \frac{\Pi_{x^2a^4}}{(\Pi_{xa})^4} - 6 \frac{\Pi_{x^2a^3} \Pi_{xa^2}}{(\Pi_{xa})^5} + 15 \frac{\Pi_{x^2a^2} (\Pi_{xa^2})^2}{(\Pi_{xa})^6} - 4 \frac{\Pi_{x^2a^2} \Pi_{xa^3}}{(\Pi_{xa})^5} \\ &\quad - \frac{\Pi_{x^2a} \Pi_{xa^4}}{(\Pi_{xa})^5} + 10 \frac{\Pi_{xa^3} \Pi_{x^2a} \Pi_{xa^2}}{(\Pi_{xa})^6} - 15 \frac{\Pi_{x^2a} (\Pi_{xa^2})^3}{(\Pi_{xa})^7} \quad \text{and} \\ 0 &= \frac{\Pi_{x^4a^2}}{(\Pi_{xa})^2} - 6 \frac{\Pi_{x^3a^2} \Pi_{x^2a}}{(\Pi_{xa})^3} - 4 \frac{\Pi_{x^3a} \Pi_{x^2a^2}}{(\Pi_{xa})^3} - \frac{\Pi_{x^4a} \Pi_{xa^2}}{(\Pi_{xa})^3} + \\ &\quad + 15 \frac{\Pi_{x^2a^2} (\Pi_{x^2a})^2}{(\Pi_{xa})^4} + 10 \frac{\Pi_{x^3a} \Pi_{x^2a} \Pi_{xa^2}}{(\Pi_{xa})^4} - 15 \frac{(\Pi_{x^2a})^3 \Pi_{xa^2}}{(\Pi_{xa})^5}. \end{aligned}$$

As an application, this characterizes local sphericity of a rigid hypersurface $w = \bar{w} + i\Theta(z, \bar{z})$ of \mathbb{C}^2 . The answer for a general $y = \Pi(x, a, b)$, together with a proof, will appear elsewhere.

A modern restitution of Lie's original proof of Theorem 1.1 may be found in [Me2004]. In this reference, we generalize Theorem 1.1 to several dependent variables $y = (y^1, y^2, \dots, y^m)$. In the present Part III, we will instead pass to several independent variables $x = (x^1, x^2, \dots, x^n)$.

Theorem 1.7. *Let $\mathbb{K} = \mathbb{R}$ or \mathbb{C} , let $n \in \mathbb{N}$, suppose $n \geq 2$ and consider a system of completely integrable partial differential equations in n independent variables $x = (x^1, \dots, x^n) \in \mathbb{K}^n$ and in one dependent variable $y \in \mathbb{K}$ of the form:*

$$(1.8) \quad y_{x^{j_1}x^{j_2}}(x) = F^{j_1, j_2}(x, y(x), y_{x^1}(x), \dots, y_{x^n}(x)), \quad 1 \leq j_1, j_2 \leq n,$$

where $F^{j_1, j_2} = F^{j_2, j_1}$. Under a local change of coordinates $(x, y) \mapsto (X, Y) = (X(x, y), Y(x, y))$, this system (1.8) is equivalent to the simplest "flat" system

$$(1.9) \quad Y_{X^{j_1}X^{j_2}} = 0, \quad 1 \leq j_1, j_2 \leq n,$$

if and only if there exist arbitrary functions G_{j_1, j_2} , $H_{j_1, j_2}^{k_1}$, $L_{j_1}^{k_1}$ and M^{k_1} of the variables (x, y) , for $1 \leq j_1, j_2, k_1 \leq n$, satisfying the two symmetry conditions $G_{j_1, j_2} = G_{j_2, j_1}$ and $H_{j_1, j_2}^{k_1} = H_{j_2, j_1}^{k_1}$, such that the equation (1.8) is of the specific cubic polynomial form:

$$(1.10) \quad y_{x^{j_1}x^{j_2}} = G_{j_1, j_2} + \sum_{k_1=1}^n y_{x^{k_1}} \left(H_{j_1, j_2}^{k_1} + \frac{1}{2} y_{x^{j_1}} L_{j_2}^{k_1} + \frac{1}{2} y_{x^{j_2}} L_{j_1}^{k_1} + y_{x^{j_1}} y_{x^{j_2}} M^{k_1} \right),$$

for $j_1, j_2 = 1, \dots, n$.

It may seem quite paradoxical and counter-intuitive (or even false?) that every system (1.10), for arbitrary choices of functions G_{j_1, j_2} , $H_{j_1, j_2}^{k_1}$, $L_{j_1}^{k_1}$ and M^{k_1} , is automatically equivalent to $Y_{X^{j_1}X^{j_2}} = 0$. However, a strong hidden assumption holds: that of complete integrability. Shortly, this crucial condition amounts to say that

$$(1.11) \quad D_{j_3} (F^{j_1, j_2}) = D_{j_2} (F^{j_1, j_3}),$$

for all $j_1, j_2, j_3 = 1, \dots, n$, where, for $j = 1, \dots, n$, the D_j are the total differentiation operators defined by

$$(1.12) \quad D_j := \frac{\partial}{\partial x^j} + y_{x^j} \frac{\partial}{\partial y} + \sum_{l=1}^n F^{j, l} \frac{\partial}{\partial y_{x^l}}.$$

These conditions are non-void precisely when $n \geq 2$. More concretely, developing out (1.11) when the F^{j_1, j_2} are of the specific cubic polynomial form (1.10), after some nontrivial manual computation, we obtain the complicated cubic differential polynomial in the variables y_{x^k} . Equating to zero all the coefficients of this cubic polynomial, we obtain four families (I'), (II'), (III') and (IV') of first order partial differential equations satisfied by G_{j_1, j_2} , $H_{j_1, j_2}^{k_1}$, $L_{j_1}^{k_1}$ and M^{k_1} :

$$(I') \quad \left\{ 0 = G_{j_1, j_2, x^{j_3}} - G_{j_1, j_3, x^{j_2}} + \sum_{k_1=1}^n H_{j_1, j_2}^{k_1} G_{k_1, j_3} - \sum_{k_1=1}^n H_{j_1, j_3}^{k_1} G_{k_1, j_2} \right.$$

$$(II') \quad \left\{ \begin{aligned} 0 = & \delta_{j_3}^{k_1} G_{j_1, j_2, y} - \delta_{j_2}^{k_1} G_{j_1, j_3, y} + H_{j_1, j_2, x^{j_3}}^{k_1} - H_{j_1, j_3, x^{j_2}}^{k_1} + \\ & + \frac{1}{2} G_{j_1, j_3} L_{j_2}^{k_1} - \frac{1}{2} G_{j_1, j_2} L_{j_3}^{k_1} + \\ & + \frac{1}{2} \delta_{j_1}^{k_1} \sum_{k_2=1}^n G_{k_2, j_3} L_{j_2}^{k_2} - \frac{1}{2} \delta_{j_1}^{k_1} \sum_{k_2=1}^n G_{k_2, j_2} L_{j_3}^{k_2} + \\ & + \frac{1}{2} \delta_{j_2}^{k_1} \sum_{k_2=1}^n G_{k_2, j_3} L_{j_1}^{k_2} - \frac{1}{2} \delta_{j_3}^{k_1} \sum_{k_2=1}^n G_{k_2, j_2} L_{j_1}^{k_2} + \\ & + \sum_{k_2=1}^n H_{k_2, j_3}^{k_1} H_{j_1, j_2}^{k_2} - \sum_{k_2=1}^n H_{k_2, j_2}^{k_1} H_{j_1, j_3}^{k_2} \end{aligned} \right.$$

$$(III') \quad \left\{ \begin{aligned} 0 = & \sum_{\sigma \in \mathfrak{S}_2} \left(\delta_{j_3}^{k_{\sigma(2)}} H_{j_1, j_2, y}^{k_{\sigma(1)}} - \delta_{j_2}^{k_{\sigma(2)}} H_{j_1, j_3, y}^{k_{\sigma(1)}} + \right. \\ & + \frac{1}{2} \delta_{j_2}^{k_{\sigma(2)}} L_{j_1, x^{j_3}}^{k_{\sigma(1)}} - \frac{1}{2} \delta_{j_3}^{k_{\sigma(2)}} L_{j_1, x^{j_2}}^{k_{\sigma(1)}} + \\ & + \frac{1}{2} \delta_{j_1}^{k_{\sigma(2)}} L_{j_2, x^{j_3}}^{k_{\sigma(1)}} - \frac{1}{2} \delta_{j_1}^{k_{\sigma(2)}} L_{j_3, x^{j_2}}^{k_{\sigma(1)}} + \\ & + \delta_{j_2}^{k_{\sigma(2)}} G_{j_1, j_3} M^{k_{\sigma(1)}} - \delta_{j_3}^{k_{\sigma(2)}} G_{j_1, j_2} M^{k_{\sigma(1)}} + \\ & + \delta_{j_1, j_2}^{k_{\sigma(1)}, k_{\sigma(2)}} \sum_{k_3=1}^n G_{k_3, j_3} M^{k_3} - \delta_{j_1, j_3}^{k_{\sigma(1)}, k_{\sigma(2)}} \sum_{k_3=1}^n G_{k_3, j_2} M^{k_3} + \\ & + \frac{1}{2} \delta_{j_1}^{k_{\sigma(1)}} \sum_{k_3=1}^n H_{k_3, j_3}^{k_{\sigma(2)}} L_{j_2}^{k_3} - \frac{1}{2} \delta_{j_1}^{k_{\sigma(1)}} \sum_{k_3=1}^n H_{k_3, j_2}^{k_{\sigma(2)}} L_{j_3}^{k_3} + \\ & + \frac{1}{2} \delta_{j_2}^{k_{\sigma(1)}} \sum_{k_3=1}^n H_{k_3, j_3}^{k_{\sigma(2)}} L_{j_1}^{k_3} - \frac{1}{2} \delta_{j_3}^{k_{\sigma(1)}} \sum_{k_3=1}^n H_{k_3, j_2}^{k_{\sigma(2)}} L_{j_1}^{k_3} + \\ & \left. + \frac{1}{2} \delta_{j_3}^{k_{\sigma(1)}} \sum_{k_3=1}^n H_{j_1, j_2}^{k_3} L_{k_3}^{k_{\sigma(2)}} - \frac{1}{2} \delta_{j_2}^{k_{\sigma(1)}} \sum_{k_3=1}^n H_{j_1, j_3}^{k_3} L_{k_3}^{k_{\sigma(2)}} \right) \end{aligned} \right.$$

$$(IV') \left\{ \begin{array}{l} 0 = \sum_{\sigma \in \mathfrak{S}_3} \left(\frac{1}{2} \delta_{j_3, j_1}^{k_{\sigma(3)}, k_{\sigma(2)}} L_{j_2, y}^{k_{\sigma(1)}} - \frac{1}{2} \delta_{j_2, j_1}^{k_{\sigma(3)}, k_{\sigma(2)}} L_{j_3, y}^{k_{\sigma(1)}} + \right. \\ \left. + \delta_{j_2, j_1}^{k_{\sigma(3)}, k_{\sigma(2)}} M_{x^{j_3}}^{k_{\sigma(1)}} - \delta_{j_3, j_1}^{k_{\sigma(3)}, k_{\sigma(2)}} M_{x^{j_2}}^{k_{\sigma(1)}} + \right. \\ \left. + \delta_{j_2, j_1}^{k_{\sigma(3)}, k_{\sigma(1)}} \sum_{k_4=1}^n H_{k_4, j_3}^{k_{\sigma(2)}} M^{k_4} - \delta_{j_3, j_1}^{k_{\sigma(3)}, k_{\sigma(1)}} \sum_{k_4=1}^n H_{k_4, j_2}^{k_{\sigma(2)}} M^{k_4} + \right. \\ \left. + \frac{1}{4} \delta_{j_1, j_3}^{k_{\sigma(1)}, k_{\sigma(3)}} \sum_{k_4=1}^n L_{k_4}^{k_{\sigma(2)}} L_{j_2}^{k_4} - \frac{1}{4} \delta_{j_1, j_2}^{k_{\sigma(1)}, k_{\sigma(3)}} \sum_{k_4=1}^n L_{k_4}^{k_{\sigma(2)}} L_{j_3}^{k_4} \right). \end{array} \right.$$

(These systems (I'), (II'), (III') and (IV') should be distinguished from the systems (I), (II), (III) and (IV) of Theorem 1.7 in [Me2004], although they are quite similar.) Here, the indices $j_1, j_2, j_3, k_1, k_2, k_3$ vary in $\{1, 2, \dots, n\}$. By \mathfrak{S}_2 and by \mathfrak{S}_3 , we denote the permutation group of $\{1, 2\}$ and of $\{1, 2, 3\}$. To facilitate hand- and Latex-writing, partial derivatives are denoted as indices after a comma; for instance, $G_{j_1, j_2, x^{j_3}}$ is an abbreviation for $\partial G_{j_1, j_2} / \partial x^{j_3}$. To deduce (I'), (II'), (III') and (IV') from equation (1.11), we use the fact that every cubic polynomial equation of the form

$$(1.13) \quad 0 \equiv A + \sum_{k_1=1}^n B_{k_1} \cdot y_{x^{k_1}} + \sum_{k_1=1}^n \sum_{k_2=1}^n C_{k_1, k_2} \cdot y_{x^{k_1}} y_{x^{k_2}} + \sum_{k_1=1}^n \sum_{k_2=1}^n \sum_{k_3=1}^n D_{k_1, k_2, k_3} \cdot y_{x^{k_1}} y_{x^{k_2}} y_{x^{k_3}}$$

is equivalent to the annihilation of the following symmetric sums of its coefficients:

$$(1.14) \quad \left\{ \begin{array}{l} 0 = A, \\ 0 = B_{k_1}, \\ 0 = C_{k_1, k_2} + C_{k_2, k_1}, \\ 0 = D_{k_1, k_2, k_3} + D_{k_3, k_1, k_2} + D_{k_2, k_3, k_1} + D_{k_2, k_1, k_3} + D_{k_3, k_2, k_1} + D_{k_1, k_3, k_2}. \end{array} \right.$$

for all $k_1, k_2, k_3 = 1, \dots, n$.

In conclusion, the functions G_{j_1, j_2} , $H_{j_1, j_2}^{k_1}$, $L_{j_1}^{k_1}$ and M^{k_1} in the statement of Theorem 1.7 are far from being arbitrary: they satisfy the complicated system of first order partial differential equations (I'), (II'), (III') and (IV') above.

Our proof of Theorem 1.7 is similar to the one provided in [Me2004], in the case of systems of second order ordinary differential equations, so that most steps of the proof will be summarized.

In the end of this paper, we will delineate a complicated system of *second* order partial differential equations satisfied by G_{j_1, j_2} , $H_{j_1, j_2}^{k_1}$, $L_{j_1}^{k_1}$ and M^{k_1} which is the exact analog of the system described in the abstract. The main technical part of the proof of Theorem 1.7 will be to establish that this second order system is a consequence, by linear combinations and by differentiations, of the first order system (I'), (II'), (III') and (IV').

Open question 1.15. *Are Theorems 1.1 and 1.7 true under weaker smoothness assumptions, namely with a C^2 or a $W_{\text{loc}}^{1, \infty}$ right-hand side ?*

We refer to [Ma2003] for inspiration and appropriate tools.

Open question 1.16. *Deduce from Theorem 1.7 an explicit necessary and sufficient condition for the associated submanifold of solutions $y = b + \Pi(x^i, a^k, b)$ to be locally equivalent to $Y = B + X^1 A^1 + \dots + X^n A^n$.*

As an application, this would characterize local sphericity of a Levi nondegenerate hypersurface $M \subset \mathbb{C}^{n+1}$ with $n \geq 2$.

Generalizing the Lie-Tresse classification would be a great achievement.

Open problem 1.17. *For $n = 2$ establish a complete list of normal forms of all possible systems (1.?) according to their Lie symmetry group. In case of success, classify Levi nondegenerate real analytic hypersurfaces of \mathbb{C}^3 up to biholomorphisms.*

§2. COMPLETELY INTEGRABLE SYSTEMS OF SECOND ORDER ORDINARY DIFFERENTIAL EQUATIONS

2.1. Prolongation of a point transformation to the second order jet space. Let $\mathbb{K} = \mathbb{R}$ or \mathbb{C} , let $n \in \mathbb{N}$, suppose $n \geq 2$, let $x = (x^1, \dots, x^n) \in \mathbb{K}^n$ and let $y \in \mathbb{K}$. According to the main assumption of Theorem 1.7, we have to consider a local \mathbb{K} -analytic diffeomorphism of the form

$$(2.2) \quad (x^{j_1}, y) \longmapsto (X^j(x^{j_1}, y), Y(x^{j_1}, y)),$$

which transforms the system (1.8) to the system $Y_{X^{i_1 X^{j_2}}} = 0$, $1 \leq j_1, j_2 \leq n$. Without loss of generality, we shall assume that this transformation is close to the identity. To obtain the precise expression (2.35) of the transformed system (1.8), we have to prolong the above diffeomorphism to the second order jet space. We introduce the coordinates $(x^j, y, y_{x^{j_1}}, y_{x^{j_1 x^{j_2}}})$ on the second order jet space. Let

$$(2.3) \quad D_k := \frac{\partial}{\partial x^k} + y_{x^k} \frac{\partial}{\partial y} + \sum_{l=1}^n y_{x^k x^l} \frac{\partial}{\partial y_{x^l}},$$

be the k -th total differentiation operator. According to [O11986, BK1989, O11995], for the first order partial derivatives, one has the (implicit, compact) expression:

$$(2.4) \quad \begin{pmatrix} Y_{X^1} \\ \vdots \\ Y_{X^n} \end{pmatrix} = \begin{pmatrix} D_1 X^1 & \cdots & D_1 X^n \\ \vdots & \cdots & \vdots \\ D_n X^1 & \cdots & D_n X^n \end{pmatrix}^{-1} \begin{pmatrix} D_1 Y \\ \vdots \\ D_n Y \end{pmatrix},$$

where $(\cdot)^{-1}$ denotes the inverse matrix, which exists, since the transformation (2.2) is close to the identity. For the second order partial derivatives, again according to [O11986, BK1989, O11995], one has the (implicit, compact) expressions:

$$(2.5) \quad \begin{pmatrix} Y_{X^j X^1} \\ \vdots \\ Y_{X^j X^n} \end{pmatrix} = \begin{pmatrix} D_1 X^1 & \cdots & D_1 X^n \\ \vdots & \cdots & \vdots \\ D_n X^1 & \cdots & D_n X^n \end{pmatrix}^{-1} \begin{pmatrix} D_1 Y_{X^j} \\ \vdots \\ D_n Y_{X^j} \end{pmatrix},$$

for $j = 1, \dots, n$. Let DX denote the matrix $(D_i X^j)_{\substack{1 \leq j \leq n \\ 1 \leq i \leq n}}$, where i is the index of lines and j the index of columns, let Y_X denote the column matrix $(Y_{X^i})_{1 \leq i \leq n}$ and let DY be the column matrix $(D_i Y)_{1 \leq i \leq n}$.

By inspecting (2.5) above, we see that the equivalence between (i), (ii) and (iii) just below is obvious:

Lemma 2.6. *The following conditions are equivalent:*

- (i) *the differential equations $Y_{X^j X^k} = 0$ hold for $1 \leq j, k \leq n$;*
- (ii) *the matrix equations $D_k(Y_X) = 0$ hold for $1 \leq k \leq n$;*
- (iii) *the matrix equations $DX \cdot D_k(Y_X) = 0$ hold for $1 \leq k \leq n$;*
- (iv) *the matrix equations $0 = D_k(DX) \cdot Y_X - D_k(DY)$ hold for $1 \leq k \leq n$.*

Formally, in the sequel, it will be more convenient to achieve the explicit computations starting from condition **(iv)**, since no matrix inversion at all is involved in it.

Proof. Indeed, applying the total differentiation operator D_k to the matrix equation (2.4) written under the equivalent form $0 = DX \cdot Y_X - DY$, we get:

$$(2.7) \quad 0 = D_k(DX) \cdot Y_X + DX \cdot D_k(Y_X) - D_k(DY),$$

so that the equivalence between **(iii)** and **(iv)** is now clear. \square

2.8. An explicit formula in the case $n = 2$. Thus, we can start to develop explicitly the matrix equations

$$(2.9) \quad 0 = D_k(DX) \cdot Y_X - D_k(DY).$$

In it, some huge formal expressions are hidden behind the symbol D_k . Proceeding inductively, we start by examining the case $n = 2$ thoroughly. By direct computations which require to be clever, we reconstitute some 3×3 determinants in the four (in fact three) developed equations (2.9). After some work, the first equation is:

$$(2.10) \quad \begin{aligned} 0 = & y_{x^1 x^1} \cdot \left| \begin{array}{ccc} X_{x^1}^1 & X_{x^2}^1 & X_y^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_y^2 \\ Y_{x^1} & Y_{x^2} & Y_y \end{array} \right| + \left| \begin{array}{ccc} X_{x^1}^1 & X_{x^2}^1 & X_{x^1 x^1}^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_{x^1 x^1}^2 \\ Y_{x^1} & Y_{x^2} & Y_{x^1 x^1} \end{array} \right| + \\ & + y_{x^1} \cdot \left\{ 2 \left| \begin{array}{ccc} X_{x^1}^1 & X_{x^2}^1 & X_{x^1 y}^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_{x^1 y}^2 \\ Y_{x^1} & Y_{x^2} & Y_{x^1 y} \end{array} \right| - \left| \begin{array}{ccc} X_{x^1 x^1}^1 & X_{x^2}^1 & X_y^1 \\ X_{x^1 x^1}^2 & X_{x^2}^2 & X_y^2 \\ Y_{x^1 x^1} & Y_{x^2} & Y_y \end{array} \right| \right\} + \\ & + y_{x^2} \cdot \left\{ - \left| \begin{array}{ccc} X_{x^1}^1 & X_{x^1 x^1}^1 & X_y^1 \\ X_{x^1}^2 & X_{x^1 x^1}^2 & X_y^2 \\ Y_{x^1} & Y_{x^1 x^1} & Y_y \end{array} \right| \right\} + \\ & + y_{x^1} y_{x^1} \cdot \left\{ \left| \begin{array}{ccc} X_{x^1}^1 & X_{x^2}^1 & X_{yy}^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_{yy}^2 \\ Y_{x^1} & Y_{x^2} & Y_{yy} \end{array} \right| - 2 \left| \begin{array}{ccc} X_{x^1 y}^1 & X_{x^2}^1 & X_y^1 \\ X_{x^1 y}^2 & X_{x^2}^2 & X_y^2 \\ Y_{x^1 y} & Y_{x^2} & Y_y \end{array} \right| \right\} + \\ & + y_{x^1} y_{x^2} \cdot \left\{ -2 \left| \begin{array}{ccc} X_{x^1}^1 & X_{x^1 y}^1 & X_y^1 \\ X_{x^1}^2 & X_{x^1 y}^2 & X_y^2 \\ Y_{x^1} & Y_{x^1 y} & Y_y \end{array} \right| \right\} + \\ & + y_{x^1} y_{x^1} y_{x^1} \cdot \left\{ - \left| \begin{array}{ccc} X_{yy}^1 & X_{x^2}^1 & X_y^1 \\ X_{yy}^2 & X_{x^2}^2 & X_y^2 \\ Y_{yy} & Y_{x^2} & Y_y \end{array} \right| \right\} + \\ & + y_{x^1} y_{x^1} y_{x^2} \cdot \left\{ - \left| \begin{array}{ccc} X_{x^1}^1 & X_{yy}^1 & X_y^1 \\ X_{x^1}^2 & X_{yy}^2 & X_y^2 \\ Y_{x^1} & Y_{yy} & Y_y \end{array} \right| \right\}. \end{aligned}$$

This formula and the two next (2.22), (2.23) have been checked by Sylvain Neut and Michel Petitot with the help of Maple.

2.11. Comparison with the coefficients of the second prolongation of a vector field.

At present, it is useful to make an illuminating digression which will help us to devise what is the general form of the development of the equations (2.9). Consider an arbitrary vector field of the form

$$(2.12) \quad \mathcal{L} := \sum_{k=1}^n \mathcal{X}^k \frac{\partial}{\partial x^k} + \mathcal{Y} \frac{\partial}{\partial y},$$

where the coefficients \mathcal{X}^k and \mathcal{Y} are functions of (x^i, y) . According to [Ol1986, BK1989, Ol1995], there exists a unique prolongation $\mathcal{L}^{(2)}$ of this vector field to the second order jet space, of the form

$$(2.13) \quad \mathcal{L}^{(2)} := \mathcal{L} + \sum_{j_1=1}^n \mathbf{Y}_{j_1} \frac{\partial}{\partial y_{x^{j_1}}} + \sum_{j_1=1}^n \sum_{j_2=1}^n \mathbf{Y}_{j_1, j_2} \frac{\partial}{\partial y_{x^{j_1} x^{j_2}}},$$

where the coefficients \mathbf{Y}_{j_1} , \mathbf{Y}_{j_1, j_2} may be computed by means of formulas (3.4) of Section 3(II). In Part II, we obtained the following perfect formulas:

$$(2.14) \quad \left\{ \begin{array}{l} \mathbf{Y}_{j_1, j_2} = \mathcal{Y}_{x^{j_1} x^{j_2}} + \sum_{k_1=1}^n y_{x^{k_1}} \cdot \left\{ \delta_{j_1}^{k_1} \mathcal{Y}_{x^{j_2} y} + \delta_{j_2}^{k_1} \mathcal{Y}_{x^{j_1} y} - \mathcal{X}_{x^{j_1} x^{j_2}}^{k_1} \right\} + \\ + \sum_{k_1=1}^n \sum_{k_2=1}^n y_{x^{k_1}} y_{x^{k_2}} \cdot \left\{ \delta_{j_1, j_2}^{k_1, k_2} \mathcal{Y}_{yy} - \delta_{j_1}^{k_1} \mathcal{X}_{x^{j_2} y}^{k_2} - \delta_{j_2}^{k_1} \mathcal{X}_{x^{j_1} y}^{k_2} \right\} + \\ + \sum_{k_1=1}^n \sum_{k_2=1}^n \sum_{k_3=1}^n y_{x^{k_1}} y_{x^{k_2}} y_{x^{k_3}} \cdot \left\{ -\delta_{j_1, j_2}^{k_1, k_2} \mathcal{X}_{yy}^{k_3} \right\}, \end{array} \right.$$

for $j_1, j_2 = 1, \dots, n$. The expression of \mathbf{Y}_{j_1} does not matter for us here. Specifying this formula to the case $n = 2$ and taking account of the symmetry $\mathbf{Y}_{1,2} = \mathbf{Y}_{2,1}$ we get the following three second order coefficients:

$$(2.15) \quad \left\{ \begin{array}{l} \mathbf{Y}_{1,1} = \mathcal{Y}_{x^1 x^1} + y_{x^1} \cdot \left\{ 2 \mathcal{Y}_{x^1 y} - \mathcal{X}_{x^1 x^1}^1 \right\} + y_{x^2} \cdot \left\{ -\mathcal{X}_{x^1 x^1}^2 \right\} + \\ + y_{x^1} y_{x^1} \cdot \left\{ \mathcal{Y}_{yy} - 2 \mathcal{X}_{x^1 y}^1 \right\} + y_{x^1} y_{x^2} \cdot \left\{ -2 \mathcal{X}_{x^1 y}^2 \right\} + \\ + y_{x^1} y_{x^1} y_{x^1} \cdot \left\{ -\mathcal{X}_{yy}^1 \right\} + y_{x^1} y_{x^1} y_{x^2} \cdot \left\{ -\mathcal{X}_{yy}^2 \right\}, \\ \mathbf{Y}_{1,2} = \mathcal{Y}_{x^1 x^2} + y_{x^1} \cdot \left\{ \mathcal{Y}_{x^2 y} - \mathcal{X}_{x^1 x^2}^1 \right\} + y_{x^2} \cdot \left\{ \mathcal{Y}_{x^1 y} - \mathcal{X}_{x^1 x^2}^2 \right\} + \\ + y_{x^1} y_{x^1} \cdot \left\{ -\mathcal{X}_{x^2 y}^1 \right\} + y_{x^1} y_{x^2} \cdot \left\{ \mathcal{Y}_{yy} - \mathcal{X}_{x^1 y}^1 - \mathcal{X}_{x^2 y}^2 \right\} + \\ + y_{x^2} y_{x^2} \cdot \left\{ -\mathcal{X}_{x^1 y}^2 \right\} + \\ + y_{x^1} y_{x^1} y_{x^2} \cdot \left\{ -\mathcal{X}_{yy}^1 \right\} + y_{x^1} y_{x^2} y_{x^2} \cdot \left\{ -\mathcal{X}_{yy}^2 \right\}, \\ \mathbf{Y}_{2,2} = \mathcal{Y}_{x^2 x^2} + y_{x^1} \cdot \left\{ -\mathcal{X}_{x^2 x^2}^1 \right\} + y_{x^2} \cdot \left\{ 2 \mathcal{Y}_{x^2 y} - \mathcal{X}_{x^2 x^2}^2 \right\} + \\ + y_{x^1} y_{x^2} \cdot \left\{ -2 \mathcal{X}_{x^2 y}^1 \right\} + y_{x^2} y_{x^2} \cdot \left\{ \mathcal{Y}_{yy} - 2 \mathcal{X}_{x^2 y}^2 \right\} + \\ + y_{x^1} y_{x^2} y_{x^2} \cdot \left\{ -\mathcal{X}_{yy}^1 \right\} + y_{x^2} y_{x^2} y_{x^2} \cdot \left\{ -\mathcal{X}_{yy}^2 \right\}. \end{array} \right.$$

We would like to mention that the computation of \mathbf{Y}_{j_1, j_2} , $1 \leq j_1, j_2 \leq 2$, above is easier than the verification of (2.10). Based on the three formulas (2.15), we claim that we can guess the second and the third equations, which would be obtained by developing and by simplifying (2.9), namely with $y_{x^1 x^2}$ and with $y_{x^2 x^2}$ instead of $y_{x^1 x^2}$ in (2.10). Our dictionary to translate from the first formula (2.15) to (2.10) may be described as follows. Begin with the *Jacobian determinant*

$$(2.16) \quad \left| \begin{array}{ccc} X_{x^1}^1 & X_{x^2}^1 & X_y^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_y^2 \\ Y_{x^1} & Y_{x^2} & Y_y \end{array} \right|$$

of the change of coordinates (2.2). Since this change of coordinates is close to the identity, we may consider that the following Jacobian matrix approximation holds:

$$(2.17) \quad \left(\begin{array}{ccc} X_{x^1}^1 & X_{x^2}^1 & X_y^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_y^2 \\ Y_{x^1} & Y_{x^2} & Y_y \end{array} \right) \cong \left(\begin{array}{ccc} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{array} \right).$$

The jacobian matrix has three columns. There are six possible second order derivatives with respect to the variables (x^1, x^2, y) , namely

$$(2.18) \quad (\cdot)_{x^1x^1}, \quad (\cdot)_{x^1x^2}, \quad (\cdot)_{x^2x^2}, \quad (\cdot)_{x^1y}, \quad (\cdot)_{x^2y}, \quad (\cdot)_{yy}.$$

In the Jacobian determinant (2.16), by replacing any one of the three columns of first order derivatives with a column of second order derivatives, we obtain exactly $3 \times 6 = 18$ possible determinants. For instance, by replacing the third column by the second order derivative $(\cdot)_{x^1y}$ or the first column by the second order derivative $(\cdot)_{x^1x^1}$, we get:

$$(2.19) \quad \begin{vmatrix} X_{x^1}^1 & X_{x^2}^1 & X_{x^1y}^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_{x^1y}^2 \\ Y_{x^1} & Y_{x^2} & Y_{x^1y} \end{vmatrix} \quad \text{or} \quad \begin{vmatrix} X_{x^1x^1}^1 & X_{x^2}^1 & X_y^1 \\ X_{x^1x^1}^2 & X_{x^2}^2 & X_y^2 \\ Y_{x^1x^1} & Y_{x^2} & Y_y \end{vmatrix}.$$

We recover the two determinants appearing in the second line of (2.10). On the other hand, according to the approximation (2.17), these two determinants are essentially equal to

$$(2.20) \quad \begin{vmatrix} 1 & 0 & X_{x^1y}^1 \\ 0 & 1 & X_{x^1y}^2 \\ 0 & 0 & Y_{x^1y} \end{vmatrix} = Y_{x^1y} \quad \text{or to} \quad \begin{vmatrix} X_{x^1x^1}^1 & 0 & 0 \\ X_{x^1x^1}^2 & 1 & 0 \\ Y_{x^1x^1} & 0 & 1 \end{vmatrix} = X_{x^1x^1}^1.$$

Consequently, in the second line of (2.10), up to a change to calligraphic letters, we recover the coefficient

$$(2.21) \quad 2\mathcal{Y}_{x^1y} - \mathcal{X}_{x^1x^1}^1$$

of y_{x_1} in the expression of $\mathbf{Y}_{1,1}$ in (2.15). In conclusion, we have discovered how to pass symbolically from the first equation (2.15) to the equation (2.10) and conversely.

Translating the second equation (2.15), we deduce, *without any further computation*, that the second equation which would be obtained by developing (2.9) in length, is:

$$(2.22) \quad \begin{aligned} 0 = & y_{x^1x^2} \cdot \left[\begin{vmatrix} X_{x^1}^1 & X_{x^2}^1 & X_y^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_y^2 \\ Y_{x^1} & Y_{x^2} & Y_y \end{vmatrix} + \begin{vmatrix} X_{x^1}^1 & X_{x^2}^1 & X_{x^1x^2}^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_{x^1x^2}^2 \\ Y_{x^1} & Y_{x^2} & Y_{x^1x^2} \end{vmatrix} + \right. \\ & + y_{x^1} \cdot \left\{ \begin{vmatrix} X_{x^1}^1 & X_{x^2}^1 & X_{x^2y}^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_{x^2y}^2 \\ Y_{x^1} & Y_{x^2} & Y_{x^2y} \end{vmatrix} - \begin{vmatrix} X_{x^1x^2}^1 & X_{x^2}^1 & X_y^1 \\ X_{x^1x^2}^2 & X_{x^2}^2 & X_y^2 \\ Y_{x^1x^2} & Y_{x^2} & Y_y \end{vmatrix} \right\} + \\ & + y_{x^2} \cdot \left\{ \begin{vmatrix} X_{x^1}^1 & X_{x^2}^1 & X_{x^1y}^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_{x^1y}^2 \\ Y_{x^1} & Y_{x^2} & Y_{x^1y} \end{vmatrix} - \begin{vmatrix} X_{x^1}^1 & X_{x^1x^2}^1 & X_y^1 \\ X_{x^1}^2 & X_{x^1x^2}^2 & X_y^2 \\ Y_{x^1} & Y_{x^1x^2} & Y_y \end{vmatrix} \right\} + \end{aligned}$$

$$\begin{aligned}
& + y_{x^1} y_{x^1} \cdot \left\{ - \left| \begin{array}{ccc} X_{x^2 y}^1 & X_{x^2}^1 & X_y^1 \\ X_{x^2 y}^2 & X_{x^2}^2 & X_y^2 \\ Y_{x^2 y} & Y_{x^2} & Y_y \end{array} \right| \right\} + \\
& + y_{x^1} y_{x^2} \cdot \left\{ \left| \begin{array}{ccc} X_{x^1}^1 & X_{x^2}^1 & X_{yy}^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_{yy}^2 \\ Y_{x^1} & Y_{x^2} & Y_{yy} \end{array} \right| - \left| \begin{array}{ccc} X_{x^1 y}^1 & X_{x^2}^1 & X_y^1 \\ X_{x^1 y}^2 & X_{x^2}^2 & X_y^2 \\ Y_{x^1 y} & Y_{x^2} & Y_y \end{array} \right| - \right. \\
& - \left. \left| \begin{array}{ccc} X_{x^1}^1 & X_{x^2 y}^1 & X_y^1 \\ X_{x^1}^2 & X_{x^2 y}^2 & X_y^2 \\ Y_{x^1} & Y_{x^2 y} & Y_y \end{array} \right| \right\} + y_{x^2} y_{x^2} \left\{ - \left| \begin{array}{ccc} X_{x^1}^1 & X_{x^1 y}^1 & X_y^1 \\ X_{x^1}^2 & X_{x^1 y}^2 & X_y^2 \\ Y_{x^1} & Y_{x^1 y} & Y_y \end{array} \right| \right\} + \\
& + y_{x^1} y_{x^1} y_{x^2} \cdot \left\{ - \left| \begin{array}{ccc} X_{yy}^1 & X_{x^2}^1 & X_y^1 \\ X_{yy}^2 & X_{x^2}^2 & X_y^2 \\ Y_{yy} & Y_{x^2} & Y_y \end{array} \right| \right\} + \\
& + y_{x^1} y_{x^2} y_{x^2} \cdot \left\{ - \left| \begin{array}{ccc} X_{x^1}^1 & X_{yy}^1 & X_y^1 \\ X_{x^1}^2 & X_{yy}^2 & X_y^2 \\ Y_{x^1} & Y_{yy} & Y_y \end{array} \right| \right\}.
\end{aligned}$$

Using the third equation (2.15), we also deduce, *without any further computation*, that the third equation which would be obtained by developing (2.9) in length, is:

$$\begin{aligned}
(2.23) \quad 0 & = y_{x^2 x^2} \cdot \left| \begin{array}{ccc} X_{x^1}^1 & X_{x^2}^1 & X_y^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_y^2 \\ Y_{x^1} & Y_{x^2} & Y_y \end{array} \right| + \left| \begin{array}{ccc} X_{x^1}^1 & X_{x^2}^1 & X_{x^2 x^2}^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_{x^2 x^2}^2 \\ Y_{x^1} & Y_{x^2} & Y_{x^2 x^2} \end{array} \right| + \\
& + y_{x^1} \cdot \left\{ - \left| \begin{array}{ccc} X_{x^2 x^2}^1 & X_{x^2}^1 & X_y^1 \\ X_{x^1 x^2}^2 & X_{x^2}^2 & X_y^2 \\ Y_{x^2 x^2} & Y_{x^2} & Y_y \end{array} \right| \right\} + \\
& + y_{x^2} \cdot \left\{ 2 \left| \begin{array}{ccc} X_{x^1}^1 & X_{x^2}^1 & X_{x^2 y}^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_{x^2 y}^2 \\ Y_{x^1} & Y_{x^2} & Y_{x^2 y} \end{array} \right| - \left| \begin{array}{ccc} X_{x^1}^1 & X_{x^2 x^2}^1 & X_y^1 \\ X_{x^1}^2 & X_{x^2 x^2}^2 & X_y^2 \\ Y_{x^1} & Y_{x^2 x^2} & Y_y \end{array} \right| \right\} + \\
& + y_{x^1} y_{x^2} \cdot \left\{ -2 \left| \begin{array}{ccc} X_{x^2 y}^1 & X_{x^2}^1 & X_y^1 \\ X_{x^2 y}^2 & X_{x^2}^2 & X_y^2 \\ Y_{x^2 y} & Y_{x^2} & Y_y \end{array} \right| \right\} + \\
& + y_{x^2} y_{x^2} \cdot \left\{ \left| \begin{array}{ccc} X_{x^1}^1 & X_{x^2}^1 & X_{yy}^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_{yy}^2 \\ Y_{x^1} & Y_{x^2} & Y_{yy} \end{array} \right| - 2 \left| \begin{array}{ccc} X_{x^1}^1 & X_{x^2 y}^1 & X_y^1 \\ X_{x^1}^2 & X_{x^2 y}^2 & X_y^2 \\ Y_{x^1} & Y_{x^2 y} & Y_y \end{array} \right| \right\} + \\
& + y_{x^1} y_{x^2} y_{x^2} \cdot \left\{ - \left| \begin{array}{ccc} X_{yy}^1 & X_{x^2}^1 & X_y^1 \\ X_{yy}^2 & X_{x^2}^2 & X_y^2 \\ Y_{yy} & Y_{x^2} & Y_y \end{array} \right| \right\} + \\
& + y_{x^2} y_{x^2} y_{x^2} \cdot \left\{ - \left| \begin{array}{ccc} X_{x^1}^1 & X_{yy}^1 & X_y^1 \\ X_{x^1}^2 & X_{yy}^2 & X_y^2 \\ Y_{x^1} & Y_{yy} & Y_y \end{array} \right| \right\}.
\end{aligned}$$

2.24. Appropriate formalism. To describe the combinatorics underlying formulas (2.10), (2.22) and (2.23), as in [Me2004], let us introduce the following notation for

the Jacobian determinant:

$$(2.25) \quad \Delta(x^1|x^2|y) := \begin{vmatrix} X_{x^1}^1 & X_{x^2}^1 & X_y^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_y^2 \\ Y_{x^1} & Y_{x^2} & Y_y \end{vmatrix}.$$

Here, in the notation $\Delta(x^1|x^2|y)$, the three spaces between the two vertical lines $|$ refer to the three columns of the Jacobian determinant, and the terms x^1 , x^2 , y in $(x^1|x^2|y)$ designate the partial derivatives appearing in each column. Accordingly, in the following two examples of *modified Jacobian determinants*:

$$(2.26) \quad \left\{ \begin{array}{l} \Delta(\underline{x^1x^2}|x^2|y) := \begin{vmatrix} X_{\underline{x^1x^2}}^1 & X_{x^2}^1 & X_y^1 \\ X_{\underline{x^1x^2}}^2 & X_{x^2}^2 & X_y^2 \\ Y_{\underline{x^1x^2}} & Y_{x^2} & Y_y \end{vmatrix} \\ \Delta(x^1|x^2|\underline{x^1y}) := \begin{vmatrix} X_{x^1}^1 & X_{x^2}^1 & X_{\underline{x^1y}}^1 \\ X_{x^1}^2 & X_{x^2}^2 & X_{\underline{x^1y}}^2 \\ Y_{x^1} & Y_{x^2} & Y_{\underline{x^1y}} \end{vmatrix}, \end{array} \right. \quad \text{and}$$

we simply mean which column of first order derivatives is replaced by a column of second order derivatives in the original Jacobian determinant.

As there are 6 possible second order derivatives $(\cdot)_{x^1x^1}$, $(\cdot)_{x^1x^2}$, $(\cdot)_{x^1xy}$, $(\cdot)_{x^2x^2}$, $(\cdot)_{x^2y}$ and $(\cdot)_{yy}$ together with 3 columns, we obtain $3 \times 6 = 18$ possible modified Jacobian determinants:

$$(2.27) \quad \left\{ \begin{array}{lll} \Delta(x^1x^1|x^2|y) & \Delta(x^1|x^1x^1|y) & \Delta(x^1|x^2|x^1x^1) \\ \Delta(x^1x^2|x^2|y) & \Delta(x^1|x^1x^2|y) & \Delta(x^1|x^2|x^1x^2) \\ \Delta(x^1y|x^2|y) & \Delta(x^1|x^1y|y) & \Delta(x^1|x^2|x^1y) \\ \Delta(x^2x^2|x^2|y) & \Delta(x^1|x^2x^2|y) & \Delta(x^1|x^2|x^2x^2) \\ \Delta(x^2y|x^2|y) & \Delta(x^1|x^2y|y) & \Delta(x^1|x^2|x^2y) \\ \Delta(yy|x^2|y) & \Delta(x^1|yy|y) & \Delta(x^1|x^2|yy). \end{array} \right.$$

Next, we observe that if we want to solve with respect to $y_{x^1x^1}$ in (2.10), with respect to $y_{x^1x^2}$ in (2.22) and with respect to $y_{x^2x^2}$ in (2.23), we have to divide by the Jacobian determinant $\Delta(x^1|x^2|y)$. Consequently, we introduce 18 new *square functions* as follows:

$$(2.28) \quad \left\{ \begin{array}{lll} \square_{x^1x^1}^1 := \frac{\Delta(x^1x^1|x^2|y)}{\Delta(x^1|x^2|y)} & \square_{x^1x^2}^1 := \frac{\Delta(x^1x^2|x^2|y)}{\Delta(x^1|x^2|y)} & \square_{x^1y}^1 := \frac{\Delta(x^1y|x^2|y)}{\Delta(x^1|x^2|y)} \\ \square_{x^2x^2}^1 := \frac{\Delta(x^2x^2|x^2|y)}{\Delta(x^1|x^2|y)} & \square_{x^2y}^1 := \frac{\Delta(x^2y|x^2|y)}{\Delta(x^1|x^2|y)} & \square_{yy}^1 := \frac{\Delta(yy|x^2|y)}{\Delta(x^1|x^2|y)} \\ \square_{x^1x^1}^2 := \frac{\Delta(x^1|x^1x^1|y)}{\Delta(x^1|x^2|y)} & \square_{x^1x^2}^2 := \frac{\Delta(x^1|x^1x^2|y)}{\Delta(x^1|x^2|y)} & \square_{x^1y}^2 := \frac{\Delta(x^1|x^1y|y)}{\Delta(x^1|x^2|y)} \\ \square_{x^2x^2}^2 := \frac{\Delta(x^1|x^2x^2|y)}{\Delta(x^1|x^2|y)} & \square_{x^2y}^2 := \frac{\Delta(x^1|x^2y|y)}{\Delta(x^1|x^2|y)} & \square_{yy}^2 := \frac{\Delta(x^1|yy|y)}{\Delta(x^1|x^2|y)} \\ \square_{x^1x^1}^3 := \frac{\Delta(x^1|x^2|x^1x^1)}{\Delta(x^1|x^2|y)} & \square_{x^1x^2}^3 := \frac{\Delta(x^1|x^2|x^1x^2)}{\Delta(x^1|x^2|y)} & \square_{x^1y}^3 := \frac{\Delta(x^1|x^2|x^1y)}{\Delta(x^1|x^2|y)} \\ \square_{x^2x^2}^3 := \frac{\Delta(x^1|x^2|x^2x^2)}{\Delta(x^1|x^2|y)} & \square_{x^2y}^3 := \frac{\Delta(x^1|x^2|x^2y)}{\Delta(x^1|x^2|y)} & \square_{yy}^3 := \frac{\Delta(x^1|x^2|yy)}{\Delta(x^1|x^2|y)}. \end{array} \right.$$

Thanks to these notations, we can rewrite the three equations (2.10), (2.22) and (2.23) in a more compact style.

Lemma 2.29. *A completely integrable system of three second order partial differential equations*

$$(2.30) \quad \begin{cases} y_{x^1 x^1}(x) = F^{1,1}(x^1, x^2, y(x), y_{x^1}(x), y_{x^2}(x)), \\ y_{x^1 x^2}(x) = F^{1,2}(x^1, x^2, y(x), y_{x^1}(x), y_{x^2}(x)), \\ y_{x^2 x^2}(x) = F^{2,2}(x^1, x^2, y(x), y_{x^1}(x), y_{x^2}(x)), \end{cases}$$

is equivalent to the simplest system $Y_{X^1 X^1} = 0$, $Y_{X^1 X^2} = 0$, $Y_{X^2 X^2} = 0$, if and only if there exist local \mathbb{K} -analytic functions X^1 , X^2 , Y such that it may be written under the specific form:

$$(2.31) \quad \begin{cases} y_{x^1 x^1} = -\square_{x^1 x^1}^3 + y_{x^1} \cdot \left(-2\square_{x^1 y}^3 + \square_{x^1 x^1}^1\right) + y_{x^2} \cdot \left(\square_{x^1 x^1}^2\right) + \\ \quad + y_{x^1} y_{x^1} \cdot \left(-\square_{yy}^3 + 2\square_{x^1 y}^1\right) + y_{x^1} y_{x^2} \cdot \left(2\square_{x^1 y}^2\right) + \\ \quad + y_{x^1} y_{x^1} y_{x^1} \cdot \left(\square_{yy}^1\right) + y_{x^1} y_{x^1} y_{x^2} \cdot \left(\square_{yy}^2\right), \\ y_{x^1 x^2} = -\square_{x^1 x^2}^3 + y_{x^1} \cdot \left(-\square_{x^2 y}^3 + \square_{x^1 x^2}^1\right) + y_{x^2} \cdot \left(-\square_{x^1 y}^3 + \square_{x^1 x^2}^2\right) + \\ \quad + y_{x^1} y_{x^1} \cdot \left(\square_{x^2 y}^1\right) + y_{x^1} y_{x^2} \cdot \left(-\square_{yy}^3 + \square_{x^1 y}^1 + \square_{x^2 y}^2\right) + \\ \quad + y_{x^2} y_{x^2} \cdot \left(\square_{x^1 y}^2\right) + y_{x^1} y_{x^1} y_{x^2} \cdot \left(\square_{yy}^1\right) + y_{x^1} y_{x^2} y_{x^2} \cdot \left(\square_{yy}^2\right), \\ y_{x^2 x^2} = -\square_{x^2 x^2}^3 + y_{x^1} \cdot \left(\square_{x^2 x^2}^1\right) + y_{x^2} \cdot \left(-2\square_{x^2 y}^3 + \square_{x^2 x^2}^2\right) + \\ \quad + y_{x^1} y_{x^2} \cdot \left(2\square_{x^2 y}^1\right) + y_{x^2} y_{x^2} \cdot \left(-\square_{yy}^3 + 2\square_{x^2 y}^2\right) + \\ \quad + y_{x^1} y_{x^2} y_{x^2} \cdot \left(\square_{yy}^1\right) + y_{x^2} y_{x^2} y_{x^2} \cdot \left(\square_{yy}^2\right). \end{cases}$$

2.32. General formulas. The formal dictionary between the original determinantal formulas (2.10), (2.22), (2.23), between the coefficients (2.15) of the second order prolongation of a vector field and between the new square formulas (2.31) above is evident. Consequently, *without any computation*, just by translating the family of formulas (2.14), we may deduce the exact formulation of the desired generalization of Lemma 2.29 above.

Lemma 2.33. *A completely integrable system of second order partial differential equations of the form*

$$(2.34) \quad y_{x^{j_1} x^{j_2}}(x) = F^{j_1, j_2}(x, y(x), y_{x^1}(x), \dots, y_{x^n}(x)), \quad j_1, j_2 = 1, \dots, n,$$

is equivalent to the simplest system $Y_{X^{j_1} X^{j_2}} = 0$, $j_1, j_2 = 1, \dots, n$, if and only if there exist local \mathbb{K} -analytic functions X^l , Y such that it may be written under the specific form:

$$(2.35) \quad \begin{cases} y_{x^{j_1} x^{j_2}} = -\square_{x^{j_1} x^{j_2}}^{n+1} + \sum_{k_1=1}^n y_{x^{k_1}} \cdot \left\{ \left(\square_{x^{j_1} x^{j_2}}^{k_1} - \delta_{j_1}^{k_1} \square_{x^{j_2} y}^{n+1} - \delta_{j_2}^{k_1} \square_{x^{j_1} y}^{n+1} \right) + \right. \\ \quad + y_{x^{j_1}} \cdot \left(\square_{x^{j_2} y}^{k_1} - \frac{1}{2} \delta_{j_2}^{k_1} \square_{yy}^{n+1} \right) + y_{x^{j_2}} \cdot \left(\square_{x^{j_1} y}^{k_1} - \frac{1}{2} \delta_{j_1}^{k_1} \square_{yy}^{n+1} \right) + \\ \quad \left. + y_{x^{j_1}} y_{x^{j_2}} \cdot \left(\square_{yy}^{k_1} \right) \right\}. \end{cases}$$

Of course, to define the square functions in the context of $n \geq 2$ independent variables (x^1, x^2, \dots, x^n) , we introduce the Jacobian determinant

$$(2.36) \quad \Delta(x^1 | x^2 | \dots | x^n | y) := \begin{vmatrix} X_{x^1}^1 & \cdots & X_{x^n}^1 & X_y^1 \\ \vdots & \cdots & \vdots & \vdots \\ X_{x^1}^n & \cdots & X_{x^n}^n & X_y^n \\ Y_{x^1} & \cdots & Y_{x^n} & Y_y \end{vmatrix},$$

together with its modifications

$$(2.37) \quad \Delta (x^1 | \dots |^{k_1} x^{j_1} x^{j_2} | \dots | y),$$

in which the k_1 -th column of partial first order derivatives $|^{k_1} x^{k_1}|$ is replaced by the column $|^{k_1} x^{j_1} x^{j_2}|$ of partial derivatives. Here, the indices k_1, j_1, j_2 satisfy $1 \leq k_1, j_1, j_2 \leq n+1$, with the convention that we adopt the notational equivalence

$$(2.38) \quad \boxed{x^{n+1} \equiv y}.$$

This convention will be convenient to write some of our general formulas in the sequel.

As we promised to only summarize the proof of Theorem 1.7 in this paper, we will not develop the proof of Lemma 2.33: it is similar to the proof of Lemma 3.32 in [Me2004].

§3. FIRST AND SECOND AUXILIARY SYSTEM

3.1. Functions $G_{j_1, j_2}, H_{j_1, j_2}^{k_1}, L_{j_1}^{k_1}$ and M^{k_1} . To discover the four families of functions appearing in the statement of Theorem 1.7, by comparing (2.35) and (1.10), it suffices (of course) to set:

$$(3.2) \quad \begin{cases} G_{j_1, j_2} := -\square_{x^{j_1} x^{j_2}}^{n+1}, \\ H_{j_1, j_2}^{k_1} := \square_{x^{j_1} x^{j_2}}^{k_1} - \delta_{j_1}^{k_1} \square_{x^{j_2} y}^{n+1} - \delta_{j_2}^{k_1} \square_{x^{j_1} y}^{n+1}, \\ L_{j_1}^{k_1} := 2 \square_{x^{j_1} y}^{k_1} - \delta_{j_1}^{k_1} \square_{yy}^{n+1}, \\ M^{k_1} := \square_{yy}^{k_1}. \end{cases}$$

Consequently, we have shown the “only if” part of Theorem 1.7, which is the easiest implication.

To establish the “if” part, by far the most difficult implication, the very main lemma can be stated as follows.

Lemma 3.3. *The partial differential relations (I’), (II’), (III’) and (IV’) which express in length the compatibility conditions (1.11) are necessary and sufficient for the existence of functions X^l, Y of (x^{l_1}, y) satisfying the second order nonlinear system of partial differential equations (3.2) above.*

Indeed, the collection of equations (3.2) is a system of partial differential equations with unknowns X^l, Y , by virtue of the definition of the square functions.

3.4. First auxiliary system. To proceed further, we observe that there are $(m+1)$ more square functions than functions $G_{j_1, j_2}, H_{j_1, j_2}^{k_1}, L_{j_1}^{k_1}$ and M^{k_1} . Indeed, a simple counting yields:

$$(3.5) \quad \begin{cases} \#\{\square_{x^{j_1} x^{j_2}}^{k_1}\} = \frac{n^2(n+1)}{2}, & \#\{\square_{x^{j_1} y}^{k_1}\} = n^2, \\ \#\{\square_{yy}^{k_1}\} = n, & \#\{\square_{x^{j_1} x^{j_2}}^{n+1}\} = \frac{n(n+1)}{2}, \\ \#\{\square_{x^{j_1} y}^{n+1}\} = n, & \#\{\square_{yy}^{n+1}\} = 1, \end{cases}$$

whereas

$$(3.6) \quad \begin{cases} \#\{G^{j_1, j_2}\} = \frac{n(n+1)}{2}, & \#\{H_{j_1, j_2}^{k_1}\} = \frac{n^2(n+1)}{2}, \\ \#\{L_{j_1}^{k_1}\} = n^2, & \#\{M^{k_1}\} = n. \end{cases}$$

Here, the indices j_1, j_2, k_1 satisfy $1 \leq j_1, j_2, k_1 \leq n$. Similarly as in [Me2004], to transform the system (3.2) in a true complete system, let us introduce functions $\Pi_{j_1, j_2}^{k_1}$

(x^{j_1}, y) , where $1 \leq j_1, j_2, k_1 \leq n+1$, which satisfy the symmetry $\Pi_{j_1, j_2}^{k_1} = \Pi_{j_1, j_1}^{k_1}$, and let us introduce the following *first auxiliary system*:

$$(3.7) \quad \begin{cases} \square_{x^{j_1} x^{j_2}}^{k_1} = \Pi_{j_1, j_2}^{k_1}, & \square_{x^{j_1} y}^{k_1} = \Pi_{j_1, n+1}^{k_1}, & \square_{yy}^{k_1} = \Pi_{n+1, n+1}^{k_1}, \\ \square_{x^{j_1} x^{j_2}}^{n+1} = \Pi_{j_1, j_2}^{n+1}, & \square_{x^{j_1} y}^{n+1} = \Pi_{j_1, n+1}^{n+1}, & \square_{yy}^{n+1} = \Pi_{n+1, n+1}^{n+1}. \end{cases}$$

It is complete. The necessary and sufficient conditions for the existence of solutions X^l , Y follow by cross differentiations.

Lemma 3.8. *For all $j_1, j_2, j_3, k_1 = 1, 2, \dots, n+1$, we have the cross differentiation relations*

$$(3.9) \quad \left(\square_{x^{j_1} x^{j_2}}^{k_1} \right)_{x^{j_3}} - \left(\square_{x^{j_1} x^{j_3}}^{k_1} \right)_{x^{j_2}} = - \sum_{k_2=1}^{n+1} \square_{x^{j_1} x^{j_2}}^{k_2} \square_{x^{j_3} x^{k_2}}^{k_1} + \sum_{k_2=1}^{n+1} \square_{x^{j_1} x^{j_3}}^{k_2} \square_{x^{j_2} x^{k_2}}^{k_1}.$$

The proof of this lemma is exactly the same as the proof of Lemma 3.40 in [Me2004].

As a direct consequence, we deduce that a necessary and sufficient condition for the existence of solutions $\Pi_{j_1, j_2}^{k_1}$ to the first auxiliary system is that they satisfy the following compatibility partial differential relations:

$$(3.10) \quad \frac{\partial \Pi_{j_1, j_2}^{k_1}}{\partial x^{j_3}} - \frac{\partial \Pi_{j_1, j_3}^{k_1}}{\partial x^{j_2}} = - \sum_{k_2=1}^{n+1} \Pi_{j_1, j_2}^{k_2} \cdot \Pi_{j_3, k_2}^{k_1} + \sum_{k_2=1}^{n+1} \Pi_{j_1, j_3}^{k_2} \cdot \Pi_{j_2, k_2}^{k_1},$$

for all $j_1, j_2, j_3, k_1 = 1, \dots, n+1$.

We shall have to specify this system in length according to the splitting $\{1, 2, \dots, n\}$ and $\{n+1\}$ of the indices of coordinates. We obtain six families of equations equivalent

to (3.10) just above:

(3.11)

$$\left. \begin{aligned}
 (\Pi_{j_1, j_2}^{n+1})_{xj_3} - (\Pi_{j_1, j_3}^{n+1})_{xj_2} &= - \sum_{k_2=1}^n \Pi_{j_1, j_2}^{k_2} \Pi_{j_3, k_2}^{n+1} - \Pi_{j_1, j_2}^{n+1} \Pi_{j_3, n+1}^{n+1} + \\
 &\quad + \sum_{k_2=1}^n \Pi_{j_1, j_3}^{k_2} \Pi_{j_2, k_2}^{n+1} + \Pi_{j_1, j_3}^{n+1} \Pi_{j_2, n+1}^{n+1}, \\
 (\Pi_{j_1, j_2}^{n+1})_y - (\Pi_{j_1, n+1}^{n+1})_{xj_2} &= - \sum_{k_2=1}^n \Pi_{j_1, j_2}^{k_2} \Pi_{n+1, k_2}^{n+1} - \Pi_{j_1, j_2}^{n+1} \Pi_{n+1, n+1}^{n+1} + \\
 &\quad + \sum_{k_2=1}^n \Pi_{j_1, n+1}^{k_2} \Pi_{j_2, k_2}^{n+1} + \Pi_{j_1, n+1}^{n+1} \Pi_{j_2, n+1}^{n+1}, \\
 (\Pi_{j_1, n+1}^{n+1})_y - (\Pi_{n+1, n+1}^{n+1})_{xj_1} &= - \sum_{k_2=1}^n \Pi_{j_1, n+1}^{k_2} \Pi_{n+1, k_2}^{n+1} - \underbrace{\Pi_{j_1, n+1}^{n+1} \Pi_{n+1, n+1}^{n+1}}_{\circ} + \\
 &\quad + \sum_{k_2=1}^n \Pi_{n+1, n+1}^{k_2} \Pi_{j_1, k_2}^{n+1} + \underbrace{\Pi_{n+1, n+1}^{n+1} \Pi_{j_1, n+1}^{n+1}}_{\circ}, \\
 (\Pi_{j_1, j_2}^{k_1})_{xj_3} - (\Pi_{j_1, j_3}^{k_1})_{xj_2} &= - \sum_{k_2=1}^n \Pi_{j_1, j_2}^{k_2} \Pi_{j_3, k_2}^{k_1} - \Pi_{j_1, j_2}^{n+1} \Pi_{j_3, n+1}^{k_1} + \\
 &\quad + \sum_{k_2=1}^n \Pi_{j_1, j_3}^{k_2} \Pi_{j_2, k_2}^{k_1} + \Pi_{j_1, j_3}^{n+1} \Pi_{j_2, n+1}^{k_1}, \\
 (\Pi_{j_1, j_2}^{k_1})_y - (\Pi_{j_1, n+1}^{k_1})_{xj_2} &= - \sum_{k_2=1}^n \Pi_{j_1, j_2}^{k_2} \Pi_{n+1, k_2}^{k_1} - \Pi_{j_1, j_2}^{n+1} \Pi_{n+1, n+1}^{k_1} + \\
 &\quad + \sum_{k_2=1}^n \Pi_{j_1, n+1}^{k_2} \Pi_{j_2, k_2}^{k_1} + \Pi_{j_1, n+1}^{n+1} \Pi_{j_2, n+1}^{k_1}, \\
 (\Pi_{j_1, n+1}^{k_1})_y - (\Pi_{n+1, n+1}^{k_1})_{xj_1} &= - \sum_{k_2=1}^n \Pi_{j_1, n+1}^{k_2} \Pi_{n+1, k_2}^{k_1} - \Pi_{j_1, n+1}^{n+1} \Pi_{n+1, n+1}^{k_1} + \\
 &\quad + \sum_{k_2=1}^n \Pi_{n+1, n+1}^{k_2} \Pi_{j_1, k_2}^{k_1} + \Pi_{n+1, n+1}^{n+1} \Pi_{j_1, n+1}^{k_1}.
 \end{aligned} \right\}$$

where the indices j_1, j_2, j_3, k_1 vary in the set $\{1, 2, 1, \dots, n\}$.

3.12. Principal unknowns. As there are $(m + 1)$ more square (or Pi) functions than the functions $G_{j_1, j_2}^{k_1}, H_{j_1, j_2}^{k_1}, L_{j_1}^{k_1}$ and M^{k_1} , we cannot invert directly the linear system (3.2). To quasi-invert it, we choose the $(m + 1)$ specific square functions

$$(3.13) \quad \Theta^1 := \square_{x^1 x^1}^1, \quad \Theta^2 := \square_{x^2 x^2}^2, \dots, \Theta^{n+1} := \square_{x^{n+1} x^{n+1}}^{n+1},$$

calling them *principal unknowns*, and we get the quasi-inversion:

$$(3.14) \quad \left\{ \begin{array}{l} \Pi_{j_1, j_2}^{k_1} = \square_{x^{j_1} x^{j_2}}^{k_1} = H_{j_1, j_2}^{k_1} - \frac{1}{2} \delta_{j_1}^{k_1} H_{j_2, j_2}^{j_2} - \frac{1}{2} \delta_{j_2}^{k_1} H_{j_1, j_1}^{j_1} + \frac{1}{2} \delta_{j_1}^{k_1} \Theta^{j_2} + \frac{1}{2} \delta_{j_2}^{k_1} \Theta^{j_1}, \\ \Pi_{j_1, n+1}^{k_1} = \square_{x^{j_1} y}^{k_1} = \frac{1}{2} L_{j_1}^{k_1} + \frac{1}{2} \delta_{j_1}^{k_1} \Theta^{n+1}, \\ \Pi_{n+1, n+1}^{k_1} = \square_{yy}^{k_1} = M^{k_1}, \\ \Pi_{j_1, j_2}^{n+1} = \square_{x^{j_1} x^{j_2}}^{n+1} = -G_{j_1, j_2}, \\ \Pi_{j_1, n+1}^{n+1} = \square_{x^{j_1} y}^{n+1} = -\frac{1}{2} H_{j_1, j_1}^{j_1} + \frac{1}{2} \Theta^{j_1}. \end{array} \right.$$

3.15. Second auxiliary system. Replacing the five families of functions $\Pi_{j_1, j_2}^{k_1}$, $\Pi_{j_1, n+1}^{k_1}$, $\Pi_{n+1, n+1}^{k_1}$, Π_{j_1, j_2}^{n+1} , $\Pi_{j_1, n+1}^{n+1}$ by their values obtained in (3.14) just above together with the principal unknowns

$$(3.16) \quad \Pi_{j_1, j_1}^{j_1} = \Theta^{j_1}, \quad \Pi_{n+1, n+1}^{n+1} = \Theta^{n+1},$$

in the six equations (3.11)₁, (3.11)₂, (3.11)₃, (3.11)₄, (3.11)₅ and (3.11)₆, after hard computations that we will not reproduce here, we obtain six families of equations. From now on, we abbreviate every sum $\sum_{k=1}^n$ as \sum_{k_1} .

Firstly:

$$(3.17) \quad 0 = \underline{G_{j_1, j_2, x^{j_3}} - G_{j_1, j_3, x^{j_2}}} + \sum_{k_1} G_{j_3, k_1} H_{j_1, j_2}^{k_1} - \sum_{k_1} G_{j_2, k_1} H_{j_1, j_3}^{k_1}.$$

This is (I') of Theorem 1.7. Just above and below, we plainly underline the monomials involving a first order derivative. Secondly:

$$(3.18) \quad \left\{ \begin{array}{l} \Theta_{x^{j_2}}^{j_1} = \underline{-2 G_{j_1, j_2, y} + H_{j_1, j_1, x^{j_2}}^{j_1}} + \\ + \sum_{k_1} G_{j_2, k_1} L_{j_1}^{k_1} + \frac{1}{2} H_{j_1, j_1}^{j_1} H_{j_2, j_2}^{j_2} - \sum_{k_1} H_{j_1, j_2}^{k_1} H_{k_1, k_1}^{k_1} - \\ - G_{j_1, j_2} \Theta^{n+1} - \frac{1}{2} H_{j_1, j_1}^{j_1} \Theta^{j_2} - \frac{1}{2} H_{j_2, j_2}^{j_2} \Theta^{j_1} + \sum_{k_1} H_{j_1, j_2}^{k_1} \Theta^{k_1} + \\ + \frac{1}{2} \Theta^{j_1} \Theta^{j_2}. \end{array} \right.$$

Thirdly:

$$(3.19) \quad \left\{ \begin{array}{l} -\Theta_{x^{j_1}}^{n+1} + \frac{1}{2} \Theta_y^{j_1} = \underline{\frac{1}{2} H_{j_1, j_1, y}^{j_1}} - \\ - \sum_{k_1} G_{j_1, k_1} M^{k_1} + \frac{1}{4} \sum_{k_1} H_{k_1, k_1}^{k_1} L_{j_1}^{k_1} + \\ + \frac{1}{4} H_{j_1, j_1}^{j_1} \Theta^{n+1} - \frac{1}{4} \sum_{k_1} L_{j_1}^{k_1} \Theta^{k_1} - \frac{1}{4} \Theta^{j_1} \Theta^{n+1}. \end{array} \right.$$

Fourthly:

(3.20)

$$\left\{ \begin{aligned}
 & \frac{1}{2} \delta_{j_1}^{k_1} \Theta_{x^{j_3}}^{j_2} - \frac{1}{2} \delta_{j_1}^{k_1} \Theta_{x^{j_2}}^{j_3} + \frac{1}{2} \delta_{j_2}^{k_1} \Theta_{x^{j_3}}^{j_1} - \frac{1}{2} \delta_{j_3}^{k_1} \Theta_{x^{j_2}}^{j_1} = \\
 & \quad = -\frac{H_{j_1, j_2, x^{j_3}}^{k_1}}{H_{j_1, j_3, x^{j_2}}^{k_1}} + \frac{H_{j_1, j_3, x^{j_2}}^{k_1}}{H_{j_3, j_3, x^{j_2}}^{j_3}} + \frac{1}{2} \delta_{j_1}^{k_1} H_{j_2, j_2, x^{j_3}}^{j_2} - \\
 & \quad - \frac{\frac{1}{2} \delta_{j_3}^{k_1} H_{j_1, j_1, x^{j_2}}^{j_1}}{\frac{1}{2} \delta_{j_2}^{k_1} H_{j_1, j_1, x^{j_3}}^{j_1}} - \\
 & \quad - \frac{1}{2} G_{j_1, j_2} L_{j_3}^{k_1} + \frac{1}{2} G_{j_1, j_3} L_{j_2}^{k_1} - \frac{1}{4} \delta_{j_3}^{k_1} H_{j_1, j_1}^{j_1} H_{j_2, j_2}^{j_2} + \frac{1}{4} \delta_{j_2}^{k_1} H_{j_1, j_1}^{j_1} H_{j_3, j_3}^{j_3} - \\
 & \quad - \sum_{k_2} H_{j_1, j_2}^{k_2} H_{j_3, k_2}^{k_1} + \sum_{k_2} H_{j_1, j_3}^{k_2} H_{j_2, k_2}^{k_1} - \frac{1}{2} \delta_{j_2}^{k_1} H_{j_1, j_3}^{k_2} H_{k_2, k_2}^{k_2} + \frac{1}{2} \delta_{j_3}^{k_1} H_{j_1, j_2}^{k_2} H_{k_2, k_2}^{k_2} - \\
 & \quad - \frac{1}{2} \delta_{j_2}^{k_1} G_{j_1, j_3} \Theta^{n+1} + \frac{1}{2} \delta_{j_3}^{k_1} G_{j_1, j_2} \Theta^{n+1} - \\
 & \quad - \frac{1}{4} \delta_{j_2}^{k_1} H_{j_1, j_1}^{j_1} \Theta^{j_3} + \frac{1}{4} \delta_{j_3}^{k_1} H_{j_1, j_1}^{j_1} \Theta^{j_2} - \frac{1}{4} \delta_{j_2}^{k_1} H_{j_3, j_3}^{j_3} \Theta^{j_1} + \frac{1}{4} \delta_{j_3}^{k_1} H_{j_2, j_2}^{j_2} \Theta^{j_1} - \\
 & \quad - \frac{1}{2} \delta_{j_3}^{k_1} \sum_{k_1} H_{j_1, j_2}^{k_2} \Theta^{k_2} + \frac{1}{2} \delta_{j_2}^{k_1} \sum_{k_1} H_{j_1, j_3}^{k_2} \Theta^{k_2} - \\
 & \quad - \frac{1}{4} \delta_{j_3}^{k_1} \Theta^{j_1} \Theta^{j_2} + \frac{1}{4} \delta_{j_2}^{k_1} \Theta^{j_1} \Theta^{j_3}.
 \end{aligned} \right.$$

Fifthly:

(3.21)

$$\left\{ \begin{aligned}
 & \frac{1}{2} \delta_{j_1}^{k_1} \Theta_y^{j_2} + \frac{1}{2} \delta_{j_2}^{k_1} \Theta_y^{j_1} - \frac{1}{2} \delta_{j_1}^{k_1} \Theta_{x^{j_2}}^{n+1} = \\
 & \quad = G_{j_1, j_2} M^{k_1} + \frac{1}{2} \sum_{k_2} H_{j_1, k_2}^{k_1} L_{j_1}^{k_2} - \frac{1}{2} \sum_{k_2} H_{j_1, j_2}^{k_2} L_{k_2}^{k_1} - \frac{1}{4} \delta_{j_2}^{k_1} \sum_{k_2} H_{k_2, k_2}^{k_2} L_{j_1}^{k_2} - \\
 & \quad - \frac{1}{4} \delta_{j_2}^{k_1} H_{j_1, j_1}^{j_1} \Theta^{n+1} + \frac{1}{4} \delta_{j_2}^{k_1} \sum_{k_2} L_{j_1}^{k_2} \Theta^{k_2} + \frac{1}{4} \delta_{j_2}^{k_1} \Theta^{j_1} \Theta^{n+1}.
 \end{aligned} \right.$$

Sixthly:

$$(3.22) \quad \left\{ \begin{aligned}
 & \delta_{j_1}^{k_1} \Theta_y^{n+1} = -\frac{L_{j_1, y}^{k_1}}{x^{j_1}} + 2 M_{x^{j_1}}^{k_1} + \\
 & \quad + 2 \sum_{k_2} H_{j_1, k_2}^{k_1} M^{k_2} - \delta_{j_1}^{k_1} \sum_{k_2} H_{k_2, k_2}^{k_2} M^{k_2} - \frac{1}{2} \sum_{k_2} L_{j_1}^{k_2} L_{k_2}^{k_1} + \\
 & \quad + \delta_{j_1}^{k_1} \sum_{k_2} M^{k_2} \Theta^{k_2} + \frac{1}{2} \delta_{j_1}^{k_1} \Theta^{n+1} \Theta^{n+1}.
 \end{aligned} \right.$$

3.23. Solving $\Theta_{x^{j_2}}^{j_1}$, $\Theta_y^{j_1}$, $\Theta_{x^{j_1}}^{n+1}$ and Θ_y^{n+1} . From the six families of equations (3.17), (3.18), (3.19), (3.20), (3.21) and (3.22), we can solve $\Theta_{x^{j_2}}^{j_1}$, $\Theta_y^{j_1}$, $\Theta_{x^{j_1}}^{n+1}$ and Θ_y^{n+1} . Not mentioning the (hard) intermediate computations, we obtain firstly:

(3.24)

$$\left\{ \begin{aligned}
 & \Theta_{x^{j_2}}^{j_1} = -\frac{2 G_{j_1, j_2, y} + H_{j_1, j_1, x^{j_2}}^{j_1}}{L_{j_1}^{j_1}} + \sum_l G_{j_2, l} L_{j_1}^l + \frac{1}{2} H_{j_1, j_1}^{j_1} H_{j_2, j_2}^{j_2} - \sum_l H_{j_1, j_2}^l H_{l, l}^l - \\
 & \quad - G_{j_1, j_2} \Theta^{n+1} - \frac{1}{2} H_{j_1, j_1} \Theta^{j_1} - \frac{1}{2} H_{j_2, j_2}^{j_2} \Theta^{j_1} + \sum_l H_{j_1, j_2}^l \Theta^l + \frac{1}{2} \Theta^{j_1} \Theta^{j_2}.
 \end{aligned} \right.$$

Secondly:

(3.25)

$$\left\{ \begin{aligned} \Theta_y^{j_1} &= -\frac{1}{3} \frac{H_{j_1, j_1, y}^{j_1}}{L_{j_1, x^{j_1}}^{j_1}} + \frac{2}{3} \frac{L_{j_1, x^{j_1}}^{j_1}}{L_{j_1, x^{j_1}}^{j_1}} + \frac{4}{3} G_{j_1, j_1} M^{j_1} + \frac{2}{3} \sum_l G_{j_1, l} M^l - \frac{1}{2} \sum_l H_{l, l}^l L_{j_1}^l + \\ &+ \frac{2}{3} \sum_l H_{j_1, l}^{j_1} L_{j_1}^l - \frac{2}{3} \sum_l H_{j_1, j_1}^l L_l^{j_1} - \frac{1}{2} H_{j_1, j_1}^{j_1} \Theta^{n+1} + \frac{1}{2} \sum_l L_{j_1}^l \Theta^l + \\ &+ \frac{1}{2} \Theta^{j_1} \Theta^{n+1}. \end{aligned} \right.$$

Thirdly:

(3.26)

$$\left\{ \begin{aligned} \Theta_{x^{j_1}}^{n+1} &= -\frac{2}{3} \frac{H_{j_1, j_1, y}^{j_1}}{L_{j_1, x^{j_1}}^{j_1}} + \frac{1}{3} \frac{L_{j_1, x^{j_1}}^{j_1}}{L_{j_1, x^{j_1}}^{j_1}} + \frac{2}{3} G_{j_1, j_1} M^{j_1} + \frac{4}{3} \sum_l G_{j_1, l} M^l - \frac{1}{2} \sum_l H_{l, l}^l L_{j_1}^l + \\ &+ \frac{1}{3} \sum_l H_{j_1, l}^{j_1} L_{j_1}^l - \frac{1}{3} \sum_l H_{j_1, j_1}^l L_l^{j_1} - \frac{1}{2} H_{j_1, j_1}^{j_1} \Theta^{n+1} + \frac{1}{2} \sum_l L_{j_1}^l \Theta^l + \\ &+ \frac{1}{2} \Theta^{j_1} \Theta^{n+1}. \end{aligned} \right.$$

Fourthly:

(3.27)

$$\left\{ \begin{aligned} \Theta_y^{n+1} &= -\frac{L_{j_1, y}^{j_1}}{L_{j_1, y}^{j_1}} + 2 \frac{M_{x^{j_1}}^{j_1}}{L_{j_1, x^{j_1}}^{j_1}} + 2 \sum_l H_{j_1, l}^{j_1} M^l - \sum_l H_{l, l}^l M^l - \frac{1}{2} \sum_l L_{j_1}^l L_l^{j_1} + \\ &+ \sum_l M^l \Theta^l + \frac{1}{2} \Theta^{n+1} \Theta^{n+1}. \end{aligned} \right.$$

These four families of partial differential equations constitute the *second auxiliary system*. By replacing these solutions in the three remaining families of equations (3.20), (3.21) and (3.22), we obtain supplementary equations (which we do not copy) that are direct consequences of (I'), (II'), (III'), (IV').

To complete the proof of the main Lemma 3.3 above, it suffices now to establish the first implication of the following list, since the other three have been already established.

- Some given functions G_{j_1, j_2} , $H_{j_1, j_2}^{k_1}$, $L_{j_1}^{k_1}$ and M^{k_1} of (x^{l_1}, y) satisfy the four families of partial differential equations (I'), (II'), (III') and (IV') of Theorem 1.7.
- ↓
- There exist functions Θ^{j_1} , Θ^{n+1} satisfying the second auxiliary system (3.24), (3.25), (3.26) and (3.27).
- ↓
- These solution functions Θ^{j_1} , Θ^{n+1} satisfy the six families of partial differential equations (3.17), (3.18), (3.19), (3.20), (3.21) and (3.22).
- ↓
- There exist functions $\Pi_{j_1, j_2}^{k_1}$ of (x^{l_1}, y) , $1 \leq j_1, j_2, k_1 \leq m+1$, satisfying the first auxiliary system (3.7) of partial differential equations.
- ↓
- There exist functions X^{l_2} , Y of (x^{l_1}, y) transforming the system $y_{x^{j_1} x^{j_2}} = F^{j_1, j_2}(x^{l_1}, y, y_{x^{l_2}})$, $j_1, j_2 = 1, \dots, n$, to the simplest system $Y_{X^{j_1} X^{j_2}} = 0$, $j_1, j_2 = 1, \dots, n$.

3.28. Compatibility conditions for the second auxiliary system. We notice that the second auxiliary system is also a complete system. Thus, to establish the first above

implication, it suffices to show that the four families of compatibility conditions:

$$(3.29) \quad \begin{cases} 0 = (\Theta_{x^j_2}^{j_1})_{x^j_3} - (\Theta_{x^j_3}^{j_1})_{x^j_2}, \\ 0 = (\Theta_{x^j_2}^{j_1})_y - (\Theta_y^{j_1})_{x^j_2}, \\ 0 = (\Theta_{x^j_1}^{n+1})_{x^j_2} - (\Theta_{x^j_2}^{n+1})_{x^j_1}, \\ 0 = (\Theta_{x^j_2}^{n+1})_y - (\Theta_y^{n+1})_{x^j_2}, \end{cases}$$

are a consequence of (I'), (I''), (III'), (IV').

For instance, in (3.29)₁, replacing $\Theta_{x^j_2}^{j_1}$ by its expression (3.24), differentiating it with respect to x^j_3 , replacing $\Theta_{x^j_3}^{j_1}$ by its expression (3.24), differentiating it with respect to x^j_2 and subtracting, we get:

$$(3.30) \quad \left\{ \begin{aligned} 0 = & -2 G_{j_1, j_2, y x^j_3} + 2 G_{j_1, j_3, y x^j_2} + \frac{H_{j_1, j_1, x^j_2 x^j_3}^{j_1}}{a} - \frac{H_{j_1, j_1, x^j_3 x^j_2}^{j_1}}{a} + \\ & + \frac{1}{2} \frac{\Theta_{x^j_3}^{j_1}}{x^j_3} \Theta^{j_2} + \frac{1}{2} \Theta^{j_1} \frac{\Theta^{j_2}}{x^j_3} - \frac{1}{2} \frac{\Theta_{x^j_2}^{j_1}}{x^j_2} \Theta^{j_3} - \frac{1}{2} \Theta^{j_1} \frac{\Theta^{j_3}}{x^j_2} - \\ & - \frac{1}{2} H_{j_1, j_1, x^j_3}^{j_1} \Theta^{j_2} - \frac{1}{2} H_{j_1, j_1}^{j_1} \frac{\Theta^{j_2}}{x^j_3} + \frac{1}{2} H_{j_1, j_1, x^j_2}^{j_1} \Theta^{j_3} + \frac{1}{2} H_{j_1, j_1}^{j_1} \frac{\Theta^{j_3}}{x^j_2} - \\ & - \frac{1}{2} H_{j_2, j_2, x^j_3}^{j_2} \Theta^{j_1} - \frac{1}{2} H_{j_2, j_2}^{j_2} \frac{\Theta^{j_1}}{x^j_3} + \frac{1}{2} H_{j_3, j_3, x^j_2}^{j_3} \Theta^{j_1} + \frac{1}{2} H_{j_3, j_3}^{j_3} \frac{\Theta^{j_1}}{x^j_2} - \\ & - G_{j_1, j_2, x^j_3} \Theta^{n+1} - G_{j_1, j_2} \frac{\Theta^{n+1}}{x^j_3} + G_{j_1, j_3, x^j_2} \Theta^{n+1} + G_{j_1, j_3} \frac{\Theta^{n+1}}{x^j_2} + \\ & + \sum_l H_{j_1, j_2, x^j_3}^l \Theta^l + \sum_l H_{j_1, j_2}^l \frac{\Theta^l}{x^j_3} - \sum_l H_{j_1, j_3, x^j_2}^l \Theta^l - \sum_l H_{j_1, j_3}^l \frac{\Theta^l}{x^j_2} + \\ & + \frac{1}{2} H_{j_1, j_1, x^j_3}^{j_1} H_{j_2, j_2}^{j_2} + \frac{1}{2} H_{j_1, j_1}^{j_1} H_{j_2, j_2, x^j_3}^{j_2} - \frac{1}{2} H_{j_1, j_1, x^j_2}^{j_1} H_{j_3, j_3}^{j_3} - \frac{1}{2} H_{j_1, j_1}^{j_1} H_{j_3, j_3, x^j_2}^{j_3} - \\ & - \sum_l H_{j_1, j_2, x^j_3}^l H_{l, l}^l - \sum_l H_{j_1, j_2}^l H_{l, l, x^j_3}^l + \sum_l H_{j_1, j_3, x^j_2}^l H_{l, l}^l + \sum_l H_{j_1, j_3}^l H_{l, l, x^j_2}^l + \\ & + \sum_l G_{j_2, l, x^j_3} L_{j_1}^l + \sum_l G_{j_2, l} L_{j_1, x^j_3}^l - \sum_l G_{j_3, l, x^j_2} L_{j_1}^l - \sum_l G_{j_3, l} L_{j_1, x^j_2}^l. \end{aligned} \right.$$

Next, replacing the twelve first order partial derivatives underlined just above:

$$(3.31) \quad \left\{ \begin{aligned} \frac{\Theta_{x^j_3}^{j_1}}{x^j_3}, \quad \frac{\Theta_{x^j_3}^{j_2}}{x^j_3}, \quad \frac{\Theta_{x^j_2}^{j_1}}{x^j_2}, \quad \frac{\Theta_{x^j_2}^{j_3}}{x^j_2}, \quad \frac{\Theta_{x^j_3}^{j_2}}{x^j_3}, \quad \frac{\Theta_{x^j_2}^{j_3}}{x^j_2}, \\ \frac{\Theta_{x^j_3}^{j_1}}{x^j_3}, \quad \frac{\Theta_{x^j_2}^{j_1}}{x^j_2}, \quad \frac{\Theta_{x^j_3}^{n+1}}{x^j_3}, \quad \frac{\Theta_{x^j_2}^{n+1}}{x^j_2}, \quad \frac{\Theta^l}{x^j_3}, \quad \frac{\Theta^l}{x^j_2}. \end{aligned} \right.$$

by their values issued from (3.24), (3.26) and adapting the summation indices, we get the explicit developed form of the first family of compatibility conditions (3.29)₁:

(3.32)

$$\begin{aligned}
 0 =? = & \frac{-2 G_{j_1, j_2, x^{j_3} y} + 2 G_{j_1, j_3, x^{j_2} y}}{} \\
 & - \frac{\sum_l G_{j_3, l, x^{j_2}} L_{j_1}^l + \sum_l G_{j_2, l, x^{j_3}} L_{j_1}^l - G_{j_1, j_2, y} H_{j_3, j_3}^{j_3} + G_{j_1, j_3, y} H_{j_2, j_2}^{j_2}}{} \\
 & - \frac{2 \sum_l G_{l, j_3} H_{j_1, j_2}^l + 2 \sum_l G_{l, j_2} H_{j_1, j_3}^l - \sum_l H_{j_1, j_2, x^{j_3}}^l H_{l, l}^l + \sum_l H_{j_1, j_3, x^{j_2}}^l H_l^l}{} \\
 & - \frac{\frac{2}{3} H_{j_2, j_2, y}^{j_2} G_{j_1, j_3} + \frac{2}{3} H_{j_3, j_3, y}^{j_3} G_{j_1, j_2} - \frac{2}{3} L_{j_3, x^{j_3}}^{j_3} G_{j_1, j_2} + \frac{2}{3} L_{j_2, x^{j_2}}^{j_2} G_{j_1, j_3}}{} \\
 & - \frac{\sum_l L_{j_1, x^{j_2}}^l G_{j_3, l} + \sum_l L_{j_1, x^{j_3}}^l G_{j_2, l}}{} \\
 & - \frac{2}{3} G_{j_1, j_2} G_{j_3, j_3} M^{j_3} + \frac{2}{3} G_{j_1, j_3} G_{j_2, j_2} M^{j_2} - \frac{4}{3} \sum_l G_{j_1, j_2} G_{j_3, l} M^l + \\
 & + \frac{4}{3} \sum_l G_{j_1, j_3} G_{j_2, l} M^l - \frac{1}{2} \sum_l G_{j_3, l} H_{j_1, j_1}^{j_1} L_{j_2}^l + \frac{1}{2} \sum_l G_{j_2, l} H_{j_1, j_1}^{j_1} L_{j_3}^l - \\
 & - \frac{1}{2} \sum_l G_{j_3, l} H_{j_2, j_2}^{j_2} L_{j_1}^l + \frac{1}{2} \sum_l G_{j_2, l} H_{j_3, j_3}^{j_3} L_{j_1}^l - \frac{1}{2} \sum_l G_{j_1, j_3} H_{l, l}^l L_{j_2}^l + \\
 & + \frac{1}{2} \sum_l G_{j_1, j_2} H_{l, l}^l L_{j_3}^l - \frac{1}{3} \sum_l G_{j_1, j_2} H_{j_3, l}^{j_3} L_{j_3}^l + \frac{1}{3} \sum_l G_{j_1, j_3} H_{j_2, l}^{j_2} L_{j_2}^l - \\
 & - \frac{1}{3} G_{j_1, j_3} H_{j_2, j_2}^l L_l^{j_2} + \frac{1}{3} G_{j_1, j_2} H_{j_3, j_3}^l L_l^{j_3} - \\
 & - \sum_l \sum_p G_{j_2, p} H_{j_1, j_3}^l L_l^p + \sum_l \sum_p G_{j_3, p} H_{j_1, j_2}^l L_l^p - \\
 & - \sum_l \sum_p H_{j_1, j_2}^l H_{l, j_3}^p H_{p, p}^p + \sum_l \sum_p H_{j_1, j_3}^l H_{l, j_2}^p H_{p, p}^p.
 \end{aligned}$$

Lemma 3.33. ([Me2003, Me2004]) *This first family of compatibility conditions for the second auxiliary system obtained by developing (3.29)₁ in length, together with the three remaining families obtained by developing (3.29)₂, (3.29)₃, (3.29)₄ in length, are consequences, by linear combinations and by differentiations, of (I'), (II'), (III'), (IV'), of Theorem 1.7.*

The summarized proof of Theorem 1.7 is complete. □

IV: Bibliography

REFERENCES

- [Ar1988] ARNOL'D, V.I.: *Dynamical systems. I. Ordinary differential equations and smooth dynamical systems*, Translated from the Russian. Edited by D. V. Anosov and V. I. Arnol'd. Encyclopaedia of Mathematical Sciences, vol. 1. Springer-Verlag, Berlin, 1988. x+233 pp.
- [Ar1968] ARTIN, M.: *On the solutions of analytic equations*, *Invent. Math.* **5** (1968), 277–291.
- [BER1999] BAOUENDI, M.S.; EBENFELT, P.; ROTHSCCHILD, L.P.: *Real submanifolds in complex space and their mappings*. Princeton Mathematical Series, vol. 47, Princeton University Press, Princeton, NJ, 1999, xii+404 pp.
- [BJT1985] BAOUENDI, M.S.; JACOBOWITZ, H.; TREVES, F.: *On the analyticity of CR mappings*, *Ann. of Math.* **122** (1985), no. 2, 365–400.
- [Bel1996] BELLAÏCHE, A.: *SubRiemannian Geometry*, Progress in Mathematics, vol. 144, Birkhäuser Verlag, Basel/Switzerland, 1996, 1–78.
- [Be1979] BELOSHAPKA, V.K.: *On the dimension of the group of automorphisms of an analytic hypersurface*, *Izv. Akad. Nauk SSSR Ser. Mat.* **43** (1979), 243–266; English transl. in *Math. USSR-Izv.* **14** (1980), 223–245.
- [Be1988] BELOSHAPKA, V.K.: *Finite-dimensionality of the group of automorphisms of a real-analytic surface*, *Izv. Akad. Nauk SSSR Ser. Mat.* **52** (1988), 437–442; English transl. in *Math. USSR-Izv.* **32** (1989), 443–448.
- [Be1997] BELOSHAPKA, V.K.: *CR-varieties of the type (1,2) as varieties of 'super-high' codimension*, *Russian J. Math. Phys.* **5** (1997), 399–404.
- [Be2002] BELOSHAPKA, V.K.: *Real submanifolds in complex space: polynomial models, automorphisms, and classification problems*, *Uspekhi Mat. Nauk* **57** (2002), no. 1, 3–44; English transl. in *Russian Math. Surveys* **57** (2002), no. 1, 1–41.
- [BES2005] BELOSHAPKA, V.K.; EZHOV, V.; SCHMALZ, G.: *Canonical Cartan connection and holomorphic invariants of Engel CR manifolds*, arxiv.org/abs/math.CV/0508084.
- [BK1989] BLUMAN, G.W.; KUMEI, S.: *Symmetries and differential equations*, Springer Verlag, Berlin, 1989.
- [Bo1991] BOGGESS, A.: *CR manifolds and the tangential Cauchy-Riemann complex*. Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1991, xviii+364 pp.
- [Bo1972] BOURBAKI, N.: *Groupes at algèbres de Lie*, chapitre 2, Hermann, Paris, 1972.
- [BSW1978] BURNS, D.Jr.; SHNIDER, S.; WELLS, R.O.Jr.: *Deformations of strictly pseudoconvex domains* *Invent. Math.* **46** (1978), no. 3, 237–253.
- [Ca1922] CARTAN, É.: *Sur les équations de la gravitation d'Einstein*, *J. Math. pures et appl.* **1** (1922), 141–203.
- [Ca1924] CARTAN, É.: *Sur les variétés à connexion projective*, *Bull. Soc. Math. France* **52** (1924), 205–241.
- [Ca1932a] CARTAN, É.: *Sur la géométrie pseudo-conforme des hypersurfaces de l'espace de deux variables complexes, I*, *Ann. Math. Pura Appl.* **11** (1932), 17–90.
- [Ca1932b] CARTAN, É.: *Sur la géométrie pseudo-conforme des hypersurfaces de l'espace de deux variables complexes, II*, *Ann. Scuola Norm. Sup. Pisa* **1** (1932), 333–354.
- [Ca1937] CARTAN, É.: *Les problèmes d'équivalence* (Séminaire de Math., 4^e année, 1936–37), *Œuvres complètes*, II, 1311–1334, Gauthier-Villars, Paris, 1953.
- [CM1974] CHERN, S.S.; MOSER, J.K.: *Real hypersurfaces in complex manifolds*, *Acta Math.* **133** (1974), no. 2, 219–271.
- [Ch1975] CHERN, S.-S.: *On the projective structure of a real hypersurface in \mathbb{C}^{n+1}* , *Math. Scand.* **36** (1975), 74–82.
- [Ch1989] CHIRKA, E.M.: *Complex analytic sets*, Mathematics and its applications (Soviet Series), vol. 46. Kluwer Academic Publishers Group, Dordrecht, 1989. xx+372 pp.
- [Ch1991] CHIRKA, E.M.: *An introduction to the geometry of CR manifolds* (Russian), *Uspekhi Mat. Nauk* **46** (1991), no. 1(277), 81–164, 240; translation in *Russian Math. Surveys* **46** (1991), no. 1, 95–197
- [CS1996] CONSTANTINE, G.M.; SAVITS, T.H.: *A multivariate Faà di Bruno formula with applications*, *Trans. Amer. Math. Soc.* **348** (1996), no. 2, 503–520.

- [DF1988] DIEDERICH, K.; FORNÆSS, J.E.: *Proper holomorphic mappings between real-analytic pseudoconvex domains in \mathbb{C}^n* , Math. Ann **282** (1988), no. 4, 681–700.
- [DP2003] DIEDERICH, K.; PINCHUK, S.: *Regularity of continuous CR-maps in arbitrary dimension*, Michigan Math. J. **51** (2003), 111–140. *Erratum: ib.*, no. 3, 667–668.
- [DW1980] DIEDERICH, K.; WEBSTER, S.M.: *A reflection principle for degenerate real hypersurfaces*, Duke Math. J. **47** (1980), no. 4, 835–843.
- [Eb1998] EBENFELT, P.: *Normal forms and biholomorphic equivalences of real hypersurfaces in \mathbb{C}^3* , Indiana Univ. Math. J. **47** (1998), 311–366.
- [Eb2006] EBENFELT, P.: *Correction to "Uniformly Levi degenerate CR manifolds: the 5-dimensional case"*, Duke Math. J. **131** (2006), 589–591.
- [EL1890] ENGEL, F.; LIE, S.: *Theorie der Transformationsgruppen, I, II, II*, Teubner, Leipzig, 1889, 1891, 1893.
- [EKV1985] EZHOV, V.V.; KRUZHILIN, N.G.; VITUSHKIN, A.G.: *Continuation of holomorphic mappings along real-analytic hypersurfaces* (Russian). Current problems in mathematics. Mathematical analysis, algebra, topology. Trudy Mat. Inst. Steklov **167** (1985), 60–95, 276.
- [Fa1980] FARAN, J.: *Segre families and real hypersurfaces*, Invent. Math. **60** (1980), no. 2, 135–172.
- [F1969] FEDERER, H.: *Geometric measure theory*, Die Grundlehren der Mathematischen Wissenschaften, Band 153, Springer Verlag, New York, 1969, xiv+676 pp. Springer-Verlag, Berlin, 1969.
- [Fe1995] FELS, M.: *The equivalence problem for systems of second-order ordinary differential equations*, Proc. London Math. Soc. **71** (1995), no. 2, 221–240.
- [FK2005a] FELS, G.; KAUP, W.: *CR-manifolds of dimension 5: A Lie algebra approach*, J. Reine Angew. Math., to appear. arxiv.org/abs/math.DG/050811.
- [FK2005b] FELS, G.; KAUP, W.: *Homogeneous Levi degenerate CR-manifolds in dimension 5*, to appear.
- [Fr1877] FROBENIUS, G.: *Ueber das Pfaffsche Problem*, J. Reine Angew. Math. **82** (1877), 230–315.
- [G1989] GARDNER, R.B.: *The method of equivalence and its applications*, CBMS-NSF Regional Conference Series in Applied Mathematics, vol. 58 (SIAM, Philadelphia, 1989), 127 pp.
- [GM2003a] GAUSSIER, H.; MERKER, J.: *Symmetries of partial differential equations*, J. Korean Math. Soc. **40** (2003), no. 3, 517–561; e-print: <http://fr.arxiv.org/abs/math.CV/0404127>.
- [GM2003b] GAUSSIER, H.; MERKER, J.: *A new example of uniformly Levi nondegenerate hypersurface in \mathbb{C}^3* , Ark. Mat. **41** (2003), no. 1, 85–94.
- [GM2004] GAUSSIER, H.; MERKER, J.: *Nonalgebraizable real analytic tubes in \mathbb{C}^n* , Math. Z. **247** (2004), no. 2, 337–383.
- [GM2006] GAUSSIER, H.; MERKER, J.: *Erratum to "A new example of a uniformly Levi degenerate hypersurface in \mathbb{C}^3 "*, 2006, to appear.
- [GV1987] GERSHKOVICH, V.Ya.; VERSHIK, A.M.: *Nonholonomic dynamical systems. Geometry of distributions and variational problems*. Dynamical Systems VII, Encyclopædia of mathematical sciences, vol. 16, V.I. Arnol'd and S.P. Novikov (Eds.), 1–81, Springer-Verlag, Berlin, 1994.
- [Gr2005] DE GRAAF, W.A.: *Classification of solvable Lie algebras*, Experiment. Math. **14** (2005), no. 1, 15–25.
- [GTW1989] GRISSOM, C.; THOMPSON, G.; WILKENS, G.: *Linearization of second order ordinary differential equations via Cartan's equivalence method*, J. Diff. Eq. **77** (1989), no. 1, 1–15.
- [Gr2000] GROSSMAN, D.A.: *Torsion-free path geometries and integrable second order ODE systems*, Selecta Math. (N.S.) **6** (2000), no. 4, 399–442.
- [Ha1937] HACHTROUDI, M.: *Les espaces d'éléments à connexion projective normale*, Actualités Scientifiques et Industrielles, vol. 565, Paris, Hermann, 1937.
- [Ha1982] HARTMAN, P.: *Ordinary Differential Equations*. Birkhäuser, Boston 1982.
- [Ha2003] HAUSER, H.: *The Hironaka theorem on resolution of singularities (or: A proof we always wanted to understand)*, Bull. Amer. Math. Soc. (N.S.) **40** (2003), no. 3, 323–403.
- [Hi1976] HIRSCH, M.W.: *Differential topology*, Graduate Texts in Mathematics, 33, Springer-Verlag, Berlin, 1976, x+222 pp.
- [HK1989] HSU, L.; KAMRAN, N.: *Classification of second order ordinary differential equations admitting Lie groups of fibre-preserving point symmetries*, Proc. London Math. Soc. **58** (1989), no. 3, 387–416.
- [Ib1992] IBRAGIMOV, N.H.: *Group analysis of ordinary differential equations and the invariance principle in mathematical physics*, Russian Math. Surveys **47:4** (1992), 89–156.

- [Ib1999] IBRAGIMOV, N.H.: *Elementary Lie group analysis and ordinary differential equations*, Wiley Series in Mathematical Methods in Practice, 4. John Wiley & Sons, Ltd., Chichester, 1999. xviii+347 pp.
- [IL2003] IVEY, J.A.; LANDSBERG, J.M.: *Cartan for beginners: differential geometry via moving frames and exterior differential systems*, Graduate Studies in Mathematics, 61. American Mathematical Society, Providence, RI, 2003. xiv+378 pp.
- [Ja1990] JACOBOWITZ, *An introduction to CR structures*, Math. Surveys and Monographs, 32. Amer. Math. Soc., Providence, 1990. x+237 pp.
- [Ji2002] JI, S.: *Algebraicity of real analytic hypersurfaces with maximal rank*, Amer. J. Math. **124** (2002), no. 6, 1083–1102.
- [JoPf2000] DE JONG, T.; PFISTER, G.: *Local analytic geometry. Basic theory and applications*, Advanced Lectures in Mathematics. Friedr. Vieweg & Sohn, Braunschweig, 2000. xii+382 pp.
- [Kn2004] KNAPP, A.W.: *Lie groups beyond an introduction*, Progress in Mathematics, 140, Birkhäuser, Basel, third edition, 2004, xviii+812 pp.
- [KN1963] KOBAYASHI, S.; NOMIZU, K.m: *Foundations of differential geometry*, I, Interscience publishers, John Wiley & Sons, New York, 1963. xi+329 pp.
- [Kr1985] KRUZHILIN, N.G.: *Local automorphisms and mappings of smooth strictly pseudoconvex hypersurfaces* (Russian). Izv. Akad. Nauk SSSR Ser. Mat. **49** (1985), no. 3, 566–591, 672.
- [Kr1987] KRUZHILIN, N.G.: *Description of the local automorphism groups of real hypersurfaces*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986), 749–758, Amer. Math. Soc., Providence, RI, 1987.
- [KV1987] KRUZHILIN, N.G.; VITUSHKIN, A.G.: *Description of the automorphism groups of real hypersurfaces in complex space* (Russian). Investigations in the theory of the approximation of functions, 26–69, Akad. Nauk SSSR Bashkir Filial, Otdel. Fiz. Mat., Ufa, 1987.
- [Lie1880] LIE, S.: *Theorie der Transformationsgruppen*, Math. Ann. **16** (1880), 441–528.
- [Lie1883] LIE, S.: *Klassifikation und Integration vo gewöhnlichen Differentialgleichungen zwischen x, y , die eine Gruppe von Transformationen gestaten I-IV*. In: Gesammelte Abhandlungen, Vol. 5, B.G. Teubner, Leipzig, 1924, pp. 240–310; 362–427, 432–448.
- [LS1893] LIE, S.; SCHEFFERS, G.: *Vorlesungen über continuierliche Gruppen mit Geometrischen und anderen Anwendungen*. (German) Nachdruck der Auflage des Jahres 1893. Chelsea Publishing Co., Bronx, New York, 1971. xii+810 pp.
- [Lo1981] LOBODA, A.V.: *Local automorphisms of real-analytic hypersurfaces* (Russian), Izv. Akad. Nauk SSSR Ser. Mat. **45** (1981), no. 3, 620–645.
- [Lo2001] LOBODA, A.V.: *Homogeneous strictly pseudoconvex hypersurfaces in \mathbb{C}^3 with two-dimensional isotropy groups* (Russian) Mat. Sb. **192** (2001), no. 12, 3–24; translation in Sb. Math. **192** (2001), no. 11-12, 1741–1761.
- [Lo2002] LOBODA, A.V.: *Homogeneous nondegenerate surfaces in \mathbb{C}^3 with two-dimensional isotropy groups* (Russian) Mat. Sb. **192** (2001), no. 12, 3–24; translation in Sb. Math. **192** (2001), no. 11–12, 1741–1761.
- [Lo2003] LOBODA, A.V.: *On the determination of a homogeneous strictly pseudoconvex hypersurface from the coefficients of its normal form* (Russian) Mat. Zametki **73** (2003), no. 3, 453–456; translation in Math. Notes **73** (2003), no. 3-4, 419–423.
- [Ma2003] MARDARE, S.: *On isometric immersions of a Riemannian space under a weak regularity assumption*, C. R. Acad. Sci. Paris, Sér. I **337** (2003), 785–790.
- [Me2001] MERKER, J.: *On the partial algebraicity of holomorphic mappings between two real algebraic sets in the complex euclidean spaces of different dimensions*, Bull. Soc. Math. France **129** (2001), no. 4, 547–591.
- [Me2003] MERKER, J.: *hand manuscript I*, 212 pp., May–July 2003; *hand manuscript II*, 114 pp., August 2003.
- [Me2004] MERKER, J.: *Explicit differential characterization of the Newtonian free particle system in $m \geq 2$ dependent variables*, Acta Mathematicæ Applicandæ, to appear, 73 pp; e-print: arxiv.org/abs/math.DG/0411165.
- [Me2005a] MERKER, J.: *On the local geometry of generic submanifolds of \mathbb{C}^n and the analytic reflection principle* (Part I), Journal of Mathematical Sciences (N.Y.) **125** (2005), no. 6, 751–824.
- [Me2005b] MERKER, J.: *Étude de la régularité analytique de l'application de réflexion CR formelle*, Annales Fac. Sci. Toulouse, **XIV** (2005), no. 2, 215–330.

- [MP2005] MERKER, J.; PORTEN, E.: *Holomorphic extension of CR functions, envelopes of holomorphy and removable singularities*, 432 pp., to appear. Downloadable at: www.cmi.univ-mrs.fr/~merker/index.html.
- [N2003] NEUT, S.: *Implantation et nouvelles applications de la méthode d'équivalence d'Élie Cartan*, Thèse, Université Lille 1, October 2003.
- [OL1979] OLVER, P.J.: *Symmetry groups and group invariant solutions of partial differential equations*, J. Diff. Geom. **14** (1979), 497–542.
- [OI1986] OLVER, P.J.: *Applications of Lie groups to differential equations*. Springer Verlag, New York, 1986. xxvi+497 pp.
- [OI1995] OLVER, P.J.: *Equivalence, Invariance and Symmetries*. Cambridge, Cambridge University Press, 1995, xvi+525 pp.
- [OV1994] ONISHCHIK, A.L.; VINBERG, E.B.: *Lie groups and Lie algebras, III*. Encyclopædia of mathematical sciences, 41. Springer Verlag, Berlin, 248 pp.
- [Pi1975] PINCHUK, S.: *On the analytic continuation of holomorphic mappings* (Russian), Mat. Sb. (N.S.) **98(140)** (1975) no.3(11), 375–392, 416–435, 495–496.
- [Pi1978] PINCHUK, S.: *Holomorphic mappings of real-analytic hypersurfaces* (Russian), Mat. Sb. (N.S.) **105(147)** (1978), no. 4, 574–593, 640; English translation in Math. USSR Sbornik **34** (1978), 503–519.
- [Re1993] REUTENAUER, C.: *Free Lie algebras*, London Mathematical Society Monograph, New Series, 7. Oxford Science Publications, The Clarendon Press, Oxford University Press, New York, 1993. xviii+269 pp.
- [Se1931] SEGRE, B.: *Intorno al problema di Poincaré della rappresentazione pseudoconforme*, Rend. Acc. Lincei, VI, Ser. **13** (1931), 676–683.
- [Se1932] SEGRE, B.: *Questioni geometriche legate colla teoria delle funzioni di due variabili complesse*, Rendiconti del Seminario di Matematici di Roma, II, Ser. **7** (1932), no. 2, 59–107.
- [Sh1997] SHARPE, R.W.: *Differential geometry; Cartan's generalization of Klein's Erlangen program*, Graduate texts in mathematics, vol. 166, Springer Verlag, Berlin, 1997, xix+421 pp.
- [Sp1970] SPIVAK, M.: *A comprehensive introduction to differential geometry*, vols. 1 and 2. Published by M. Spivak, Brandeis Univ., Waltham, Mass. 1970.
- [St1996] STANTON, N.: *Infinitesimal CR automorphisms of real hypersurfaces*, Amer. J. Math. **118** (1996), no. 1, 209–233.
- [Ste1983] STERNBERG, S.: *Lectures in differential geometry*, Second edition, Chelsea publishing co., New York, 1983, xviii+442 pp.
- [Stk1982] STORMARK, O.: *On the theorem of Frobenius for complex vector fields*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **9** (1982), no. 1, 57–90.
- [Stk2000] STORMARK, O.: *Lie's structural approach to PDE systems*, Encyclopædia of mathematics and its applications, vol. 80, Cambridge University Press, Cambridge, 2000, xv+572 pp.
- [Su2001] SUKHOV, A.: *Segre varieties and Lie symmetries*, Math. Z. **238** (2001), no. 3, 483–492.
- [Su2002] SUKHOV, A.: *CR maps and point Lie transformations*, Michigan Math. J. **50** (2002), 369–379.
- [Su2003] SUKHOV, A.: *On transformations of analytic structures* (Russian). Izv. Ross. Akad. Nauk Ser. Mat. **67** (2003), no. 2, 101–132; transl. in Izv. Math. **67** (2003), no. 2, 303–332.
- [Su1973] SUSSMANN, H.J.: *Orbits of families of vector fields and integrability of distributions*, Trans. Amer. Math. Soc. **180** (1973), no. 1, 171–188.
- [Tr1896] TRESSE, A.: *Détermination des invariants ponctuels de l'équation différentielle du second ordre $y'' = \omega(x, y, y')$* , Hirzel, Leipzig, 1896.
- [Ve1924] VESSIOT, E.: *Sur une théorie nouvelle des problèmes généraux d'intégration*, Bull. Soc. Math. France **52** (1924), 336–395.
- [Vi1990] VITUSHKIN, A.G.: *Holomorphic mappings and the Geometry of Hypersurfaces*, Encyclopædia of Mathematical Sciences, Volume 7, Several Complex Variables, I, Springer-Verlag, Berlin, 1990, pp. 159–214.
- [We1977] WEBSTER, S.M.: *On the mapping problem for algebraic real hypersurfaces*, Invent. Math. **43** (1977), no.1, 53–68.