



Designation: E 1242 – 97

Standard Practice for Using Octanol-Water Partition Coefficient to Estimate Median Lethal Concentrations for Fish Due to Narcosis¹

This standard is issued under the fixed designation E 1242; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon (ϵ) indicates an editorial change since the last revision or reapproval.

1. Scope

1.1 This practice covers a procedure for estimating the fathead minnow (*Pimephales promelas*) 96-h LC₅₀ of non-reactive (that is, covalently bonded without unsaturated residues) and nonelectrolytic (that is, require vigorous reagents to facilitate substitution, addition, replacement reactions and are non-ionic, non-dissociating in aqueous solutions) organic chemicals acting solely by narcosis, also referred to as Meyer-Overton toxicity relationship.²

1.2 This procedure is accurate for organic chemicals that are toxic due to narcosis and are non-reactive and non-electrolytic. Examples of appropriate chemicals are: alcohols, ketones, ethers, simple halogenated aliphatics, aromatics, and aliphatic substituted aromatics. It is not appropriate for chemicals whose structures include a potential toxiphore (that structural component of a chemical molecule that has been identified to show mammalian toxicity, for example CN is known to be responsible for inactivation of enzymes, NO₂ for decoupling of oxidative phosphorylation, both leading to mammalian toxicity). Examples of inappropriate chemicals are: carbamates, organophosphates, phenols, beta-gamma unsaturated alcohols, electrophiles, and quaternary ammonium salts.

2. Referenced Documents

2.1 ASTM Standards:

E 729 Guide for Conducting Acute Toxicity Tests with Fishes, Macroinvertebrates, and Amphibians³

E 943 Terminology Relating to Biological Effects and Environmental Fate³

E 1023 Guide for Assessing the Hazard of a Material to Aquatic Organisms and Their Uses³

E 1147 Test Method for Partition Coefficient (N-Octanol/Water) Estimation by Liquid Chromatography³

3. Terminology

3.1 Definitions:

3.1.1 *narcosis*—a reversible state of stupor, unconsciousness, or arrested activity produced by the influence of chemicals on critical sites within membranes or by disrupting the normal functioning of certain proteins by means of nonspecific binding of organic chemical(s) to hydrophobic sites. Death results if exposure is not terminated after a length of time which varies with concentration.

3.1.2 *octanol-water partition coefficient* (K_{ow})—referred to as P in some literature.

3.1.3 *toxiphore*—a chemical structure substituent group that when present gives rise to an adverse effect in exposed organisms.

3.2 For definitions of other terms used in this standard, refer to Guide E 729, Terminology E 943, and Guide E 1023.

4. Summary of Practice

4.1 The hydrophobicity of a non-reactive and non-electrolytic organic chemical as quantified by log of the octanol-water partition coefficient is substituted into an experimentally derived equation and an approximate 96-h log LC₅₀ for fathead minnow is calculated. This value is a maximum value. The actual LC₅₀ could be lower, but should not be higher.

5. Significance and Use

5.1 This procedure can be used to limit the need for screening tests prior to performing a test for estimating the LC₅₀ of a non-reactive and non-electrolytic chemical to the fathead minnow. By eliminating the screening test, fewer fish need be tested. The time used for preparing and performing the screening test can also be saved. The value obtained in this procedure can be used as the preliminary estimate of the LC₅₀ in a full-scale test.

5.2 Estimates can be used to set testing priority of groups of non-reactive and non-electrolytic chemicals.

5.3 If the estimated value is more than 0.3 times the experimental value, the mechanism of action is probably narcosis. If less, the effect concentration is considered to reflect a different mechanism of action.

5.4 This practice estimates a maximum LC₅₀, that is,

¹ This practice is under the jurisdiction of ASTM Committee E-47 on Biological Effects and Environmental Fate and is the direct responsibility of Subcommittee E47.01 on Aquatic Assessment and Toxicology.

Current edition approved Feb. 10, 1997. Published April 1997. Originally published as E 1242 – 88. Last previous edition E 1242 – 88.

² Lipnick, Robert L., "Validation and Extension of Fish Toxicity QSARs and Interspecies Comparisons for Certain Classes of Organic Chemicals," *QSAR in Toxicology and Xenobiochemistry*, Elsevier, 1985.

³ *Annual Book of ASTM Standards*, Vol 11.05.



non-reactive and non-electrolytic chemicals are at least as toxic as the practice predicts, but may have a lower LC50 if acting by a more specific mechanism. Data on a chemical indicating a lower toxicity than predicted should be considered suspect or an artifact because of limited solubility of the test material.

6. Procedure

6.1 Review other toxicity data on the test material to see if the test material is likely to be acutely toxic by a mechanism other than narcosis. Also, review the structure of the test material for toxic substructures. If there are no toxiphores present or other data which indicate the non-reactive and non-electrolytic chemical will be acutely toxic by a mechanism other than simple narcosis, this procedure will provide a maximum 96-h LC50 estimate or hypothetical narcosis 96-h LC50.

6.2 Obtain the octanol-water partition coefficient (K_{ow}), by measurement (see Test Method E 1147), from a literature source or by estimation based on structure.⁴ Since the K_{ow} value is critical to calculating the LC50, it is important to obtain a measured value rather than depending completely on literature or estimation based on structure.

6.3 Derive the relationship between the LC50 and K_{ow} in the form:

$$\text{Log LC50} = a \log K_{ow} + b \log (c K_{ow} + 1) - d \quad (1)$$

⁴ Great Lakes Water Quality Initiative Technical Support Document for the Procedure to Determine Bioaccumulation Factors Appendix A, Procedure for Deriving Recommended Values for Log K_{ow} , EPA-820-B-95-005, 1995.

using the bilinear model of Kubinyi.^{5,6} Calculate the Coefficient of Determination (R^2).

6.4 Using this relationship, calculate an estimated LC50 for additional non-reactive and non-electrolytic chemicals for which the K_{ow} is known or estimated.

6.5 Compare the calculated 96-h LC50 with solubility to be sure it is equal to or less than the solubility limit. If the estimated LC50 is greater than the solubility, the chemical is probably not lethal at the highest possible dissolved concentration in water.

7. Report

7.1 Report the following information:

7.1.1 The coefficients derived using the relationship in 6.3 and the coefficient of determination,

7.1.2 The estimated 96-h LC50 for a non-reactive and non-electrolytic chemical of interest based on the derived relationship,

7.1.3 Sources of data for LC50 and K_{ow} from which the relationship was defined,

7.1.4 Source of the K_{ow} for a non-reactive and non-electrolytic chemical of interest,

7.1.5 Name and address of the persons making each derivation of the LC50 K_{ow} relationship and estimates, and

7.1.6 Date of the estimates.

⁵ Kubinyi, H., "Non-Linear Dependence of Biological Activity on Hydrophobic Character: The Bilinear Model," *Farmaco (Pavia) Ed. Sci.*, Vol 34, pp. 248–276.

⁶ Veith, G. D., Call, D. J., and Brooke, L. T., "Structure-Toxicity Relationships for the Fathead Minnow, *Pimephales promelas*: Narcotic Industrial Chemicals," *Canadian Journal of Fisheries and Aquatic Sciences*, Vol 40, 1983, pp. 743–748.

ASTM International takes no position respecting the validity of any patent rights asserted in connection with any item mentioned in this standard. Users of this standard are expressly advised that determination of the validity of any such patent rights, and the risk of infringement of such rights, are entirely their own responsibility.

This standard is subject to revision at any time by the responsible technical committee and must be reviewed every five years and if not revised, either reapproved or withdrawn. Your comments are invited either for revision of this standard or for additional standards and should be addressed to ASTM International Headquarters. Your comments will receive careful consideration at a meeting of the responsible technical committee, which you may attend. If you feel that your comments have not received a fair hearing you should make your views known to the ASTM Committee on Standards, at the address shown below.

This standard is copyrighted by ASTM International, 100 Barr Harbor Drive, PO Box C700, West Conshohocken, PA 19428-2959, United States. Individual reprints (single or multiple copies) of this standard may be obtained by contacting ASTM at the above address or at 610-832-9585 (phone), 610-832-9555 (fax), or service@astm.org (e-mail); or through the ASTM website (www.astm.org).