



Designation: E 1367 – 9903

Standard Guide Test Method for Conducting 10-day Static Sediment Measuring the Toxicity T of Sediment-Associated Contaminants with Marine and Estuarine Amphipods and Marine Invertebrates¹

This standard is issued under the fixed designation E 1367; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon (ϵ) indicates an editorial change since the last revision or reapproval.

¹ This guide test method is under the jurisdiction of ASTM Committee E-47 on Biological Effects and Environmental Fate and is the direct responsibility of Subcommittee E47.03 on Sediment Assessment and Toxicology.

Current edition approved ~~April 10, 1999~~; Nov. 1, 2003. Published ~~July 1999~~; February 2004. Originally published as E 1367 – 90. Last previous edition E 1367 – 92~~9~~.

1. Scope

1.1 This guide¹ describes

1.1 This test method covers procedures for testing estuarine or marine organisms in the laboratory to evaluate the short-term adverse effects toxicity of potentially contaminated sediment, contaminants associated with whole sediments. Sediments may be collected from the field or of a test material experimentally added spiked with compounds in the laboratory. General guidance is presented in Sections 1-151 to contaminated 15 for conducting sediment toxicity tests with estuarine or uncontaminated sediment, on marine amphipods. Specific guidance for conducting 10-d sediment toxicity tests with estuarine infaunal or marine amphipods during static 10-day exposures. These procedures is outlined in Annex A1 and specific guidance for conducting 28-d sediment toxicity tests with *Leptocheirus plumulosus* is outlined in Annex A2.

1.2 Procedures are useful described for testing estuarine or marine amphipod crustaceans in 10-d laboratory exposures to evaluate the effects toxicity of various geochemical characteristics of contaminants associated with whole sediments (Annex A1; USEPA 1994a (1)). Sediments may be collected from the field or spiked with compounds in the laboratory. A toxicity method is outlined for four species of estuarine or marine sediment-burrowing amphipods found within United States coastal waters. The species are *Ampelisca abdita*, a marine species that inhabits marine and mesohaline portions of the Atlantic coast, the Gulf of Mexico, and San Francisco Bay; *Eohaustorius estuarius*, a Pacific coast estuarine amphipods species; *Leptocheirus plumulosus*, an Atlantic coast estuarine species; and *Rhepoxynius abronius*, a Pacific coast marine species. Generally, the method described may be applied to all four species, although acclimation procedures and some test conditions (that is, temperature and salinity) will be species-specific (Sections 12 and Annex A1). The toxicity test is conducted in 1-L glass chambers containing 175 mL of sediment and 775 mL of overlying seawater. Exposure is static (that is, water is not renewed), and the animals are not fed over the 10-d exposure period. The endpoint in the toxicity test is survival with reburial of surviving amphipods as an additional measurement that can be used as an endpoint for some of the test species (for *R. abronius* and *E. estuarius*). Performance criteria established for this test include the average survival of amphipods in negative control treatment must be greater than or equal to 90 %. Procedures are described for use with sediments with pore-water salinity ranging from >0 ‰ to fully marine.

1.3 A procedure is also described for determining the chronic toxicity of contaminants associated with whole sediments with the amphipod *Leptocheirus plumulosus* in laboratory exposures (Annex A2; USEPA-USACE 2001(2)). The toxicity test is conducted for 28 d in 1-L glass chambers containing 175 mL of sediment and about 775 mL of overlying water. Test temperature is $25^{\circ} \pm 2^{\circ}\text{C}$, and the recommended overlying water salinity is $5 \text{ ‰} \pm 2 \text{ ‰}$ (for test sediment with pore water at 1 ‰ to 10 ‰) or $20 \text{ ‰} \pm 2 \text{ ‰}$ (for test sediment with pore water >10 ‰). Four hundred millilitres of overlying water is renewed three times per week, at which times test organisms are fed. The endpoints in the toxicity test are survival, growth, and reproduction of amphipods. Performance criteria established for this test include the average survival of amphipods in negative control treatment must be greater than or equal to other infaunal taxa, although modifications 80 % and there must be measurable growth and reproduction in all replicates of the procedures appropriate negative control treatment. This test is applicable for use with sediments from oligohaline to fully marine environments, with a silt content greater than 5 % and a clay content less than 85 %.

1.4 A salinity of 5 or 20 ‰ is recommended for routine application of 28-d test with *L. plumulosus* (Annex A2; USEPA-USACE 2001 (2)) and a salinity of 20 ‰ is recommended for routine application of the 10-d test with *E. estuarius* or *L. plumulosus* (Annex A1). However, the salinity of the overlying water for tests with these two species might can be adjusted to a specific salinity of interest (for example, salinity representative of site of interest or the objective of the study may be to evaluate the influence of salinity on the bioavailability of chemicals in sediment). More importantly, the salinity tested must be within the tolerance range of the test organisms (as outlined in Annex A1 and Annex A2). If tests are conducted with procedures different from those described section 1.3 or in Table A1.1 (for example, different salinity, lighting, temperature, feeding conditions), additional tests are required to determine comparability of results (section 1.10). If there is not a need to make comparisons among studies, then the test could be conducted just at a selected salinity for 10-day static the sediment of interest.

1.5 Future revisions of this standard may include additional annexes describing whole-sediment toxicity tests with other groups of estuarine or marine invertebrates (for example, information presented in Guide E 1611 on sediment testing with polychaetes could be added as an annex to future revisions to this standard). Future editions to this standard may also include methods for conducting the toxicity tests in smaller chambers with less sediment (Ho et al. 2000 (3) , Ferretti et al. 2002 (4)).

1.6 Procedures outlined in this standard are based primarily on procedures described in the USEPA (1994a (1)), USEPA-USACE (2001(2)), Test Method E 1706, and Guides E 1391, E 1525, E 1688, Environment Canada (1992 (5)), DeWitt et al. (1992a (6); 1997a (7)), Emery et al. (1997 (8)), and Emery and Moore (1996 (9)), Swartz et al. (1985 (10)), DeWitt et al. (1989(11)), Scott and Redmond (1989 (12)), and Schlekot et al. (1992 (13)).

1.7 Additional sediment toxicity research and methods development are now in progress to (1) refine sediment spiking procedures, (2) refine sediment dilution procedures, (3) refine sediment Toxicity Identification Evaluation (TIE) procedures, (4) produce additional data on confirmation of responses in laboratory tests with natural populations of benthic organisms (that is, field

¹ Boldface numbers in parentheses refer to the list

² Microtox is a trademark of references at the end of this guide. Strategic Diagnostics Inc. 111 Pencader Drive Newark, Delaware 19702-3322.

³ This guide is based largely on Guide E 729 and Ref

⁴ For referenced ASTM standards, visit the ASTM website, www.astm.org, or contact ASTM Customer Service at service@astm.org. For (3): Annual Book of ASTM Standards volume information, refer to the standard's Document Summary page on the ASTM website.

validation studies), and (5) evaluate relative sensitivity of endpoints measured in 10- and 28-d toxicity tests using estuarine or marine amphipods. This information will be described in future editions of this standard.

1.8 Although standard procedures are described in Annex A2 of this standard for conducting chronic sediment tests with *L. plumulosus*, further investigation of certain issues could aid in the interpretation of test results. Some of these issues include further investigation to evaluate the relative toxicological sensitivity of the lethal and sublethal endpoints to a wide variety of chemicals spiked in sediment and to mixtures of chemicals in sediments from contamination gradients in the field (USEPA-USACE 2001 R(2)). Additional research is needed to evaluate the ability of the lethal and sublethal endpoints to estimate the responses of populations and communities of benthic invertebrates to contaminated sediments. Research is also needed to link the toxicity test endpoints to a field-validated population model of *L. plumulosus* that would then generate estimates of population-level responses of the amphipod to test sediments and thereby provide additional ecologically relevant interpretive guidance for the laboratory toxicity test.

1.9 This standard outlines specific test methods for evaluating the toxicity of sediments with *A. abdita*, *Eohaustorius*, *E. estuarius*, *Ampelisca abdita*, *Grandidierella japonica*, *L. plumulosus*, and *R. abronius*. While standard procedures are described in this standard, further investigation of certain issues could aid in the interpretation of test results. Some of these issues include the effect of shipping on organism sensitivity, additional performance criteria for organism health, sensitivity of various populations of the same test species, and confirmation of responses in laboratory tests with natural benthos populations.

1.10 General procedures described in this standard might be useful for conducting tests with other estuarine or marine organisms (for example, *Corophium* spp., *Grandidierella japonica*, *Lepidactylus dytiscus*, *Streblospio benedicti*), although modifications may be necessary. Results of tests, even those with the same species, using procedures different from those described in the test method may not be comparable and using these different procedures may alter bioavailability. Comparison of results obtained using modified versions of these procedures might provide useful information concerning new concepts and procedures for conducting sediment tests with aquatic organisms. If tests are conducted with procedures different from those described in this test method, additional tests are required to determine comparability of results. General procedures described in this test method might be useful for conducting tests with other aquatic organisms; however, modifications may be necessary.

1.11 Selection of Toxicity Testing Organisms:

1.11.1 The choice of a test organism has a major influence on the relevance, success, and interpretation of a test. Furthermore, no one organism is best suited for all sediments. The following criteria were considered when selecting test organisms to be described in this standard (Table 1 and Guide E 1525). Ideally, a test organism should: (1) have a toxicological database demonstrating relative sensitivity to a range of contaminants of interest in sediment, (2) have a database for interlaboratory comparisons of procedures (for example, round-robin studies), (3) be in direct contact with sediment, (4) be readily available from culture or through field collection, (5) be easily maintained in the laboratory, (6) be easily identified, (7) be ecologically or economically important, (8) have a broad geographical distribution, be indigenous (either present or historical) to the site being evaluated, or have a niche similar to organisms of concern (for example, similar feeding guild or behavior to the indigenous organisms), (9) be tolerant of a broad range of sediment physico-chemical characteristics (for example, grain size), and (10) be compatible with selected exposure methods and endpoints (Guide E 1525). Methods utilizing selected organisms should also be (11) peer reviewed (for example, journal articles) and (12) confirmed with responses with natural populations of benthic organisms.

1.11.2 Of these criteria (Table 1), a database demonstrating relative sensitivity to contaminants, contact with sediment, ease of culture in the laboratory or availability for field-collection, ease of handling in the laboratory, tolerance to varying sediment physico-chemical characteristics, and confirmation with responses with natural benthic populations were the primary criteria used for selecting *A. abdita*, *E. estuarius*, *L. plumulosus*, and *R. abronius* for the current edition of this standard for 10-d sediment tests (Annex A1). The species chosen for this method are intimately associated with sediment, due to their tube-dwelling or free-burrowing, and sediment ingesting nature. Amphipods have been used extensively to test the toxicity of marine, estuarine, and freshwater sediments (Swartz et al., 1985 (10); DeWitt et al., 1989 (11); Scott and Redmond, 1989 (12); DeWitt et al., 1992a; Schlegel et al., 1992 (13)). The selection of test species for this standard followed the consensus of experts in the field of sediment toxicology who participated in a workshop entitled "Testing Issues for Freshwater and Marine Sediments". The workshop was sponsored by USEPA Office of Water, Office of Science and Technology, and Office of Research and Development, and was held in Washington, D.C. from 16-18 September 1992 (USEPA, 1992 (14)). Of the candidate species discussed at the workshop, *A. abdita*, *E. estuarius*, *L. plumulosus*, and *R. abronius* best fulfilled the selection criteria, and presented the availability of a combination of one estuarine and one marine species each for both the Atlantic (the estuarine *L. plumulosus* and the marine *A. abdita*) and Pacific (the estuarine *E. estuarius* and the marine *R. abronius*) coasts. *Ampelisca abdita* is also native to portions of the Gulf of Mexico and San Francisco Bay. Many other organisms that might be appropriate for sediment testing do not now meet these selection criteria because little emphasis has been placed on developing standardized testing procedures for benthic organisms. For example, a fifth species, *Grandidierella japonica* was not selected because workshop participants felt that the use of this species was not sufficiently broad to warrant standardization of the method. Environment Canada (1992 (5)) has recommended the use of the following amphipod species for sediment toxicity testing: *Amphiporeia virginiana*, *Corophium volutator*, *Eohaustorius washingtonianus*, *Foxiphilus xiximeus*, and *Leptocheirus pinguis*. A database similar to those available for *A. abdita*, *E. estuarius*, *L. plumulosus*, and *R. abronius* must be developed in order for these and other organisms to be included in future editions of this standard.

1.11.3 The primary criterion used for selecting *L. plumulosus* for chronic testing of sediments was that this species is found in both oligohaline and mesohaline regions of estuaries on the East Coast of the United States and is tolerant to a wide range of sediment grain size distribution (USEPA-USACE 2001 (2), Annex Annex A2). This species is easily cultured in the laboratory and has a relatively short generation time (that is, about 24 d at 23°C, DeWitt et al. 1992a (6)) that makes this species adaptable to chronic testing (Section 12).

1.11.4 An important consideration in the selection of specific species for test method development is the existence of information concerning relative sensitivity of the organisms both to single chemicals and complex mixtures. Several studies have evaluated the sensitivities of *A. abdita*, *E. estuarius*, *L. plumulosus*, or *R. abronius*, either relative to one another, or to other commonly tested estuarine or marine species. For example, the sensitivity of marine amphipods was compared to other species that were used in generating saltwater Water Quality Criteria. Seven amphipod genera, including *Ampelisca abdita* and *Rhepoxynius abronius*, were among the test species used to generate saltwater Water Quality Criteria for 12 chemicals. Acute amphipod toxicity data from 4-d water-only tests for each of the 12 chemicals was compared to data for (1) all other species, (2) other benthic species, and (3) other infaunal species. Amphipods were generally of median sensitivity for each comparison. The average percentile rank of amphipods among all species tested was 57 %; among all benthic species, 56 %; and, among all infaunal species, 54 %. Thus, amphipods are not uniquely sensitive relative to all species, benthic species, or even infaunal species (USEPA 1994a (1)). Additional research may be warranted to develop tests using species that are consistently more sensitive than amphipods, thereby offering protection to less sensitive groups.

1.11.5 Williams et al. (1986 (15)) compared the sensitivity of the *R. abronius* 10-d whole sediment test, the oyster embryo (*Crassostrea gigas*) 48-h abnormality test, and the bacterium (*Vibrio fischeri*) 1-h luminescence inhibition test (that is, the Microtox⁴ test) to sediments collected from 46 contaminated sites in Commencement Bay, WA. *Rhepoxynius abronius* were exposed to whole sediment, while the oyster and bacterium tests were conducted with sediment elutriates and extracts, respectively. Microtox² was the most sensitive test, with 63 % of the sites eliciting significant inhibition of luminescence. Significant mortality of *R. abronius* was observed in 40 % of test sediments, and oyster abnormality occurred in 35 % of sediment elutriates. Complete concordance (that is, sediments that were either toxic or not-toxic in all three tests) was observed in 41 % of the sediments. Possible sources for the lack of concordance at other sites include interspecific differences in sensitivity among test organisms, heterogeneity in contaminant types associated with test sediments, and differences in routes of exposure inherent in each toxicity test. These results highlight the importance of using multiple assays when performing sediment assessments.

1.11.6 Several studies have compared the sensitivity of combinations of the four amphipods to sediment contaminants. For example, there are several comparisons between *A. abdita* and *R. abronius*, between *E. estuarius* and *R. abronius*, and between *A. abdita* and *L. plumulosus*. There are fewer examples of direct comparisons between *E. estuarius* and *L. plumulosus*, and no examples comparing *L. plumulosus* and *R. abronius*. There is some overlap in relative sensitivity from comparison to comparison within each species combination, which appears to indicate that all four species are within the same range of relative sensitivity to contaminated sediments.

1.11.6.1 Word et al. (1989 (16)) compared the sensitivity of *A. abdita* and *R. abronius* to contaminated sediments in a series of experiments. Both species were tested at 15°C. Experiments were designed to compare the response of the organism rather than to provide a comparison of the sensitivity of the methods (that is, *Ampelisca abdita* would normally be tested at 20°C). Sediments collected from Oakland Harbor, CA, were used for the comparisons. Twenty-six sediments were tested in one comparison, while 5 were tested in the other. Analysis of results using Kruskal Wallace rank sum test for both experiments demonstrated that *R. abronius* exhibited greater sensitivity to the sediments than *A. abdita* at 15°C. Long and Buchman (1989 (17)) also compared the sensitivity of *A. abdita* and *R. abronius* to sediments from Oakland Harbor, CA. They also determined that *A. abdita* showed less sensitivity than *R. abronius*, but they also showed that *A. abdita* was less sensitive to sediment grain size factors than *R. abronius*.

1.11.6.2 DeWitt et al. (1989 (11)) compared the sensitivity of *E. estuarius* and *R. abronius* to sediment spiked with fluoranthene and field-collected sediment from industrial waterways in Puget Sound, WA, in 10-d tests, and to aqueous cadmium (CdCl₂) in a 4-d water-only test. The sensitivity of *E. estuarius* was from two (to spiked-spiked sediment) to seven (to one Puget Sound, WA, sediment) times less sensitive than *R. abronius* in sediment tests, and ten times less sensitive to CdCl₂ in the water-only test. These results are supported by the findings of Pastorok and Becker (1990 (18)) who found the acute sensitivity of *E. estuarius* and *R. abronius* to be generally comparable to each other, and both were more sensitive than *Neanthes arenaceodentata* (survival and biomass endpoints), *Panope generosa* (survival), and *Dendraster excentricus* (survival).

1.11.6.3 *Leptocheirus plumulosus* was as sensitive as the freshwater amphipod *Hyaella azteca* to an artificially created gradient of sediment contamination when the latter was acclimated to oligohaline salinity (that is, 6 ‰; McGee et al., 1993 (19)). DeWitt et al. (1992b (20)) compared the sensitivity of *L. plumulosus* with three other amphipod species, two mollusks, and one polychaete to highly contaminated sediment collected from Baltimore Harbor, MD, that was serially diluted with clean sediment. *Leptocheirus plumulosus* was more sensitive than the amphipods *Hyaella azteca* and *Lepidactylus dytiscus* and exhibited equal sensitivity with *E. estuarius*. Schlekot et al. (1995 (21)) describe the results of an interlaboratory comparison of 10-d tests with *A. abdita*, *L. plumulosus* and *E. estuarius* using dilutions of sediments collected from Black Rock Harbor, CT. There was strong agreement among species and laboratories in the ranking of sediment toxicity and the ability to discriminate between toxic and non-toxic sediments.

1.11.6.4 Hartwell et al. (2000 (22)) evaluated the response of *Leptocheirus plumulosus* (10-d survival or growth) to the response

of the amphipod *Lepidactylus dytiscus* (10-d survival or growth), the polychaete *Streblospio benedicti* (10-d survival or growth), and lettuce germination (*Lactuca sativa* in 3-d exposure) and observed that *L. plumulosus* was relatively insensitive compared to the response of either *L. dytiscus* or *S. benedicti* in exposures to 4 sediments with elevated metal concentrations.

1.11.6.5 Ammonia is a naturally occurring compound in marine sediment that results from the degradation of organic debris. Interstitial ammonia concentrations in test sediment can range from <1 mg/L to in excess of 400 mg/L (Word et al., 1997 (23)). Some benthic infauna show toxicity to ammonia at concentrations of about 20 mg/L (Kohn et al., 1994 (24)). Based on water-only and spiked-sediment experiments with ammonia, threshold limits for test initiation and termination have been established for the *L. plumulosus* chronic test. Smaller (younger) individuals are more sensitive to ammonia than larger (older) individuals (DeWitt et al., 1997a (7),b (25)). Results of a 28-d test indicated that neonates can tolerate very high levels of pore-water ammonia (>300 mg/L total ammonia) for short periods of time with no apparent long-term effects (Moore et al., 1997 (26)). It is not surprising *L. plumulosus* has a high tolerance for ammonia given that these amphipods are often found in organic rich sediments in which diagenesis can result in elevated pore-water ammonia concentrations. Insensitivity to ammonia by *L. plumulosus* should not be construed as an indicator of the sensitivity of the *L. plumulosus* sediment toxicity test to other chemicals of concern.

1.11.7 Limited comparative data is available for concurrent water-only exposures of all four species in single-chemical tests. Studies that do exist generally show that no one species is consistently the most sensitive.

1.11.7.1 The relative sensitivity of the four amphipod species to ammonia was determined in ten-d water only toxicity tests in order to aid interpretation of results of tests on sediments where this toxicant is present (USEPA 1994a (1)). These tests were static exposures that were generally conducted under conditions (for example, salinity, photoperiod) similar to those used for standard 10-d sediment tests. Departures from standard conditions included the absence of sediment and a test temperature of 20°C for *L. plumulosus*, rather than 25°C as dictated in this standard. Sensitivity to total ammonia increased with increasing pH for all four species. The rank sensitivity was *R. abronius* = *A. abdita* > *E. estuarius* > *L. plumulosus*. A similar study by Kohn et al. (1994 (24)) showed a similar but slightly different relative sensitivity to ammonia with *A. abdita* > *R. abronius* = *L. plumulosus* > *E. estuarius*.

1.11.7.2 Cadmium chloride has been a common reference toxicant for all four species in 4-d exposures. DeWitt et al. (1992a (6)) reports the rank sensitivity as *R. abronius* > *A. abdita* > *L. plumulosus* > *E. estuarius* at a common temperature and salinity of 15°C and 28 ‰. A series of 4-d exposures to cadmium that were conducted at species-specific temperatures and salinities showed the following rank sensitivity: *A. abdita* = *L. plumulosus* = *R. abronius* > *E. estuarius* (USEPA 1994a (1), USEPA-USACE 1999).

1.11.7.3 Relative species sensitivity frequently varies among contaminants; consequently, a battery of tests including organisms representing different trophic levels may be needed to assess sediment quality (Craig, 1984 (27); Williams et al. 1986 (15); Long et al., 1990 (28); Ingersoll et al., 1990 (29); Burton and Ingersoll, 1994 (31)). For example, Reish (1988 (32)) reported the relative toxicity of six metals (arsenic, cadmium, chromium, copper, mercury, and zinc) to crustaceans, polychaetes, pelecypods, and fishes and concluded that no one species or group of test organisms was the most sensitive to all of the metals.

1.11.8 The sensitivity of an organism is related to route of exposure and biochemical response to contaminants. Sediment-dwelling organisms can receive exposure from three primary sources: interstitial water, sediment particles, and overlying water. Food type, feeding rate, assimilation efficiency, and clearance rate will control the dose of contaminants from sediment. Benthic invertebrates often selectively consume different particle sizes (Harkey et al. 1994 (33)) or particles with higher organic carbon concentrations which may have higher contaminant concentrations. Grazers and other collector-gatherers that feed on aufwuchs and detritus may receive most of their body burden directly from materials attached to sediment or from actual sediment ingestion. In some amphipods (Landrum, 1989 (34)) and clams (Boese et al., 1990 (35)) uptake through the gut can exceed uptake across the gills for conducting certain hydrophobic compounds. Organisms in direct contact with sediment may also accumulate contaminants by direct adsorption to the body wall or by absorption through the integument (Knezovich et al. 1987 (36)).

1.11.9 Despite the potential complexities in estimating the dose that an animal receives from sediment, the toxicity and bioaccumulation of many contaminants in sediment such as Kepone®, fluoranthene, organochlorines, and metals have been correlated with either the concentration of these chemicals in interstitial water or in the case of non-ionic organic chemicals, concentrations in sediment on an organic carbon normalized basis (Di Toro et al. 1990 (37); Di Toro et al. 1991 (38)). The relative importance of whole sediment and interstitial water routes of exposure depends on the test organism and the specific contaminant (Knezovich et al. 1987 (36)). Because benthic communities contain a diversity of organisms, many combinations of exposure routes may be important. Therefore, behavior and feeding habits of a test organism can influence its ability to accumulate contaminants from sediment and should be considered when selecting test organisms for sediment testing.

1.11.10 The use of *A. abdita*, *E. estuarius*, *R. abronius*, and *L. plumulosus* in laboratory toxicity studies has been field validated with natural populations of benthic organisms (Swartz et al. 1994 (39) and Anderson et al. 2001 (40) for *E. estuarius*, Swartz et al. 1982 and Anderson et al. 2001 (40) for *R. abronius*, McGee et al. 1999 (41) and McGee and Fisher 1999 (42) for *L. plumulosus*).

1.11.10.1 Data from USEPA Office of Research and Development's Environmental Monitoring and Assessment program were examined to evaluate the relationship between survival of *Ampelisca abdita* in sediment toxicity tests and the presence of amphipods, particularly ampeliscids, in field samples. Over 200 sediment samples from two years of sampling in the Virginian Province (Cape Cod, MA, to Cape Henry, VA) were available for comparing synchronous measurements of *A. abdita* survival in toxicity tests to benthic community enumeration. Although species of this genus were among the more frequently occurring taxa

in these samples, ampeliscids were totally absent from stations that exhibited *A. abdita* test survival <60 % of that in control samples. Additionally, ampeliscids were found in very low densities at stations with amphipod test survival between 60 and 80 % (USEPA 1994a (1)). These data indicate that tests with this species are predictive of contaminant effects on sensitive species under natural conditions.

1.11.10.2 Swartz et al. (1982 (43)) compared sensitivity of *R. abronius* to sediment collected from sites in Commencement Bay, WA, to benthic community structure at each site. Mortality of *R. abronius* was negatively correlated with amphipod density, and storage (Section 10.4); 2. phoxocephalid amphipods were ubiquitously absent from the most contaminated areas.

1.11.10.3 Sediment toxicity to amphipods in 10-d toxicity tests, field contamination, and field abundance of benthic amphipods were examined along a sediment spiking (Section 10.6); 3. collection, handling, contamination gradient of DDT (Swartz et al. 1994 (39)). Survival of *E. estuarius* and culturing *R. abronius* in laboratory toxicity tests was positively correlated to abundance of amphipods (Section 11.4); in the field and 4. statistical analyses (Section 16). USEPA-USACE (1999) along with the survival of *H. azteca*, was negatively correlated to DDT concentrations. The threshold for 10-d sediment toxicity in laboratory studies was about 300 ug DDT (+metabolites)/g organic carbon. The threshold for abundance of amphipods in the field was about 100 ng DDT (+metabolites)/g organic carbon. Therefore, correlations between toxicity, contamination, and biology indicate that acute 10-d sediment toxicity tests can provide reliable evidence of biologically adverse sediment contamination in the field.

1.11.10.4 As part of a comprehensive sediment quality assessment in Baltimore Harbor, MD, McGee et al. (1999 (41)) conducted 10-d toxicity tests with *L. plumulosus*. Negative relationships were detected between amphipod survival and concentrations of select sediment-associated contaminants, whereas a very strong positive association existed between survival in laboratory exposures and field density of *L. plumulosus* at test sites. A field validation study of the 10- and 28-d *L. plumulosus* tests by McGee and Fisher (1999 (42)) in Baltimore Harbor, also indicated good agreement between acute toxicity, sediment associated contaminants and responses of the *in situ* benthic community. In this study, the chronic 28-d test was less sensitive to sediment contamination than the acute 10-d test; however, the feeding regime used in this evaluation is different than the one currently recommended in Annex A2 and may have influenced the test results. Field validation studies with the revised 28-d test outlined in Annex A2 have not been conducted.

1.12 Chronic Sediment Methods with *Leptocheirus plumulosus*:

1.12.1 Most standard whole sediment toxicity tests have been developed to produce a method lethality endpoint (survival/mortality) with potential for conduction 28-d a sublethal endpoint (reburial) in some species (USEPA 1994a (1), USEPA-USACE 2001 (2)). Methods that measure sublethal effects have not been available or have not been routinely used to evaluate sediment toxicity in marine or estuarine sediments (Scott and Redmond, 1989 (12); Green and Chandler, 1996 (44); Levin et al., 1996 (45); Ciarelli et al., 1998 (46); Meador and Rice, 2001 (47)). Most assessments of contaminated sediment rely on short-term lethality tests (for example, ≤ 10 d; USEPA-USACE, 1991 (48); 1998 (49)). Short-term lethality tests are useful in identifying "hot spots" of sediment contamination, but might not be sensitive enough to evaluate moderately contaminated areas. However, sediment quality assessments using sublethal responses of benthic organisms, such as effects on growth and reproduction, have been used to successfully evaluate moderately contaminated areas (Ingersoll et al., 1998 (50); Kemble et al., 1994 (51); McGee et al., 1995 (52); Scott, 1989 (53)). The 28-d toxicity test with the amphipod *Leptocheirus plumulosus*. Endpoints measured in this 28-d test include survival, growth, has two sublethal endpoints: growth and reproduction.

1.3 Modifications of these procedures reproduction. These sublethal endpoints have potential to exhibit a toxic response from chemicals that otherwise might be appropriate for other sediment toxicity test procedures such as flow-through not cause acute effects or partial life-cycle tests. Methods outlined significant mortality in this guide should a test. Sublethal response to chronic exposure is also be useful valuable for conducting sediment toxicity tests with other aquatic taxa, although modifications might be necessary. Other test organisms might include other species population modeling of amphipods, other crustaceans, polychaetes, and bivalves.

1.4 Other modifications of these procedures might contaminant effects. These data can be justified by special needs or circumstances. Although using appropriate procedures is more important than following prescribed procedures, results used for population-level risk assessments of tests conducted using unusual procedures are not likely to be comparable to results benthic pollutant effects.

1.12.2 An evaluation of many other tests. Comparisons the distribution of results obtained using modified and unmodified versions *L. plumulosus* in Chesapeake Bay indicates that its distribution is negatively correlated with the degree of these procedures might provide useful information concerning new concepts and procedures for conducting sediment tests with infaunal organisms.

1.5 These procedures are applicable to sediments containing most chemicals, either individually or in formulations, commercial products, contamination (Pfitzenmeyer, 1975 (54); Reinharz, 1981 (55)). A field validation study of the 10- and known or unknown mixtures. With appropriate modifications these procedures can be used to conduct sediment toxicity 28-d *L. plumulosus* tests on factors such as temperature, salinity, dissolved oxygen, by McGee and natural Fisher (1999 (42)) in Baltimore Harbor, indicated good agreement between acute toxicity, sediment characteristics (for example, particle size distribution, organic carbon content, total solids). These methods can also be used to conduct bioconcentration tests associated contaminants and responses of the *in situ* tests, and benthic community. In this study, the chronic 28-d test was less sensitive to assess sediment contamination than the toxicity of potentially contaminated field sediments, or of such materials as sewage sludge, oils, particulate matter, acute 10-d test and solutions of toxicants added to sediments. A median lethal concentration (LC50) or median sublethal effect concentration

(EC50) of toxicants or of highly contaminated therefore had a poorer association between sediment mixed into uncontaminated sediment can contaminants and benthic community health. It should be determined. Materials either adhering to sediment particles or dissolved noted that the feeding regime used in interstitial water can be tested.

1.6 Results of short-term toxicity tests this evaluation is different than the one currently recommended in Annex A2 and may have influenced the test results. Field validation studies with the revised 28-d test materials experimentally added to sediments may be reported have not been conducted.

1.13 *Limitations*—While some safety considerations are included in terms this standard, it is beyond the scope of an LC50, and sometimes an EC50 where “concentration” refers this standard to dry or wet weight concentration in sediment. Results of a field survey with single samples encompass all safety requirements necessary to determine a spatial or temporal distribution of conduct sediment toxicity may be reported in terms of percent mortality (see Section 16). Field surveys can be designed to provide either a *qualitative* reconnaissance of the distribution of sediment toxicity or a *quantitative* statistical comparison of toxicity among stations.

1.7 This guide tests.

1.14 This standard is arranged as follows:

Referenced Documents	2
Terminology	3
Summary of Guide	4
Summary of Standard	4
Significance and Use	5
Interferences	6
Hazards	7
Reagents and Materials	7
Apparatus	8
Hazards	8
— Facilities	-8.1
Facilities, Equipment, and Supplies	-8.1
— Construction Materials	-8.2
— Test Chambers	-8.3
— Cleaning	-8.4
— Acceptability	-8.5
Toxicity Test Water	9
— General Requirements	-9.1
— Source	-9.2
— Preparation	-9.3
— Characterization	-9.4
Test and Control Sediments	10
Test Sample Collection, Storage, Manipulation, and Characterization	10
— General	40.1
— Characterization	40.2
— Control Sediment	40.3
— Field-Collected Test Sediment	40.4
— Reference Sediment	40.5
— Laboratory-Spiked Test Sediment	40.6
— Test Concentration(s)	40.7
— Addition of Test Material to Sediment	40.8
	Section
Quality Assurance and Quality Control	Section
Test Organisms	11
— Species	41.1
— Age	41.2
— Source	41.3
— Collection and Handling	41.4
— Collection, Culturing, and Maintaining	41.4
Test Organisms	41.5
— Quality	41.5
Experimental Design	12
— Controls	42.2
Calculation	42.2
— Field Survey Design	42.3
— Laboratory Experiments	42.4
Procedure	13
— Dissolved Oxygen	43.1
Report	43.1
— Temperature	43.2
— Salinity	43.3
— Light	43.4
— Feeding	43.5
— Beginning the Test	43.6
— Duration of Test	43.7
— Biological Data	43.8
— Other Measurements	43.9

Analytical Methodology	14
Acceptability of Test	45
Precision and Bias	15
Interpretation of Results	46
Keywords	16
Report	47
Keywords	48
Annexes	
— Annex A1— <i>Rhepoxynius abronius</i>	
A1. Procedure For Conducting A 10-d Sediment Survival Test With the Amphipods <i>Ampelisca abdita</i> , <i>Eohaustorius estuarius</i> , <i>Leptocheirus plumulosus</i> , or <i>Rhepoxynius abronius</i>	Annex A1
— Annex A2— <i>Eohaustorius estuarius</i>	
A2. Procedure For Conducting A <i>Leptocheirus plumulosus</i> 28-d Sediment For Measuring Sublethal Effects of Sediment-Associated Contaminants.	Annex A2
— Annex A3— <i>Ampelisca abdita</i>	
	— Annex
	A4— <i>Grandidierella japonica</i>
	— Annex
	A4— <i>Grandidierella japonica</i>
References	
	— Annex A5— <i>Leptocheirus plumulosus</i>

1.8 The values stated in SI units are to be regarded as standard.

1.9 This standard does not purport to address all of the safety problems, concerns, if any, associated with its use. It is the responsibility of the user of this standard to establish appropriate safety and health practices and determine the applicability of regulatory limitations prior to use. While some safety considerations are presented in this guide, it is beyond the scope of this guide to encompass all safety requirements necessary to conduct sediment toxicity tests. Specific hazard statements are given in Section 8.

TABLE 1 Rating of Selection Criteria for Estuarine or Marine Amphipod Sediment Toxicity Testing
 A “+” or “-” Rating Indicates a Positive or Negative Contribution

Criteria	<i>Ampelisca abdita</i>	<i>Eohaustorius estuarius</i>	<i>Leptocheirus plumulosus</i>	<i>Rhepoxynius abronius</i>
Relative sensitivity toxicity data base	Chemical +	Amount +		
	NaF	3 mg		
	+	+		
Round-robin studies conducted	SrCl ₂ ·6H ₂ O	20 mg		
	+	20 mg		
	H ₃ BO ₃	30 mg		
	+	+		
Contact with sediment	KBr	400 mg		
	+	400 mg		
	KCl	700 mg		
	+	+		
Laboratory culture	CaCl ₂ ·2H ₂ O	1.47 g		
	+/-	1.47 g		
	Na ₂ SO ₄	4.00 g		
	+	-		
Taxonomic identification	MgCl ₂ ·6H ₂ O	+	+	+
	+	+	+	+
Ecological importance	+	+	+	+
	NaCl	PAC	ATL	PAC
Geographical distribution	ATL, PAC, GOM	PAC	ATL	PAC
Sediment physicochemical tolerance	+	+	+	+
	+	+	+	+
	Na ₂ S	SiO ₂ ·9H ₂ O	20 mg+	+
Response confirmed with benthos populations	+	+	+	+
	NaHCO ₃	+	+	+
Peer reviewed	+	+	+	+
200 mg monitored	Survival	Survival, reburial	Survival	Survival, reburial
Endpoints monitored	Survival	Survival, reburial	Survival	Survival, reburial

^AIf the resulting solution is diluted to 1 L, the salinity should be 34 ± 0.5 g/kg and the pH 8. (20 ± 0.2). The desired test salinity is attained by dilution at time of use. The reconstituted salt water should be stripped of trace metals (1540).

ATL = Atlantic Coast, PAC = Pacific Coast, GOM = Gulf of Mexico

2. Referenced Documents

2.1 ASTM Standards:⁵

- D 1129 Terminology Relating to Water
- D 3976 Practice 4387 Guide for Preparation of Sediment Samples Selecting Grab Sampling Devices for Chemical Analysis⁶ Collecting Benthic Macroinvertebrates
- D 4447 Guide for the Disposal of Laboratory Chemicals and Samples
- E 380²⁹ Practice for Use of the International System of Units (SI) (the Modernized Metric System)⁶ Using Significant Digits in Test Data to Determine Conformance with Specifications
- E 729 Guide 105 Practice for Conducting Acute Toxicity Tests with Fishes, Macroinvertebrates and Amphibians⁶ Probability Sampling of Materials
- E 943 Terminology Relating 122 Practice for Choice of Sampling Size to Biological Effects and Environmental Fate⁶ Estimate a Measure of Quality for a Lot or Process
- E 1023 Guide 141 Practice for Acceptance of Evidence Based on Results of Probability Sampling
- E 177 Practice for Use of the Hazard Terms Precision and Bias in ASTM Test Methods
- E 178 Practice for Dealing with Outlying Observations
- E 380 Practice for Use of the International System of Units (SI) (The Modernized Metric System)
- E 456 Terminology Relating to Quality and Statistics
- E 691 Practice for Conducting an Interlaboratory Study to Determine Precision of a Test Method
- E 729 Guide for Conducting Acute Toxicity Tests with Fishes, Macroinvertebrates, and Amphibians
- E 943 Terminology Relating to Biological Effects and Environmental Fate
- E 1241 Guide for Conducting Early Life-Stage Toxicity Tests with Fishes
- E 1325 Terminology Relating to Design of Experiments
- E 1391 Guide for Collection, Storage, Characterization, and Manipulation of Sediments for Toxicological Testing
- E 1402 Terminology Relating to Sampling
- E 1525 Guide for Designing Biological Tests with Sediments
- E 1611 Guide for Conducting Sediment Toxicity Tests with Polychaetous Annelids
- E 1688 Guide for Determination of the Bioaccumulation of Sediment-Associated Contaminants by Benthic Invertebrates
- E 1706 Test Method for Measuring the Toxicity of Sediment-Associated Contaminants with Freshwater Invertebrates
- E 1847 Practice for Statistical Analysis of Toxicity Tests Conducted Under ASTM Guidelines
- E 1850 Guide for Section of Resident Species at Test Organisms for Aquatic and Their Uses⁶ Sediment Tests

3. Terminology

3.1 Definitions of Terms Specific to This Standard:

3.1.1 The term “*sediment*” is used here to denote a naturally occurring particulate material that has been transported and deposited at the bottom of a body of water. The procedures described can also be applied using an experimentally prepared substrate within which the amphipods can burrow.

3.1.1.1 *clean sediment*—denotes sediment that does not contain concentrations of toxicants that cause apparent stress to the test organisms or reduce their survival.

3.1.1.2 *solid-phase sediment*—distinguished from elutriates and resuspended sediments in that the whole, intact sediment is used to expose the organisms, not a form or derivative of the sediment.

3.1.2 *toxicity*—the property of a material or combination of materials, to adversely affect organisms (see Terminology E 943).

3.1.3 *exposure*—contact with a chemical or physical agent (see Terminology E 943).

3.1.4 *interstitial water*—the water within a wet sediment that surrounds the sediment particles. The amount of interstitial water in sediment is expressed as the percent ratio of the weight of the water in the sediment to that of the wet sediment.

3.1.5 *overlying water*—the water that is added to the test chamber over the solid phase of the sediment in a toxicity test.

3.1.6 *spiking of sediment*, refers to the experimental addition of a test material such as a chemical or mixture of chemicals, sewage sludge, oil, particulate matter, or highly contaminated sediment to a clean negative control or reference sediment to determine the toxicity of the material added. After the test material is added, sometimes with a solvent carrier, the sediment is thoroughly mixed to evenly distribute the test material throughout the sediment.

3.1.7 The LC50 is the statistically or graphically derived best estimate of the concentration of test material added to or contained in sediment that is expected to be lethal to 50 % of the test organisms under specified conditions within the test period (see Terminology E 943).

3.1.8 The EC50 is the statistically or graphically estimated concentration of test material in sediment that is expected to cause a measured sublethal effect (for example the inability of amphipods to rebury in clean sediment at the end of the test period), in 50 % of the test organisms under specified conditions (see Terminology E 943).

3.1.9 The

3.1 The words “must,” “should,” “may,” “can,” and “might” have very specific meanings in this guide.

3.1.9.1 “**Must**” standard. “**Must**” is used to express an absolute requirement, that is, to state that the a test ought to be designed

to satisfy the specified conditions, unless the purpose of the test requires a different design. “Must” is ~~only~~ used only in connection with the factors that ~~directly~~ relate directly to the acceptability of the test (see Section 15).

3.1.9.2 “Should” a test. “Should” is used to state that the specified condition is recommended and ought to be met if possible. Although the violation of one “should” is rarely a serious matter, violation of several will often render the results questionable. Terms such as “is desirable,” “is often desirable,” and “might be desirable” are used in connection with less important factors.

3.1.9.3 “May” factors. “May” is used to mean “is (are) allowed to,” “can” is used to mean “is (are) able to,” and “might” is used to mean “could possibly.” Thus, the classic distinction between “may” and “can” is preserved, and “might” is never used as a synonym for either “may” or “can.”

3.2 *Definitions*—For definitions of other terms used in this ~~guide~~, test method, refer to Guides E 729 and E 1241 and Terminology D 1129, Guide E 729, Terminology E 943, E 943 and Guide E 1023, D 1129. For an explanation of units and symbols, refer to Practice E 380.

3.3 *Definitions of Terms Specific to This Standard:*

3.3.1 *clean, n*—denotes a sediment or water that does not contain concentrations of test materials which cause apparent stress to the test organisms or reduce their survival.

3.3.2 *concentration, n*—the ratio of weight or volume of test material(s) to the weight or volume of sediment.

3.3.3 *contaminated sediment, n*—sediment containing chemical substances at concentrations that pose a known or suspected threat to environmental or human health.

3.3.4 *control sediment, n*—a sediment that is essentially free of contaminants and is used routinely to assess the acceptability of a test. Any contaminants in control sediment may originate from the global spread of pollutants and does not reflect any substantial input from local or non-point sources. Comparing test sediments to control sediments is a measure of the toxicity of a test sediment beyond inevitable background contamination.

3.3.5 *EC50, n*—a statistically or graphically estimated concentration that is expected to cause one or more specified effects in 50 % of a group of organisms under specified conditions.

3.3.6 *formulated sediment, n*—mixtures of materials used to mimic the physical components of a natural sediment.

3.3.7 *IC50, n*—a point estimate of the toxicant concentration that would cause a 50 % reduction in a non-quantal measurement such as fecundity or growth.

3.3.8 *interstitial water or pore water, n*— water occupying space between sediment or soil particles.

3.3.9 *LC50, n*—a statistically or graphically estimated concentration that is expected to be lethal to 50 % of a group of organisms under specified conditions.

3.3.10 *lowest-observable-effect concentration (LOEC), n*—in a toxicity test, the lowest tested concentration of a material at which organisms were adversely affected compared to control organisms as determined by statistical hypothesis tests—should be accompanied by a description of the statistical tests and alternative hypotheses, levels of significance, and measures of performance, for example, survival, growth, reproduction, or development—and must be above any other concentration not producing statistically significant adverse effects.

3.3.11 *no-observable-effect concentration (NOEC), n*—in a toxicity test, the highest tested concentration of a material at which organisms did as well as control organisms as determined by statistical hypothesis tests—should be accompanied by a description of the statistical tests and alternative hypotheses, levels of significance, and measures of performance, for example, survival, growth, reproduction, or development—and must be below any other concentration producing statistically significant adverse effects.

3.3.12 *overlying water, n*—the water placed over sediment in a test chamber during a test.

3.3.13 *reference sediment, n*—a whole sediment near an area of concern used to assess sediment conditions exclusive of material(s) of interest. The reference sediment may be used as an indicator of localized sediment conditions exclusive of the specific pollutant input of concern. Such sediment would be collected near the site of concern and would represent the background conditions resulting from any localized pollutant inputs as well as global pollutant input. This is the manner in which reference sediment is used in dredge material evaluations.

3.3.14 *reference-toxicity test, n*—a test conducted with reagent-grade reference chemical to assess the sensitivity of the test organisms. Deviations outside an established normal range may indicate a change in the sensitivity of the test organism population. Reference-toxicity tests are most often performed in the absence of sediment.

3.3.15 *sediment, n*—particulate material that usually lies below water. Formulated particulate material that is intended to lie below water in a test.

3.3.16 *spiked sediment, n*—a sediment to which a material has been added for experimental purposes.

3.3.17 *whole sediment, n*—sediment and associated pore water which have had minimal manipulation. The term bulk sediment has been used synonymously with whole sediment.

4. Summary of Guide

4.1 The relative Standard

4.1 *Method Description*—Procedures are described for testing estuarine or marine amphipod crustaceans in the 10-d laboratory exposures to evaluate the toxicity of contaminants associated with whole sediments. Sediments may be collected from the field or spiked with compounds in the laboratory. A toxicity method is outlined for four species of estuarine sediments can be determined

through or marine sediment-burrowing amphipods found within United States coastal waters. The species are *Ampelisca abdita*, a 10-d marine species that inhabits marine and mesohaline portions of the Atlantic coast, the Gulf of Mexico, and San Francisco Bay; *Eohaustorius estuarius*, a Pacific coast estuarine species; *Leptocheirus plumulosus*, an Atlantic coast estuarine species; and *Rhepoxynius abronius*, a Pacific coast marine species. Generally, the method described may be applied to all four species, although acclimation procedures and some test-w conditions (that is, temperature and salinity) will be species-specific (Sections 10 and 11). The toxicity test is conducted in 1-L glass chambers containing 175 mL of sediment and 775 mL of overlying seawater. Exposure is static (that is, water is not renewed), and the animals are not fed over the 10-d exposure period. The endpoint in the toxicity test is survival with reburial of surviving amphipods as an additional measurement that can be used as an endpoint for some of the test species (for *R. abronius* and *E. estuarius*). Performance criteria established for this test include the average survival of amphipods in negative control treatment must be greater than or equal to 90 %. Procedures are described for use with sediments with pore-water salinity ranging from >0 ‰ to fully marine.

4.2 A procedure is also described for determining the chronic toxicity of contaminants associated with whole sediments with the amphipod *Leptocheirus plumulosus* in laboratory exposures (USEPA-USACE 2001 (2)). The toxicity test is conducted for 28 d in 1-L glass chambers containing 175 mL of sediment and about 775 mL of overlying seawater. Four hundred millilitres of overlying water is renewed three times per week, at which time test-chambers. Mortality organisms are fed. Tests are initiated with neonate amphipods that mature and sublethal effects such reproduce during the 28-d test period. The endpoints in the 28-d toxicity test are survival, growth rate, and reproduction of amphipods. Survival is calculated as the percentage of newly born (neonate) amphipods at test initiation that survive as adults at test termination. Growth rate is calculated as the mean dry weight gain per day per adult amphipod surviving at test termination. Reproduction is calculated as the number of offspring per surviving adult. This test is applicable for use with sediment having pore-water salinity ranging from 1 ‰ to 35 ‰. Typically, endpoint selection for new toxicity tests is generally guided by methodologies for related toxicity tests (Gray et al., 1998 (56)). Sediment toxicity tests using macroinvertebrates often incorporate survival and growth endpoints (Ingersoll, 1995 (57)). Gray et al. (1998 (56)) recommend optimal endpoint measures for the *L. plumulosus* sediment toxicity test based on four criteria: relevance of each measure to its respective endpoint; signal-to-noise ratio (the ratio between the response to stressor and inability the normal variation in the response variable); redundancy to other measures of the same endpoint; and cost of labor, training, and equipment. Signal-to-noise ratios are independent of experiment design considerations (that is, Type I and Type II errors, and sample size) and are positively correlated with power (Gray et al., 1998 (56)).

4.3 *Experimental Design*—The following section is a general summary of experimental design. See Section 13 for additional detail.

4.3.1 *Control and Reference Sediment* :

4.3.1.1 Sediment tests include a control sediment (sometimes called a negative control). A control sediment is a sediment that is essentially free of contaminants and is used routinely to assess the acceptability of a test and is not necessarily collected near the site of concern. Any contaminants in clean control sediment are determined after exposure thought to originate from the global spread of pollutants and do not reflect any substantial inputs from local or non-point sources Ankley and Thomas, 1992 (58). Comparing test sediments to control sediments is a specific number (usually 20) measure of amphipods to the toxicity of a test sediment beyond inevitable background contamination and organism health Ankley and Thomas, 1992 (58). A control sediment provides a measure of test sediment. Response acceptability, evidence of test organism health, and a basis for interpreting data obtained from the amphipods test sediments. A reference sediment is collected near an area of concern and is used to assess sediment conditions exclusive of material(s) of interest. Testing a reference sediment provides a site-specific basis for evaluating toxicity.

4.3.1.2 In general, the performance of test organisms in the negative control is used to judge the acceptability of a test, and either the negative control or reference sediment may be used to evaluate performance in the experimental treatments, depending on the purpose of the study. Any study in which organisms in the negative control do not meet performance criteria must be considered questionable because it suggests that adverse factors affected the response of test organisms. Key to avoiding this situation is using only control sediments that have a demonstrated record of performance using the same test procedure. This includes testing of new collections from sediment sources that have previously provided suitable control sediment.

4.3.1.3 Because of the uncertainties introduced by poor performance in the negative control, such studies should be repeated to insure accurate results. However, the scope or sampling associated with some studies may make it difficult or impossible to repeat a study. Some researchers have reported cases where performance in the negative control is poor, but performance criteria are met in reference sediment included in the study design. In these cases, it might be reasonable to infer that other samples that show good performance are probably not toxic; however, any samples showing poor performance should not be judged to have shown toxicity, since it is unknown whether the adverse factors that caused poor control performance might have also caused poor performance in the test treatments.

4.3.1.4 Natural physico-chemical characteristics such as sediment texture may influence the response of test organisms (59). The physico-chemical characteristics of test sediment need to provide (a) be within the tolerance limits of the test organism. Ideally, the limits of a test organism should be determined in advance; however, controls for factors including grain size and organic carbon can be evaluated if the limits are exceeded in a test sediment. See section 12.1 and Annex A1 and Annex A2 for information on physico-chemical requirements of test organisms. If the acceptability physico-chemical characteristics of a test sediment exceed

the tolerance range of the test by providing evidence organism, a control sediment encompassing these characteristics can be evaluated. The effects of sediment characteristics on the results of sediment tests can be addressed with regression equations Dewitt et al. 1988, (59,) Ankley et al., 1994(60). The use of formulated sediment can also be used to evaluate physico-chemical characteristics of sediment on test organisms Walsh et al., 1991 (61) Suedel and Rodgers, 1994, (62) Kembel et al.,(63) USEPA, 2000,(64 (62), section 7.2 and Guide E 1391).

4.3.2 The experimental design depends on the purpose of the study. Variables that need to be considered include the number and type of control sediments, the number of treatments and replicates, and water quality characteristics. For instance, the purpose of the test organisms; study might be to determine a specific endpoint such as an LC50 and may include a control sediment, a positive control, a solvent control, and several concentrations of sediment spiked with a chemical (see section 10.3.2). A useful summary of field sampling design is presented by Green, 1979 (65). See Section 13 for additional guidance on experimental design and statistics.

4.3.2.1 The purpose of the study might be to determine if field-collected sediments are toxic and may include controls, reference sediments, and test sediments. Controls are used to evaluate the acceptability of the overlying water; test conditions (Table A1.3 in Annex A1 and handling procedures, etc., Table A2.3 in Annex A2) and (b) the might include a control sediment or a formulated sediment (section 7.2). Testing a reference sediment provides a site-specific basis for interpreting data obtained from evaluating toxicity of the test sediments.

4.1.1 The toxicity sediments. Comparisons of f test sediments to multiple reference or control sediments representative of the physical characteristics of the test sediment (that is, grain size, organic carbon) may be useful in these evaluations. A summary of field sampling design is indicated presented by Green, 1979 (65) . See Section 13 for additional guidance on experimental design and statistics.

4.3.2.2 If the percent mortality purpose of amphipods exposed the study is to that sediment compared conduct a reconnaissance field survey to identify sites for further investigation, the experimental design might include only one sampled from each site to control sediment. allow for sampling a larger area. The toxicity lack of f replication at a site usually precludes statistical comparisons (for example, analysis of variance (ANOVA)), but these surveys can be used to identify sites for further study or may also be assessed by testing dilutions evaluated using regression techniques.

4.3.2.3 In other instances, the purpose of the study might be to conduct a highly toxic quantitative sediment survey of chemistry and toxicity to determine statistically significant differences between effects among control and test sediments from several sites. The number of replicates/site should be based on the need for sensitivity or power (see Section 13). In a quantitative survey, field replicates (separate samples from different grabs collected at the same site) would need to be taken at each site. Chemical and physical characterizations of each of these grabs would be required for each of these field replicates used in sediment testing. Separate subsamples might be used to determine within-sample variability or for comparisons of test procedures (for example, comparative sensitivity among test organisms), but these subsamples cannot be considered to be true field replicates for statistical comparisons among sites.

4.3.2.4 Sediments often exhibit high spatial and temporal variability (66). Therefore, replicate samples may need to be collected to determine variance in sediment characteristics. Sediment should be collected with e as little disruption as possible; however, subsampling, compositing, or homogenization of sediment samples may be required for some experimental designs.

4.3.2.5 Site locations might be distributed along a known pollution gradient, in relation to obtain information on the toxicity boundary of a disposal site, or at sites identified as being contaminated in a reconnaissance survey. Comparisons can be made in both space and time. In pre-dredging studies, a sampling design can be prepared to assess the contamination of that sediment.

4.1.2 The toxicity samples representative of the project area to be dredged. Such a toxicant experimentally added design may include compositing cores collected to project depth from a specified dredged material management area.

4.3.2.6 The primary focus of the physical and experimental test design and statistical analysis of the data, is the experimental unit, which is defined as the smallest physical entity to which treatments can be expressed by analyzing independently assigned (Guide E 1241). Because overlying water or air cannot flow from one test chamber to another the test chamber is the experimental unit. The experimental unit is defined as the smallest physical entity to which treatments can be independently assigned and reburial data to determine an LC50 which air and an EC50 for water exchange between test chambers are kept to a minimum. Because of factors that might affect results within test chambers and results of a test, all test chambers should be treated as similarly as possible. Treatments should be randomly assigned to individual test chamber locations. Assignment of test organisms to test chambers should be impartial (Guide E 729). As the toxicant for number of test chambers/treatment increases, the duration number of degrees of freedom increases, and, therefore, the width of the confidence interval on a point estimate, such as an LC50, decreases, and the power of a significance test increases (see Section 13).

5. Significance and Use

5.1 The test procedure in this guide

5.1 General:

5.1.1 Sediment provides habitat for many aquatic organisms and is not intended to exactly simulate the exposure of benthic amphipods to contaminants under “natural” conditions, but rather to provide a conveniently rapid, standard toxicity test procedure yielding a reasonably sensitive indication major repository for many of the toxicity of more persistent chemicals that are introduced

into surface waters. In the aquatic environment, most anthropogenic chemicals and waste materials in marine including toxic organic and estuarine sediments:

5.2 Protection inorganic chemicals eventually accumulate in sediment. Mounting evidences exists of a community of environmental degradation in areas where USEPA Water Quality Criteria (WQC; Stephan et al.(67)) are not exceeded, yet organisms requires averting detrimental contaminant related effects on in or near sediments are adversely affected Chapman, 1989 (68). The WQC were developed to protect organisms in the number water column and health were not directed toward protecting organisms in sediment. Concentrations of individuals and species within that population. Sediment toxicity tests provide information on the toxicity of test materials contaminants in sediments. Protection sediment may be several orders of magnitude higher than in the most sensitive species within a community will theoretically protect the community as a whole.

5.3 Amphipods are an abundant component overlying water; however, whole sediment concentrations have not been strongly correlated to bioavailability Burton, 1991 (69). Partitioning or sorption of the soft bottom marine and estuarine benthic community. They are a principal prey of compound between water and sediment may depend on many fish, birds, factors including: aqueous solubility, pH, redox, affinity for sediment organic carbon and larger invertebrate species. Some species are predators dissolved organic carbon, grain size of smaller benthic invertebrates. Others ingest the sediment, sediment particles mineral constituents (oxides of iron, manganese, and aluminum), and the quantity of acid volatile sulfides in sediment Di Toro et al. 1991(70) Giesy et al. 1988 (71). Although certain chemicals are highly sorbed to sediment, these compounds may still be available to the biota. Chemicals in sediments may be directly exposed toxic to contaminants. Amphipods are among aquatic life or can be a source of chemicals for bioaccumulation in the first taxa food chain.

5.1.2 The objective of a sediment test is to disappear from benthic communities impacted by pollution, and have been shown determine whether chemicals in sediment are harmful to or are bioaccumulated by benthic organisms. The tests can be more sensitive used to measure interactive toxic effects of complex chemical mixtures in a sediment. Furthermore, knowledge of specific pathways of interactions among sediments than several other major taxa and test organisms is not necessary to conduct the tests Kemp et al. 1988, (72). Sediment tests can be used to: (1) determine the relationship between toxic effects and bioavailability, (2) investigate interactions among chemicals, (3) compare the sensitivities of different organisms, (4). The ecological importance of amphipods, their wide geographical distribution, ease of handling in the laboratory, and their sensitivity to contaminated sediments make them appropriate species for sediment toxicity testing.

5.4 An acute toxicity test is conducted to obtain information concerning the immediate effects on test organisms of a short-term exposure to a test material under specific experimental conditions. An acute toxicity test does not necessarily provide information about whether delayed effects will occur, although a post exposure observation period, with appropriate feeding if necessary, could provide such information.

5.5 Results of acute sediment toxicity tests can be used to predict acute effects likely to occur on aquatic organisms in field situations as a result of exposure under comparable conditions, except that (a) motile organisms might avoid exposure when possible and (b) toxicity to benthic organisms can be dependent on sediment characteristics, dynamics of equilibrium partitioning, and the route of exposure to the benthic organisms.

5.6 The amphipod sediment toxicity test might be used to determine the temporal or spatial and temporal distribution of sediment toxicity. Test methods can be used to detect horizontal and vertical gradients in toxicity. Mortality data can be used to indicate the relative toxicity of field collected sediments.

5.7 Results of acute tests with toxicants experimentally added to sediments can be used to compare the acute sensitivities of different species and the acute toxicities of different test materials, and to define the effects of various environmental factors on results of such tests.

5.8 Results of acute sediment toxicity tests are useful for studying biological availability of, and structure-activity relationships between, test materials in sediment.

5.9 Results of acute sediment toxicity tests might be an important consideration when assessing the hazards of materials to aquatic organisms (see Guide E-1023) or when deriving sediment quality criteria for aquatic organisms contamination, (5); Sediment evaluate hazards of dredged material, (6) measure toxicity as part of product licensing or safety testing, (7) rank areas for clean up, and (8) estimate the effectiveness of remediation or management practices.

5.1.3 A variety of methods have been developed for assessing the toxicity of chemicals in sediments using amphipods, midges, polychaetes, oligochaetes, mayflies, or cladocerans (Test Method E 1706, Guide E 1525, Guide E 1850; Annex A1, Annex A2; USEPA, 2000 (73), EPA 1994b, (74), Environment Canada 1997a, (75), Enviroment Canada 1997b,(76)). Several endpoints are suggested in these methods to measure potential effects of contaminants in sediment including survival, growth, behavior, or reproduction; however, survival of test organisms in 10-day exposures is the endpoint most commonly reported. These short-term exposures that only measure effects on survival can be used to identify high levels of contamination in sediments, but may not be able to identify moderate levels of contamination in sediments (USEPA USEPA, 2000 (73); Sibley et al.1996, (77); Sibley et al.1997a, (78); Sibley et al.1997b, (79); Benoit et al.1997, (80); Ingersoll et al.1998, (81)). Sublethal endpoints in sediment tests might also prove to be useful better estimates of responses of benthic communities to contaminants in making decisions regarding the field, Kembel et al. 1994 (82) . Insufficient information is available to determine if the long-term test conducted with *Leptocheirus plumulosus* (Annex A2) is more sensitive than 10-d toxicity tests conducted with this or other species.

5.1.3.1 The decision to conduct short-term or long-term toxicity tests depends on the goal of the assessment. In some instances,

sufficient information may be gained by measuring sublethal endpoints in 10-day tests. In other instances, the 10-day tests could be used to screen samples for toxicity before long-term tests are conducted. While the long-term tests are needed to determine direct effects on reproduction, measurement of growth in these toxicity tests may serve as an indirect estimate of reproductive effects of contaminants associated with sediments (Annex A1).

5.1.3.2 Use of sublethal endpoints for assessment of contaminant risk is not unique to toxicity testing with sediments. Numerous regulatory programs require the use of sublethal endpoints in the decision-making process (Pittinger and Adams, 1997, (83)) including: (1) Water Quality Criteria (and State Standards); (2) National Pollution Discharge Elimination System (NPDES) effluent monitoring (including chemical-specific limits and sublethal endpoints in toxicity tests); (3) Federal Insecticide, Rodenticide and Fungicide Act (FIFRA) and the Toxic Substances Control Act (TSCA, tiered assessment includes several sublethal endpoints with fish and aquatic invertebrates); (4) Superfund (Comprehensive Environmental Responses, Compensation and Liability Act; CERCLA); (5) Organization of Economic Cooperation and Development (OECD, sublethal toxicity testing with fish and invertebrates); (6) European Economic Community (EC, sublethal toxicity testing with fish and invertebrates); and (7) the Paris Commission (behavioral endpoints).

5.1.4 Results of toxicity tests on sediments spiked at different concentrations of chemicals can be used to establish cause and effect relationships between chemicals and biological responses. Results of toxicity tests with test materials spiked into sediments at different concentrations may be reported in terms of an LC50 (median lethal concentration), an EC50 (median effect concentration), an IC50 (inhibition concentration), or as a NOEC (no observed effect concentration) or LOEC (lowest observed effect concentration). However, spiked sediment may not be representative of chemicals associated with sediment in the field. Mixing time Stemmer et al. 1990b, (84), aging (Landrum et al. 1989, (85), Word et al. 1987, (86), Landrum et al., 1992,(87)), and the chemical form of the material can affect responses of test organisms in spiked sediment tests.

5.1.5 Evaluating effect concentrations for chemicals in sediment requires knowledge of factors controlling their bioavailability. Similar concentrations of a chemical in units of mass of chemical per mass of sediment dry weight often exhibit a range in toxicity in different sediments Di Toro et al. 1990, (88) Di Toro et al. 1991,(70). Effect concentrations of chemicals in sediment have been correlated to interstitial water concentrations, and effect concentrations in interstitial water are often similar to effect concentrations in water-only exposures. The bioavailability of nonionic organic compounds in sediment is often inversely correlated with the organic carbon concentration. Whatever the route of exposure, these correlations of effect concentrations to interstitial water concentrations indicate that predicted or measured concentrations in interstitial water can be used to quantify the exposure concentration to an organism. Therefore, information on partitioning of chemicals between solid and liquid phases of sediment is useful for establishing effect concentrations Di Toro et al. 1991, (70) .

5.1.6 Field surveys can be designed to provide either a qualitative reconnaissance of the distribution of sediment contamination or a quantitative statistical comparison of contamination among sites.

5.1.7 Surveys of sediment toxicity are usually part of more comprehensive analyses of biological, chemical, geological, and hydrographic data. Statistical correlations may be improved and sampling costs may be reduced if subsamples are taken simultaneously for sediment tests, chemical analyses, and benthic community structure.

5.1.8 Table 2 lists several approaches the USEPA has considered for the assessment of sediment quality USEPA, 1992, (89). These approaches include: (1) equilibrium partitioning, (2) tissue residues, (3) interstitial water toxicity, (4) whole-sediment toxicity and sediment-spiking tests, (5) benthic community structure, (6) effect ranges (for example, effect range median, ERM), and (7) sediment quality triad (see USEPA, 1989a, 1990a, 1990b and 1992b, (90, 91, 92, 93 and Wenning and Ingersoll (2002 (94)) for a critique of these methods). The sediment assessment approaches listed in Table 2 can be classified as numeric (for example, equilibrium partitioning), descriptive (for example, whole-sediment toxicity tests), or a combination of numeric and descriptive approaches (for example, ERM, USEPA, 1992c, (95). Numeric methods can be used to derive chemical-specific sediment quality guidelines (SQGs). Descriptive methods such as toxicity tests with field-collected sediment cannot be used alone to develop numerical SQGs for individual chemicals. Although each approach can be used to make site-specific decisions, no one single approach can adequately address sediment quality. Overall, an integration of several methods using the weight of evidence is the most desirable approach for assessing the effects of contaminants associated with sediment, (Long et al. 1991(96) MacDonald et al. 1996 (97) Ingersoll et al. 1996 (98) Ingersoll et al. 1997 (99) , Wenning and Ingersoll 2002 (94)). Hazard evaluations integrating data from laboratory exposures, chemical analyses, and benthic community assessments (the sediment quality triad) provide strong complementary evidence of the degree of pollution-induced degradation in aquatic communities (Burton, 1991 (69) , Chapman 1992, 1997 (100, 101).)

5.2 *Regulatory Applications*—Test Method E 1706 provides information on the regulatory applications of sediment toxicity tests.

5.3 *Performance-based Criteria* :

5.3.1 The USEPA Environmental Monitoring Management Council (EMMC) recommended the use of performance-based methods in developing standards, (Williams, 1993 (102). Performance-based methods were defined by EMMC as a monitoring approach which permits the use of appropriate methods that meet preestablished demonstrated performance standards (section 11.2).

5.3.2 The USEPA Office of Water, Office of Science and Technology, and Office of Research and Development held a workshop to provide an opportunity for experts in the field of sediment toxicology and staff from the USEPA Regional and Headquarters

TABLE 2 Precision of the Sediment Toxicity Test Using *Rhepoxynius abronius* in Remediation Pro-Sample Size and Rurepls (Modified from USEPA (378))

Number of Amphipods per Replicate		10		20	
Number of Replicates	δ^A	Type	δ/cB	h	h
Number of Replicates	δ^A	Type	Approach	h	h
Number of Replicates	δ^A	δ/c^B			
Numeric	Descriptive Combination				
-2	6.80	71.6			
Equilibrium Partitioning	*	*			
			A sediment quality value for a given contaminant is determined by calculating the sediment concentration of the contaminant that corresponds to an interstitial water concentration equivalent to the USEPA water-quality criterion for the contaminant.		
-4	2.66	28.0	2	8.55	45.0
Tissue Residues	*	*			45.0
					Safe sediment concentrations of specific chemicals are established by determining the sediment chemical concentration that results in acceptable tissue residues. ⁵⁵
-6	1.94	20.4	3	4.44	23.4
Interstitial Water Toxicity	*	*			23.4
					Toxicity of interstitial water is quantified and identification evaluation procedures are applied to identify and quantify chemical components responsible for sediment toxicity. ⁴⁴
-8	1.60	16.8	4	3.35	17.6
Benthic Community Structure	*	*			17.6
					Environmental degradation is measured by evaluating alterations in benthic community structure. ³⁵
40	1.38	14.5	5	2.80	14.7
Whole-sediment Toxicity And Sediment Spiking	*	*			14.7
					* Test organisms are exposed to sediments that may contain known or unknown quantities of potentially toxic chemicals. ⁸⁰
					14 At the end of a specified time period, organisms is examined in relation to response relationships can be established for test organisms to sediments that have been spiked with chemicals or mixtures.
12	1.25	13.2	6	2.45	12.9
Sediment Quality Triad	*	*			12.9
					* Sediment chemical contamination, sediment toxicity, and benthic community structure are measured on the same sediment sample. ⁴⁵
					12 Correspondence between sediment quality triad effects is used to determine sediment quality conditions of minimal, unacceptable, or unacceptable effects.
14	1.14	12.0	7	2.20	11.6
Sediment Quality Guidelines	*	*			11.6
					* The sediment concentration of contaminants associated with toxic responses measured in laboratory exposures or field assessments (that is, Apparent Effects Threshold (AET), Effect Range Median (ERM), Probable Effect Level (PEL)). ²⁰
16	1.05	11.0	8	2.02	10.6
18	0.98	10.3	9	1.89	10.0
20	0.93	9.8	10	1.76	9.3

^A δ is the difference between the survival means for which the toxicity test is 75 % certain of detecting statistical significance (P < 0.05) (31).

^B δ/c expresses the precision estimate as a percent of the normal control survival in *Rhepoxynius abronius* (c = 19.0 for n = 20; c = 9.5 for n = 10).

Program offices to discuss the development of standard freshwater, estuarine, and marine sediment testing procedures (USEPA, 1992a, 1994a (89,103)). Workgroup participants arrived at a consensus on several culturing and testing methods. In developing guidance for culturing test organisms to be included in the USEPA methods manual for sediment tests, it was agreed that no one method should be required to culture organisms. However, the consensus at the workshop was that success of a test depends on the health of the cultures. Therefore, having healthy test organisms of known quality and age for testing was determined to be the key consideration relative to culturing methods. A performance-based criteria approach was selected in USEPA, 2000 (73) as the preferred method through which individual laboratories could use unique culturing methods rather than requiring use of one culturing method.

5.3.3 This standard recommends the use of performance-based criteria to allow each laboratory to optimize culture methods and minimize effects of test organism health on the reliability and comparability of test results. See Annex A1 and Annex A2 for a listing of performance criteria for culturing or testing.

6. Interferences

6.1 ~~Du~~General Interferences:

6.1.1 An interference is a characteristic of a sediment or a test system that can potentially affect test organism response aside from those related to sediment-associated contaminants. These interferences can potentially confound interpretation of test results in two ways: (1) toxicity is observed in the limited time test sediment when contamination is low or there is more toxicity tests have been practiced, than expected, and (2) no toxicity is observed when contaminants are present at elevated concentrations or there is less toxicity than expected.

6.1.2 Because of the methodology continues heterogeneity of natural sediments, extrapolation from laboratory studies to develop the field can sometimes be difficult (Table 3; Burton, 1991 (69)). Sediment collection, handling, and evolve with time

TABLE 3 Advantages and Disadvantages for Use of Sediment Tests (Modified from Swartz (120))

<p>Advantages</p> <ul style="list-style-type: none"> —Measure bioavailable fraction of contaminant(s). —Provide a direct measure of benthic effects, assuming no field adaptation or amelioration of effects. —Limited special equipment is required. —Methods are rapid and inexpensive. —Legal and scientific precedence exist for use; ASTM standards are available. —Measure unique information relative to chemical analyses or benthic community analyses. —Tests with spiked chemicals provide data on cause-effect relationships. —Sediment-toxicity tests can be applied to all chemicals of concern. —Tests applied to field samples reflect cumulative effects of contaminants and contaminant interactions. —Toxicity tests are amenable to confirmation with natural benthos populations.
<p>Disadvantages</p> <ul style="list-style-type: none"> —Sediment collection, handling, and storage may alter bioavailability. —Spiked sediment may not be representative of field contaminated sediment. —Natural geochemical characteristics of sediment may affect the response of test organisms. —Indigenous animals may be present in field—collected sediments. —Route of exposure may be uncertain and data generated in sediment toxicity tests may be difficult to interpret if factors controlling the bioavailability of contaminants in sediment are unknown. —Tests applied to field samples may not discriminate effects of individual chemicals. —Few comparisons have been made of methods or species. —Only a few chronic methods for measuring sublethal effects have been developed or extensively evaluated. —Laboratory tests have inherent limitations in predicting ecological effects. —Tests do not directly address human health effects.

storage may alter bioavailability and research needs. Because concentration by changing the physical, chemical, or biological characteristics of the developmental nature sediment. Maintaining the integrity of a field-collected sediment during removal, transport, mixing, storage, and testing, is difficult and may complicate the interpretation of effects. See USEPA, 2000 (64) and Guide E 1391. An abundance of the same organism (McGee et al., 1999 (41)) or organisms taxonomically similar to the methods described test organism in this guide.

6.2 Results of acute the sediment toxicity tests will depend, in part, on sample may make interpretation of treatment effects difficult. In addition, the temperature, water quality, physical and chemical properties presence of predator may change the test sediment, condition outcome of a toxicity test. For example, Redmond and Scott, 1989 (104) showed that the polychaete *Nephtys incisa* can consume *Ampelisca abdita* under toxicity test organisms, exposure technique, and other factors. Factors potentially affecting results from static sediment conditions. Similarly, predatory isopods (*Cyathura polita*) have been observed to interfere in 10-d toxicity tests might include:

6.2.1 Alteration conducted with *Leptocheirus plumulosus* (Peter De Lisle, Coastal Bioanalysts, Gloucester, VA; personal communication).

6.1.2.1 Although disruptive of field natural sediment physical features, all test sediments in preparation for laboratory testing.

6.2.1.1 Maintaining the integrity of the *Leptocheirus plumulosus* 28-d sediment environment during its removal, transport, and test should be press-sieved sometime before testing and re-homogenized immediately before introduction to the laboratory test chambers if warranted (section 10.3 and Annex A2). Press-sieving is performed primarily to remove predatory organisms, large debris, organisms used in testing (McGee et al., 1999 (41)) or organisms taxonomically similar to the test species. Certain applications may recommend that sediments should not be press-sieved. Also, it may not be necessary to press-sieve sediments if previous experience has demonstrated the absence of potential interferences, including predatory or competitive organisms or large debris, or if large debris or predators can be removed with forceps or other suitable tools. The sediment environment is composed presence of a myriad an abundance of microenvironments, redox gradients, amphipods that are taxonomically similar to the test species should prompt press-sieving. This is particularly true if endemic Ampeliscidae are present and other interacting physiochemical and biological processes. Many *A. abdita* is the test species because it may be difficult to remove all of the reside ent amphipods from their tubes. If sediments are sieved, it is desirable to perform select analyses (for example, pore-water metals or DOC, AVS, TOC) on samples before and after sieving to document the influence of sieving on sediment toxicity chemistry (USEPA, 1994a (1)).

6.1.3 Depletion of aqueous and sediment-sorbed chemicals resulting from uptake by an organism or test chamber may also influence availability. In most cases, the organism is a minor sink for chemicals relative to benthic and planktonic organisms, microbial degradation, and the sediment. However, within the burrow of an organism, sediment desorption kinetics may limit uptake rates. Within minutes to hours, a major portion of the total chemical sorption. Any disruption may be inaccessible to the organisms because of this environment complicates interpretations depletion of treatment effects, causative factors, available residues. The desorption of a particular compound from sediment may range from easily reversible (labile; within minutes) to irreversible (non-labile; within days or months, Karickhoff and in-situ comparisons:

6.2.1.2 Testing Morris, 1985 (105)). Interparticle diffusion or advection and the quality and quantity of sediment organic carbon can also affect sorption kinetics.

6.1.4 Testing sediments at temperatures or salinities other than those at which they were collected different from the field might affect contaminant solubility, partitioning coefficients, and or other physical and chemical characteristics. 6.2.2 Interactions between the sediment particles, and overlying water, interstitial water, water and humic substances, and the ratio of sediment to overlying water ratio:

6.2.3 Interactions among chemicals that might may influence bioavailability (Stemmer and Burton, 1990b (84)).

6.1.5 Results of sediment tests can be present used to predict effects that may occur with aquatic organisms in test sediment.

6.2.4 Realism the field as a result of using spiked sediment (that is, whether exposure under comparable conditions. However, motile organisms might avoid exposure in the field. Photoinduced toxicity may be important for some compounds associated with sediment is at equilibrium (for example, polycyclic aromatic hydrocarbons (PAHs) Davenport and evenly mixed):

6.2.5 Photolysis and other processes degrading test chemicals:

6.2.6 Maintaining acceptable quality Spacie, 1991 (106)). However, lighting typically used to conduct laboratory tests does not include the appropriate spectrum of overlying water:

6.2.7 Excess food might change sediment partitioning ultraviolet radiation to photoactivate compounds (Oris and water quality parameters:

6.2.8 Resuspension Giesy, 1985 (107), Ankley et al. 1994b (108)), and thus laboratory tests may not account for toxicity expressed by this mode of action.

6.1.6 Natural physico-chemical characteristics such as sediment during texture may influence the toxicity test.

6.2.9 Limited opportunity for biological observations during the response of test because organisms bury in test sediment.

6.2.10 Natural geochemical properties (Dewitt et al. 1998, (59)). The physico-chemical characteristics of test sediment collected from the field that might not need to be within the tolerance limits of the test organisms:

6.2.11 Recovery organism. Ideally, the limits of the test organism should be determined in advance; however, control samples reflecting differences in factors such as grain size and organic carbon can be evaluated if the limits are exceeded in the test system.

6.2.12 Endemic sediment (section 12.1 and Annex A1 and Annex A2). The effects of sediment characteristics can also be addressed with regression equations Dewitt et al., 1998 (59) Ankley et al., 1994 (60). The use of formulated sediment can also be used to evaluate physico-chemical characteristics of sediment on test organisms which might (Walsh et al., 1991(61), Suedel and Rodgers, 1994 (62)).

6.1.7 The route of exposure may be uncertain and data from sediment tests may be difficult to interpret if factors controlling the bioavailability of chemicals in field collected sediments including (a) predators, (b) species that sediment are unknown. Whole-sediment chemical concentrations may be normalized to factors other than dry weight. For example, concentrations of nonionic organic compounds might be normalized to sediment organic-carbon content, (USEPA, 1992 (95)) and certain metals normalized to acid volatile sulfides, (DiToro, 1990, (88)). Even with the appropriate normalizing factors, determination of toxic effects from ingestion of sediment or from dissolved chemicals in the interstitially water can still be difficult, (Lamberson and Swartz, 1998 (109)).

6.1.8 The addition of food, water, or solvents to the test chambers might obscure the bioavailability of chemicals in sediment or might provide a substrate for bacterial or fungal growth. Without addition of food, the test organisms may starve during long-term exposures (Ankley et al., 1994, McNulty et al. 1999 (60, 110)). However, the addition of the food may alter the availability of the chemicals in the sediment, (Harkey et al. 1994, Wiederholm et al. 1987 (111,112)) microorganisms depending on the amount of food added, its composition (for example, bacteria, molds), total organic carbon (TOC)), and algae colonizing the chemical(s) of interest.

6.1.9 Laboratory sediment testing with field-collected sediments may be useful in estimating cumulative effects and interactions of multiple contaminants in a sample. Tests with field samples usually cannot discriminate between effects of individual chemicals. Many sediment samples contain a complex matrix of inorganic and organic chemicals with many unidentified compounds. 6.3 S The use of Toxicity Identification Evaluations (TIE) procedures including sediment tests might not with spiked chemicals may provide evidence of causal relationships and can be applicable applied to many chemicals of concern (Ankley and Thomas, 1992, (58)). Laboratory studies that test single compounds spiked into the sediment can be used to determine more directly the specific chemical vs causing a toxic response (Swartz et al. 1998 (113)).

6.1.10 Sediment spiking can also be used to investigate additive, antagonistic, or are rapidly biologically or chemically transformed. Furthermore, synergistic effects of specific chemical mixtures in a sediment sample (Swartz et al, 1998 (113)). However, spiked sediment may not be representative of contaminated sediment in the field. Mixing time (Stemmer et al. 1990a

(66)), and aging (Landrum 1999, Word et al. 1997, Landrum and Faust 1992 (85, 86, 87) of spiked sediment can affect responses of organisms.

6.1.11 Salinity of the overlying water quality might change considerably from is an additional factor that can affect the bioavailability of metals. Importantly, some metals (for example, cadmium) are more bioavailable at lower salinities. Therefore, if a sediment sample from a low salinity location is tested with overlying water. Because waters of high salinity, there is the potential that metal toxicity may be reduced. The suite of species provided in this standard allow these tests to be conducted over the range of pore-water salinities routinely encountered in field-collected sediments from North American estuarine or marine environments (USEPA 1994a (1)). In addition, artificial sea salts may contain chelating agents (EDTA) that can potentially influence the bioavailability of metals. Certain brands of artificial salts are available from manufacturers without the procedures addition of sodium thiosulfate that can also influence the toxicity of contaminants.

6.1.12 Most assessments of contaminated sediment rely on acute-lethality testing methods (for example, <10 d; (USEPA-USACE 1977, 1991, 1998, (114, 115, 116)). Acute-lethality tests are useful in identifying “hot spots” of sediment contamination, but may not be applied sensitive enough to evaluate moderately contaminated areas. Sediment quality assessments using sublethal responses of benthic organisms such as effects on growth and reproduction have a high oxygen demand. Materials dissolved in interstitial waters might be removed from solution in substantial quantities by adsorption been used to sediment particles successfully evaluate moderately contaminated areas (Dillon et al. 1994, Kemble et al. 1994, Ingersoll and Brunson 1998, (117, 82, 81), Annex A2). Insufficient information is available to determine if the long-term test chamber during conducted with *Leptocheirus plumulosus* (Annex A2) is more sensitive than 10-d toxicity tests conducted with this or other species.

6.1.13 Despite the test. The dynamics interferences previously listed, existing sediment testing methods that include measurement of contaminant partitioning between solid sublethal endpoints may be used to provide a rapid and dissolved phases at the initiation direct measure of the test should therefore effects of contaminants on benthic communities (for example, Canfield et al. (118)). Laboratory tests with field-collected sediment can also be considered, especially used to determine temporal, horizontal, or vertical distribution of contaminants in relation sediment. Most tests can be completed within two to assumptions four weeks. Legal and scientific precedence exist for use of sediment tests in regulatory decision making (for example, USEPA 1986a, Swartz 1989, (119, 120)). Furthermore, sediment tests with complex contaminant mixtures are important tools for making decisions about the extent of remedial action for contaminated aquatic sites and for evaluating the success of remediation activities.

6.2 Species-specific Interferences— Interferences of tests for each species are described in Annex A1 and Annex A2.

7. Apparatus Reagents and Materials

7.1 Facilities—Flow-through troughs or aquaria containing either clean (uncontaminated) natural sea water:

7.1.1 Requirements—Sea water or reconstituted used to test and culture organisms should be uniform in quality. Acceptable sea water should allow satisfactory survival, growth, or reproduction of the test organisms. Test organisms should not show signs of disease or apparent stress (for example, discoloration, unusual behavior). If problems are observed in the culturing or testing of organisms, it is desirable to evaluate the characteristics of the water. See USEPA (1993 (121)) and Guide E 729 for holding amphipods after field collection a recommended list of chemical analyses of the water supply.

7.1.2 Source:

7.1.2.1 Culture and p testing water can be natural or synthetic seawater (USEPA-USACE 2001 (2)).

7.1.2.2 The source of natural water will depend to a test. The holding tanks some extent on the objective of the test and any areas used for manipulating live amphipods the test organism that is being used. All natural waters should be located in a room or space separate obtained from an uncontaminated surface-water source beyond the influence of known discharges. It may be desirable to collect water at slack high tide, or within one h after high tide. Suitable surface water sources should have intakes that in which toxicity tests are positioned to: (1) minimize fluctuations in quality and contamination, (2) maximize the concentration of dissolved oxygen (DO), and (3) ensure low concentrations of sulfide and iron. For estuarine tests, water having a salinity as near as possible to the desired test salinity should be collected from an uncontaminated area.

7.1.2.3 Alternatively, it may be desirable to dilute full strength sea water with an appropriate fresh water source. Sources of fresh water (that is, 0 ‰) for dilutions include deionized water, uncontaminated well or spring water, or an uncontaminated surface-water source. Municipal-water supplies may be variable and may contain unacceptably high concentrations of materials are prepared, such as copper, lead, zinc, fluoride, chlorine, or equipment is cleaned. The chloramines. Chlorinated water supply system should not be equipped with salinity used to dilute water utilized for culturing or testing because residual chlorine and temperature control and aeration:

7.1.1 Test chambers containing sediment chlorine- produced oxidants are toxic to many aquatic organisms. Dechlorinated water should only be held in used as a well-lighted (at least 100 lx at last resort for diluting sea water to the test sediment surface); constant temperature room, incubator, desired salinity since dechlorination is often incomplete (Guide E 729; USEPA, 1993 (121)). It might be desirable or recirculating water bath necessary to ma dilute full strength seawater with an appropriate freshwater source to achieve 5 % or 20 % (or the experimental temperature. Air selected salinity; section 1.4) used for aeration should in culturing or testing of *L. plumulosus* (USEPA-USACE 2001 (2), Section 12.

7.1.2.4 For site-specific investigations, it may be desirable to have the water-quality characteristics of fumes, oil, and water; filters the overlying water (that is, salinity) as similar as possible to remove oil and the site water are desirable. The area containing (section 1.4). For certain applications the experimental design might require use of water from the site where sediment is

collected. In estuarine systems, however, the pore-water salinity of sediments may not be the same as the overlying water at the time of collection (Sanders et al., 1965 (122)).

7.1.2.5 Water that might be contaminated with facultative pathogens may be passed through a properly maintained ultraviolet sterilizer equipped with an intensity meter and flow controls or passed through a filter with a pore size of fumes, both to prevent contamination 0.45 µm or less.

7.1.2.6 Natural sea water might need aeration using air stones, surface aerators, or column aerators. Adequate aeration will stabilize pH, bring concentrations of test materials DO and other gases into equilibrium with air, and minimize oxygen demand and concentrations of volatiles. The concentration of DO in source water should be between 90 to protect researchers from exposure 100 % saturation to toxic volatile materials help ensure that might be released from the DO concentrations are acceptable in test sediments. Enclosures may chambers. Natural sea water used for holding or acclimating, culturing, and testing amphipods should be needed filtered (<5 µm) shortly before use to ventilate the area surrounding test chambers.

7.1.2 The exposure room remove suspended particles and organisms.

7.1.2.7 Water that is prepared from natural sea water should be equipped with a timing device stored in clean, covered containers at 4°C. USEPA-USACE (2001(2)) states that natural sea water should be used within 2 d for photoperiod control. If a photoperiod other than continuous light larval toxicity tests (Woelke, 1968 (123), 1972 (124) ; Cardwell et al., 1977 (125), 1979 (126)). However, investigators have found that when sea water is continuously aerated, it might can be desirable held for up to incorporate a 15 month before use with certain species (David Moore, MEC Analytical, Carlsbad, CA; personal communication).

7.1.3 Reconstituted/Synthetic Seawater :

7.1.3.1 Although reconstituted water is acceptable, natural seawater is preferable, especially for tests involving chemicals whose bioavailability is affected by seawater chemistry. Reconstituted water can be prepared by adding specified amounts of reagent-grade chemicals to 30-min transition period when lights go on high-purity deionized water (Guide E 729; USEPA, 1993 (121)). Acceptable high-purity water can be prepared using deionization or reverse-osmosis units (section 7.1; USEPA, 1993 (121)). Test water can also be prepared by diluting natural water with deionized water (Kemble et al., 1994 (51)).

7.1.3.2 Deionized water should be obtained from a system capable of producing at least 1 MΩ (mega-ohms) water. If large quantities of high quality deionized water are needed, it may be advisable to reduce stress to supply the organisms laboratory grade water deionizer with preconditioned water from sudden large changes in light intensity (10). If a mixed-bed water treatment system.

7.1.3.3 Reconstituted sea water is prepared by adding specified amounts of a suitable salt reagent to high-purity deionized water (Guide E 729, USEPA, 1991(127)). Suitable salt reagents can be reagent grade chemicals, or commercial sea salts. Pre-formulated brine (for example, 60 to 90 %), prepared with dry ocean salts or heat-concentrated natural sea water, can also be used. (USEPA, 1994 (1) USEPA -USACE 2001, (2))

7.1.3.4 A synthetic sea formulation called GP2 is prepared with reagent grade chemicals that can be diluted with a suitable high-quality water to have the room temperature desired salinity (USEPA, 1994b (128)).

7.1.3.5 The suitability and light controls consistency of a particular salt formulation for use in holding and testing should be verified by laboratory tests because some formulations can produce unwanted toxic effects or sequester contaminants (Environment Canada, 1992 (5) ; USEPA-USACE 2001(2)). In controlled tests with the saelt formulations mentioned above, Emery et al. (1997 (8)) found differency pes in survival, growth, and reproduction, and that laboratories can have acceptable performance (that is, survival) with any of the experiment salts evaluated. Because of higher growth rates observed in-e the Crystal Sea Marinemix® seasalt, they recommended its use for culturing and testing of *L. plumulosus* (Emery et al., 1997 (8); Annex A2).

7.1.3.6 To obtain the desired holding or acclimation salinity, sea salts or a hypersaline solution (USEPA, 1993 (121)) brine can be added to a suitable freshwater, deionized water, estuarine water, or the laboratory's sea water supply may be diluted with a suitable freshwater or deionized water.

7.1.3.7 Salinity, pH, and DO should be measured on each batch of reconstituted water. The reconstituted water should be aerated before use to adjust pH and DO to the acceptable ranges (for example, section 7.1). The artificial sea salts should be held for at least two week before use to allow pH to become more stable and reduce the activity of chelating agents (Environment Canada 1992 (5)).

7.2 Construction Materials—Equipment and facilities that contact stock solutions, test solutions, or any water or sediment into Formulated Sediment—Formulated sediments are mixtures of materials which test organisms will be placed should mimic the physical components of natural sediments. Formulated sediments have not contain substances that can been routinely applied to evaluate sediment contamination. A primary use of formulated sediment could be leached as a control sediment. Formulated sediments allow for standardization of sediment testing or provide a basis for conducting sediment research. Formulated sediment provides a basis by aqueous solutions in amounts that adversely affect test organisms, which any testing program can assess the acceptability of their procedures and facilities. In addition, equipment and facilities that contact stock or formulated sediment provides a consistent measure evaluating performance-based criteria necessary for test solutions or acceptability. The use of formulated sediment should be chosen to minimize sorption eliminates interferences caused by the presence of test materials from water. Glass, Type 316 stainless steel, nylon, high-density polyethylene, polycarbonate and fluorocarbon plastics should be used whenever possible to minimize dissolution, leaching, and sorption, except that stainless steel should not be used indigenous organisms. Spiking formulated sediments with specific chemicals would reduce variation in tests on metals in salt water. Concrete sediment physico-chemical characteristics and rigid plastics may be used would provide a consistent method for holding tanks and

in evaluating the water supply system, but they should be soaked, preferably fate of chemicals in flowing sea water, for a week or more before use sediment. See USEPA 2000, (11). Brass, copper, lead, cast iron pipe, galvanized metal, (64), Test Method E 1706 and natural rubber should not contact test sea water, stock solutions, or test sediment before or during the test. Tubing used in making up test sea water Guide E 1391 for additional detail regarding preparation and in aerating the test chambers should be nontoxic vinyl.⁶ New tubing should be aged at least one week prior to use. Separate sieves, dishes, containers, and other equipment should be used to handle test sediment or other toxic materials and these should be kept and stored separately from those used to handle live animals prior to testing. use of formulated sediment.

7.3 Test Chambers—Species specific information on test chambers is given in Annex A2-Annex A5. Reagents—Data sheets should be followed for reagents and other chemicals purchased from supply houses. The test chambers material(s) should be placed in water bath to minimize temperature fluctuations, at least reagent grade, unless a test using a formulated commercial product, technical-grade, or use-grade material is specifically needed. Reagent containers should be dated when received from the supplier, and the shelf life of the reagent should not be aerated. Aeration can exceed. Working solutions should be provided as in 13.1. dated when prepared and the recommended shelf life should not be exceeded.

7.4 Cleaning—Test chambers and other glassware, and equipment used to store and prepare test sea water, stock solutions, and test sediment should be cleaned before use. All glassware should be cleaned before each use by washing with laboratory detergent, followed by three distilled water rinses, 10 % nitric (HNO₃) Standards—Appropriate USEPA, APHA, or hydrochloric (HCl) acid rinse, and at least two distilled water rinses. Metals, sulfides, and carbonate deposits are removed by the acid rinse. Organic chemicals should be removed by a water-miscible organic solvent rinse followed by a distilled water rinse, or by baking ASTM standards for 8 h at 300 to 400°C. The use of hypochlorite solution is not recommended, because it is highly toxic to the test organisms (12) and difficult to remove from some materials. At the end of each test, all items that are to be used again should be immediately (a) emptied, (b) rinsed with water, (c) cleaned by a procedure appropriate for removing the test material, chemical and (d) rinsed at least twice with deionized, distilled, or clean sea water. Large plastic containers used only for non-toxic sediments and water may be rinsed after use with clean sea water. They physical analyses should be used only when possible. For those measurements for toxicity tests and stored in a room that is free from toxic fumes. Glassware used only for live animals, which standards do not exposed to toxicants, may be cleaned using only clean distilled exist or sea water, since the use of detergents is sometimes detrimental to live organisms.

7.5 Acceptability—The acceptability of new holding or testing facilities are not sensitive enough, methods should be demonstrated by conducting a “nontoxicant” test in which all test chambers contain control sediment and clean sea water. Survival of the test species will demonstrate whether facilities, water, control sediment, and handling techniques are adequate to result in acceptable (≥90 %) control survival in the absence of toxicants. obtained from other reliable sources.

8. Hazards

8.1 Many materials can affect humans adversely if precautions are inadequate. Therefore, skin contact

8.1 General Precautions:

8.1.1 Development and maintenance of an effective health and safety program in the laboratory requires an ongoing commitment by laboratory management and includes: (1) the appointment of a laboratory health and safety officer with all toxicants, overlying water, the responsibility and authority to develop and maintain a safety program, (2) the preparation of a formal, written health and safety plan, which is provided to each laboratory staff member, (3) an ongoing training program on laboratory safety, and (4) regular safety inspections.

8.1.2 Collection and use of sediments may involve substantial risks to personal safety and health. Chemicals in field-collected sediment may include carcinogens, mutagens, and other potentially toxic compounds. Inasmuch as sediment testing is often started before chemical analyses can be completed, worker contact with sediment needs to be minimized by: (1) using gloves, laboratory coats, safety glasses, face shields, and respirators as wearing appropriate protective gloves (especially when washing equipment or putting hands into test appropriate, (2) manipulating sediments under a ventilated hood or in an enclosed glove box, and (3) enclosing and ventilating the exposure system. Personnel collecting sediment samples and conducting tests should take all safety precautions necessary for the prevention of bodily injury and illness which might result from ingestion or invasion of infectious agents, inhalation or absorption of corrosive or toxic substances through skin contact, and asphyxiation because of lack of oxygen or presence of noxious gases.

8.1.3 Before beginning sample collection and laboratory work, personnel should determine that all required safety equipment and materials have been obtained and are in good condition.

8.2 Safety Equipment:

8.2.1 Personal Safety Gear—Personnel should use safety equipment, such as rubber aprons, laboratory coats, respirators, gloves, safety glasses, face shields, hard hats, and glasses. Special precautions, safety shoes.

8.2.2 Laboratory Safety Equipment—Each laboratory should be provided with safety equipment such as covering test chambers first-aid kits, fire extinguishers, fire blankets, emergency showers, and ventilating the area surrounding the chambers, eye wash stations. Mobile laboratories should be taken when conducting tests on volatile materials. Information on toxicity equipped with a telephone to humans (6), recommended enable personnel to summon help in case of emergency.

8.3 General Laboratory and Field Operations:

8.3.1 Special handling procedures (7); and chemical precautionary guidance in Material Safety Data Sheets (MSDS) should be

followed for reagents and other chemicals purchased from supply houses.

8.3.2 Work with some sediments may require compliance with rules pertaining to the handling of hazardous materials. Personnel collecting samples and performing tests should not work alone.

8.3.3 It is advisable to wash exposed parts of the body with bactericidal soap and water immediately after collecting or manipulating sediment samples.

8.3.4 Strong acids and volatile organic solvents should be studied before used in a test is begun. Special precautions might fume hood or under an exhaust canopy over the work area.

8.3.5 An acidic solution should not be necessary mixed with a hypochlorite solution because hazardous fumes might be produced.

8.3.6 To prepare dilute acid solutions, concentrated acid should be added to water, not vice versa. Opening a bottle of concentrated acid and adding concentrated acid to water should be performed only under a fume hood.

8.3.7 Use of ground-fault systems and leak detectors is strongly recommended to help prevent electrical shocks. Electrical equipment or extension cords not bearing the approval of Underwriter Laboratories should not be used. Ground-fault interrupters should be installed in all "wet" laboratories where electrical equipment is used.

8.3.8 All containers should be adequately labeled to indicate their contents.

8.3.9 A clean and well-organized work place contributes to safety and reliable results.

8.4 *Disease Prevention*—Personnel handling samples which are known or suspected to contain human wastes should be immunized against hepatitis B, tetanus, typhoid fever, and polio. Thorough washing of exposed skin with bactericidal soap should follow handling of samples collected from the field.

8.5 *Safety Manuals*—For further guidance on safe practices when handling sediment samples and conducting toxicity tests, check with the permittee and consult general industrial safety manuals including USEPA 1986b, Walters and Jameson 1984, (129, 130).

8.2 Field sediments to be tested, especially those from effluent areas, might contain organisms that can be pathogenic to humans. Special precautions when dealing with these sediments might include immunization prior to sampling

8.6 *Pollution Prevention, Waste Management, and use of bactericidal soaps after working with Sample Disposal*—Guidelines for the sediments:

8.3 Sediments collected from the field might be contaminated with unknown concentrations of many potentially toxic materials, handling and laboratory prepared sediments might be spiked with high concentrations disposal of toxicants. Any potentially contaminated sediments hazardous materials should be handled in a manner to minimize exposure strictly followed (Guide D 4447). The Federal Government has published regulations for the management of researchers to toxic compounds. Mixing of toxic sediments in open containers, spiking of laboratory prepared sediments, hazardous waste and loading has given the States the option of toxic sediments into test chambers should be done in a well-ventilated area, preferably a chemical fume hood. Face shields either adopting those regulations or protective goggles should developing their own. If States develop their own regulations, they are required to be worn during any operations that might involve accidental splashing of sediments, such at least as sieving, mixing and loading into test chambers:

8.4 Health and safety precautions and applicable regulations for disposal of stock solutions, overlying water from test chambers, test organisms, and sediments should be considered before beginning stringent as the Federal regulations. As a test (see Guide D 4447). Consideration handler of cost as well as detailed regulatory requirements might be necessary. For tests involving spiked sediments with known toxicants, removal or degradation of toxicants before disposal of stock solutions, test sediments, and water hazardous materials, it is sometimes desirable:

8.5 Cleaning of equipment your responsibility to know and comply with a volatile solvent such as acetone, should be performed only the pertinent regulations applicable in a well-ventilated area the State in which no smoking is allowed and no open flame, such as a pilot light, is present. Cleaning equipment with acids should be done only in a well-ventilated area, and protective gloves and safety goggles should be worn:

8.6 To prepare dilute acid solutions, concentrated acid should be added you are operating. Refer to water, not vice versa. Opening a bottle the Bureau of concentrated acid and adding concentrated acid to water should be performed only in a well-ventilated area or a chemical fume hood:

8.7 Use National Affairs Inc., 1986 (131)) for the citations of ground fault systems and leak detectors is strongly recommended to help prevent electrical shocks because salt water is a good conductor of electricity. the Federal requirements.

9. Toxicity Test Water Facilities, Equipment, and Supplies

9.1 *General Requirements*—Besides being available in adequate supply, water used in toxicity tests should be acceptable to Before a sediment test organisms and the purpose of the test. The minimum requirement for acceptable water for use in acute toxicity tests is that healthy conducted in any new test organisms survive facility, it is desirable to conduct a "non-toxicant" test, in which all test chambers contain a control sediment, and in the overlying water with sediment for the duration of holding and testing without showing signs of disease no added test material (section 11.14). Survival, growth, or apparent stress such as unusual behavior, changes in appearance, or death. The water in which reproduction of the test organisms will demonstrate whether facilities, water, control sediment, and handling techniques are held prior adequate to the test should be uniform result in quality acceptable species-specific control numbers (for example, see Table A1.3 in that the concentration of contaminants Annex A1 and

Table A2.3 in Annex A2). Evaluations may also be made on the range magnitude of temperature and salinity encountered during the holding period do not adversely affect the survival of the test organisms within-chamber and between-chamber variance in the holding tanks or in the control treatments during the test. a test (section 11.14).

9.2 Source Facilities:

9.2.1 Natural Salt Water—If natural salt water is used, it should be obtained from an uncontaminated area known

9.2.1 The facility must include separate areas for culturing and testing to support a healthy, reproducing population of reduce the possibility of contamination by test materials and other substances, especially volatile compounds. Holding, acclimation, and culture chambers should not be in a room where sediment tests are conducted, where stock solutions or sediments are prepared, or where equipment is cleaned. Test chambers may be placed in a comparable sensitive species. The temperature-controlled recirculating water intake should be positioned bath or a constant-temperature area. An enclosed test system is desirable to minimize fluctuations in provide ventilation during tests to limit exposure of laboratory personnel to volatile substances.

9.2.2 Light of the quality and illuminance normally obtained in the possibility of contamination, and laboratory is adequate (about 100 to maximize 1000 lux using wide-spectrum fluorescent lights: for example, cool-white or day-light has been used successfully to culture and test organisms). Lux is the unit selected for reporting luminance in this standard. Multiply units of dissolved oxygen lux by 0.093 to help ensure low concentrations convert to units of sulfide and iron. A specially designed system might be necessary footcandles. Multiply units of lux by 6.91×10^{-3} to obtain salt water from a natural water source. To ensure uniform quality, water convert to units of $\mu\text{mol}^{-2} \text{s}^{-1}$ (assuming an average wavelength of 550 nm ($\mu\text{mol}^{-2} \text{s}^{-1} = W \text{ m} \times \lambda(\text{nm}) \times 8.36 \times 10^{-3}$)). Illuminance should be monitored as in 9.4. These precautions are intended to ensure that test organisms are not apparently stressed by water quality during holding, acclimation, and testing and that water quality does not unnecessarily affect test results:

9.2.2 Reconstituted Salt Water—Reconstituted salt water measured at the surface of the water. A uniform photoperiod of 16L:8D can be prepared by adding a commercially available sea salt achieved in the laboratory or specified amounts (see Guide E 729 in an environmental chamber using automatic timers. A 16:8 light:dark photoperiod should be used for culturing *L. plumulosus* (section 12.6) and Table 1) of reagent-grade chemicals for holding and acclimating **(13)A. abdita in the laboratory before testing (section 12.4; USEPA 1994a to high-quality water with (a) conductivity less than 1 $\mu\text{S}/\text{cm}(1)$.**

9.2.3 During rearing, holding, and (b) either total organic carbon (TOC) less than 2 mg/L or chemical oxygen demand (COD) less than 5 mg/L. Acceptable water can usually testing, test organisms should be prepared using properly operated deionization shielded from external disturbances such as rapidly changing light or distillation units. Reconstituted salt water pedestrian traffic.

9.2.4 Air used for aeration should be intensively aerated before use, free of oil and aging for one fumes. Filters to two weeks might be remove oil, water, and bacteria are desirable. If a residue or precipitate is present, the solution The test facility should be filtered before use. The water well ventilated and free of fumes. Oil-free air pumps should meet the criteria given in 9.1.

9.2.3 Chlorinated water must never be used where possible. Particulates can be removed from the preparation of water for toxicity tests, because residual chlorine air using filters, and chlorine produced oxidants are highly toxic to many aquatic animals oil and other organic vapors can be removed using activated carbon filters (USEPA 2000, **(12)**. Dechlorinated water**(73)**). Laboratory ventilation systems should be used only as a last resort because dechlorination checked to ensure that return air from chemistry laboratories or sample handling areas is often incomplete. Municipal drinking water is not recommended for use because in addition circulated to residual chlorine, it often contains unacceptably high concentrations culture or testing areas, or that air from testing areas does not contaminate culture areas. Air pressure differentials between areas should not result in a net flow of metals, and quality is often highly variable (see Guide E 729). potentially contaminated air to sensitive areas through open or loosely fitting doors.

9.3 Preparation Equipment and Supplies:

9.3.1 Equipment and supplies that contact stock solutions, sediments, or overlying water should not contain substances that can be leached or dissolved in amounts that adversely affect the sediment toxicity test organisms. In addition, equipment and supplies that contact sediment or water should be passed through a filter effective chosen to 5 μm or less to remove suspended particles and organisms minimize sorption of test materials from the water. Water that might be contaminated with facultative pathogens Glass, Type 316 stainless steel, nylon, high-density polyethylene, polypropylene, polycarbonate, and fluorocarbon plastics should be passed through a properly maintained ultraviolet sterilizer **(16)** or a filter effective used whenever possible to 0.45 μm or less:

9.3.1.1 If necessary, minimize leaching, dissolution, and sorption. Concrete and high-density plastic containers may be used for holding, acclimation, and culture chambers, and in the salinity water-supply system. These materials should be reduced by diluting the sea washed in detergent, acid-rinsed, and soaked in flowing water with high-quality deionized for a week or distilled water (see 9.2.2). Salinity can more before use. Cast-iron pipe should not be raised by addition of clean filtered oceanic used in water-supply systems because colloidal iron will be added to the overlying water or prepared brine. Common practice is and strainers will be needed to use a 60 to 90-g/kg saltwater brine. Such brines have been successfully prepared using slow, heat-concentration of remove rust particles. Copper, brass, lead, galvanized metal, and natural rubber must not contact overlying water; or by the addition of artificial sea salts stock solutions before or reagent grade **(13)** salts to during a natural salt water (see 9.2.2).

9.3.2 Fresh sea water test. Items made of neoprene rubber and other materials not mentioned above should not be used unless it has been shown that their use will not adversely affect survival, growth, or reproduction of the test organisms.

9.3.2 New lots of plastic products should be prepared within two days of the tested for toxicity before general use by exposing organisms to them under ordinary test conditions.

9.3.3 General Equipment:

9.3.3.1 Environmental Chamber or Equivalent Facility, with photoperiod and stored in clean, covered containers temperature control (15 to 25°C).

9.3.3.2 Water Purification System, capable of producing at $4 \pm 3^\circ\text{C}$ until sediment and least 1 M Ω (mega-ohms) of water are added (USEPA, 1993a (132)).

9.3.3.3 Analytical Balance, capable of accurately weighing to 0.01 mg (for the test chambers. It might be necessary to age reconstituted sea water *L. plumulosus* test).

9.3.3.4 Reference Weights, Class S, for one to two weeks before use. Sufficient water documenting the performance of the analytical balance(s). The balance(s) should be prepared checked with reference weights that are at one time for all test chambers. Additional water might be required for sieving control sediment to adjust salinity or for holding the test amphipods prior to the test.

9.3.3 For certain applications the experimental design might require use upper and lower ends of sea water from the test sediment collection site. In other instances, experimental treatments might involve manipulation range of the test sea water conditions:

9.4 Characterization—The following items weighings made when the balance is used. A balance should be measured checked at least twice the beginning of each year and more often if such measurements have not been made semiannually for at least two years:

9.4.1 Salinity or chlorinity, pH, particulate matter, total organic carbon (TOC), organophosphorus pesticides, organic chlorine (or organic chlorine pesticides plus polychlorinated biphenyl (PCBs)), chlorinated phenoxy herbicides, ammonia, cyanide, sulfide, bromide, fluoride, iodide, nitrate, phosphate, sulfate, calcium, magnesium, potassium, aluminum, arsenic, beryllium, boron, cadmium, chromium, cobalt, copper, iron, lead, manganese, mercury, molybdenum, nickel, selenium, silver, and zinc.

9.4.2 In estuarine areas, where large diurnal, tidal, and seasonal variations in concentrations series of organics, heavy metals and water quality might occur, more frequent monitoring might be necessary. In particular, daily measurements weighings, periodically (such as every tenth weight) during a long series of salinity, temperature, weighings, and pH, and quarterly monitoring after taking the last weight of other parameters over a tidal cycle might be desirable.

9.4.3 The methods used (see 14.2) should either (a) be accurate and precise enough to adequately characterize series (for the toxicity test water *L. plumulosus* test).

9.3.3.5 Volumetric Flasks and Graduated Cylinders, Class A, borosilicate glass or (b) have detection limits below concentrations that have been shown nontoxic plastic laboratory ware, 10 to adversely affect the 1000 mL for making test solutions.

9.3.3.6 Volumetric Pipettes, Class A, 1 to 100 mL.

9.3.3.7 Serological Pipettes, 1 to 10 mL, graduated.

9.3.3.8 Pipette Bulbs and Fillers (17).

9.3.3.9 Droppers, and Glass Tubing with Fire-Polished Edges, 4 to 6-mm inside diameter, for transferring test organisms.

9.3.3.10 Wash Bottles, for rinsing small glassware, instrument electrodes and probes.

9.3.3.11 Glass or Electronic Thermometers , for measuring water temperature.

9.3.3.12 National Bureau of Standards (NBS) Certified Thermometer (see USEPA Method 170.1, 1997b, (133)).

9.3.3.13 Dissolved Oxygen (DO), pH/Selective Ion, and Specific Conductivity Meters and Probes and temperature-compensated refractometer, for routine physical and chemical measurements are needed. Unless a test is being conducted to specifically measure the effect of DO or conductivity, a portable field-grade instrument is acceptable.

9.3.3.14 Equipment for measuring ammonia (that is, an ammonia-specific probe or an ammonia test kit) is also necessary.

9.3.3.15 See USEPA (1994a (1)) and USEPA-USACE (2001 (2)) for a list of additional equipment and supplies.

9.3.4 Test Chambers—Test chambers to be used in sediment toxicity tests are 1-L glass containers (beakers or wide-mouth jars) with an internal diameter of about 10 cm. Each test chamber should have a cover. Acceptable covers include watch glasses, plastic lids, glass culture dishes, or parafilm. It may be necessary to drill a hole in the glass cover to allow the insertion of a pipette for aeration (USEPA 1994a (1)).

9.3.5 Cleaning:

9.3.5.1 All non-disposable sample containers, test chambers, and other equipment that have come in contact with sediment should be washed after use in the manner described as follows to remove surface contaminants.

9.3.5.2 Soak 15 min in tap water, and scrub with detergent, or clean in an automatic dishwasher.

9.3.5.3 Rinse twice with tap water.

9.3.5.4 Carefully rinse once with fresh, dilute (10 %, V:V) hydrochloric or nitric acid to remove scale, metals, and bases. To prepare a 10 % solution of acid, add 10 mL of concentrated acid to 90 mL of deionized water.

9.3.5.5 Rinse twice with deionized water.

9.3.5.6 Rinse once with full-strength, pesticide-grade acetone to remove organic compounds (use a fume hood or canopy). Hexane might also be used as a solvent for removing non-ionic organic compounds. However, acetone is preferable if only one organic solvent is used to clean equipment.

9.3.5.7 Rinse three times with deionized water.

9.3.5.8 All test chambers and equipment should be thoroughly rinsed with the dilution water immediately before use in a test.

9.3.5.9 Many organic solvents leave a film that is insoluble in water. A dichromate-sulfuric acid cleaning solution can be used in place of both the organic solvent and the acid (Guide E 729), but the solution might attack silicone adhesive and leave chromium residues on glass. An alternative to use of dichromate-sulfuric acid could be to heat glassware for 8 h at 450°C.

10. Test Sample Collection, Storage, Manipulation, and Control Sediments Characterization

10.1 General—Before Collection:

10.1.1 Before the preparation or collection of test sediment, an approved written procedure should be prepared established for the handling of sediment—~~ts~~ whatic might contain unknown quantities of ~~many~~ potentially toxic contaminants (see Section chemicals (Section 8)).

10.1.2 Sediments are spatially and temporally variable (Stemmer et al. 1990a (66)). Replicate samples should be collected to determine variance in sediment characteristics. Sediment should be collected with as little disruption as possible; however, subsampling, compositing, or homogenization of sediment samples may be necessary for some experimental designs. Sampling may cause loss of sediment integrity, change in chemical speciation, or disruption of chemical equilibrium (Guide E 1391). A benthic grab or core should be used rather than a dredge to minimize disruption of the sediment sample. Sediment should be collected from a depth that will represent expected exposure.

10.1.3 Exposure to direct sunlight during collection should be minimized, especially if the sediment contains photolytic compounds (Davenport and Spacie 1991, Oris and Giesy 1985, (106, 107)). Sediment samples should be cooled to 4°C in the field before shipment (Guide E 1391). Dry ice can be used to cool samples in the field; however, sediments should never be frozen. Monitors can be used to measure temperature during shipping (USEPA 2000, (73)).

10.1.4 For additional information on sediment collection and shipment see Test Method E 1706, Guide E 1391, USEPA, 2000 (64), and USEPA, 2000 (134) for additional guidance.

10.2 Characterization—Sediments chosen for use should be characterized Storage:

10.2.1 Since the chemicals of concern and influencing sediment characteristics are not always known, it is desirable to hold the sediments after collection ~~own~~ ing the dark at 4°C. Traditional convention has held that toxicity tests should be started as soon as possible following collection from the field, although actual recommended storage times range from two weeks (Guid: ~~pHe~~ E 1391) to less than eight weeks (USEPA-USACE, 1998) (135). Discrepancies in recommended storage times reflected a lack of data concerning the effects of long-term storage on the physical, chemical, and toxicological characteristics of the sediment. However, numerous studies have recently been conducted to address issues related to sediment storage (Dillon et al. 1994, (136); Becker et al. 1995, (137), Carr and Chapman 1995, (138), Moore et al. 1996, (139), Sargda and Burton 1995, (140), Sijm et al. 1997, (141), DeFoe and Ankley 1998, (142)). The conclusions and recommendations offered by these studies vary substantially and appear to depend primarily upon the type or class of chemical(s) present. Considered collectively, these studies suggest that the recommended guidance that sediments be tested sometime between the time of collection and 8 weeks storage is appropriate. Additional guidance is provided below.

10.2.2 Extended storage of sediments that contain high concentrations of labile chemicals (for example, ammonia, volatile solids), ~~partice size distribution (percent sand, silt, organics)~~ may lead to a loss of these chemicals ~~and clay~~; a corresponding reduction in toxicity. Under these circumstances, the sediment should be tested as soon as possible after collection, but not later than within two weeks (Sarda and ~~p~~ Burton 1995 (140)). Sediments that exhibit low-level to moderate toxicity can exhibit considerable ~~t~~ temporal variability in toxicity, although the direction of change is often unpredictable (Carr and Chapman 1995 (138); Moore et al. ~~Θ~~ 1996 (139); DeFoe and Ankley 1998 (142)). For these types of sediments, the ~~r~~ recommended storage time of <8 weeks may be most appropriate. In some situations, a minimum storage period for low-to-moderately contaminated sediments might include biological oxygen demand, chemical oxygen demand, Eh or pE, total inorganic carbon, metals, synthetic organic compounds, oil may help reduce variability. For example, DeFoe and ~~grease, organosilicones~~, Ankley 1998 (142) observed high variability in survival during early testing periods (for example, <2 weeks) in sediments with low toxicity. De Foe and petroleum hydrocarbons. Interstitial water might also Ankley 1998, (142) hypothesized that this variability partially reflected the presence of indigenous predators that remained alive during this relatively short storage period. Thus, if predatory species are known to exist, and the sediment does not contain labile contaminants, it may be desirable to store the sediment for a short period before testing (for example, 2 weeks) to reduce potential for interferences from indigenous organisms. Sediments that contain comparatively ~~z~~ stable compounds (for example, high molecular weight compounds such as PCBs) or which exhibit a moderate-to-high level of toxicity, typically do not vary appreciably in ~~4.4. Toxicological results~~ toxicity in relation to storage duration (Moore et al. 1996 (139), DeFoe and Ankley 1998, (142)). For these sediments, long-term storage (for example, >8 weeks) can identify samples that should be ~~subjected~~ undertaken.

10.2.3 Researchers may wish to ~~m~~ corenduct additional characterizations of sediment to evaluate possible effects of storage. Concentrations of ~~chyemicals~~ chemicals of concern could be measured periodically in pore water during the storage period and at the start of the sediment test (Kemble et al. 1994, (82)). Ingersoll et al. 1993, (143) recommend conducting a toxicity test with pore water within two weeks from ~~sediment caol~~ collection and at the start of the sediment test. Freezing might further change sediment properties such as grain size or chemical partitioning and should be avoided (Guide E 1391; Schuytema et al. 1989, (144)). Sediment should be ~~stogred~~ stored with no air over the sealed samples (no head space) at 4°C before the start of a test (Shuba et al.

1978, (145)). Sediment may be stored in containers constructed of suitable materials as outlined in Section 9.

10.3 *Control Sediment Manipulation:*

10.3.1 *Collection—Control Homogenization:*

10.3.1.1 Samples tend to settle during shipment. As a result, water above the sediment should not be collected from discarded, but should be mixed back into the amphipod collection site or from another area that sediment during homogenization. Sediment samples should not be sieved to remove indigenous organisms unless there is a good reason to believe they will influence the geochemical requirements response of the test-species organisms. Large indigenous organisms and that large debris can provide a nontoxic reference sediment for evaluation be removed using forceps. Reynoldson et al., 1994 (146), observed reduced growth of the condition of the test population subject to laboratory procedures; amphipods, midges, and for statistical comparison mayflies in sediments with test sediment. Control sediment should be brought to the elevated numbers of oligochaetes and recommended sieving area in a clean collecting basin. Any water overlying the sediment or used sediments suspected to rinse the sediment into the collecting basin should have high numbers of indigenous oligochaetes. If sediments must be saved so that fine particles contained in the water can sieved, it may be recombined into the sediment. Any sediment that shows evidence of contamination desirable to analyze samples before and after sieving (for example, oil sheen) should be discarded. As pore-water metals, dissolved organic carbon (DOC), acid volatile sulfide (AVS), total organic carbon (TOC)) to document the influence of sieving on sediment chemistry.

10.3.1.2 If sediment is collected, bottom temperature and salinity and collected from multiple field samples, the sediment temperature should can be recorded, pooled and mixed using stirring or a composite sediment sample from all shovelful, dredge hauls, rolling mill, feed mixer, or grabs should be collected for analysis other suitable apparatus (Guide E 1391). Homogenization of water content, particle size distribution, and organic content.

10.3.1.1 At least annually, control sediment should can be empirically characterized as in 10.2, accomplished using a hand-held drill outfitted with a stainless steel auger (diameter 7.6 cm, overall length 38 cm, auger bit length 25.4 cm (Kemble et al. 1994, (82)).

10.3.2 *Sieving—A separate clean container should Sediment Spiking:*

10.3.2.1 Test sediment can be set up to sieve and contain prepared by manipulating the properties of a control sediment. Control Mixing time (Stemmer et al. 1990a, (66)) and aging (Landrum 1989, Word et al. 1987, Landrum and Faust 1992, (85, 86, 87)) of spiked sediment can affect bioavailability of chemicals in sediment. Many studies with spiked sediment are often started only a few days after the chemical has been added to the sediment. This short time period may not be long enough for sediments to equilibrate with the spiked chemicals (section 10.3.2.6). Consistent spiking procedures should be sieved twice: first followed in order to make interlaboratory comparisons. Limited studies have been conducted comparing appropriate methods for spiking chemicals in sediment. Additional research is needed before more definitive recommendations for spiking of sediment can be outlined in this standard. The guidance provided in the following sections has been developed from a variety of sources. Spiking procedures that have been developed using one sediment or test-species and organism may not be applicable to other macrobenthos, sediments or test organisms. See USEPA 2000, (64) and Guide E 1391 for additional detail regarding sediment-spiking techniques.

10.3.2.2 The cause of sediment toxicity and the interactive effects of chemicals can be determined by spiking a sediment with chemicals or complex waste mixtures (Lamberson and Swartz 1988, (109)). Sediments spiked with a range of concentrations can be used to generate either point estimates (for example, LC50) or a minimum concentration at which effects are observed (lowest-observable-effect concentration; LOEC). Results of tests may be reported in terms of a BSAF (Biota-sediment accumulation factor; (Ankley et al. 1992b, (147)). The influence of sediment physico-chemical characteristics on chemical toxicity can also be determined with sediment-spiking studies Swartz et al. 1994, (148).

10.3.2.3 The test material(s) should be at least reagent grade, unless a test using a formulated commercial product, technical-grade, or use-grade material is specifically needed. Before a test is started, the following should be known about the test material: (1) the identity and concentration of major ingredients and impurities, (2) water solubility in test water, (3) log Kow, BCF (from other test species), persistence, hydrolysis, and photolysis rates of the test substrate, (4) estimated toxicity to the test-salinity organism and to humans, (5) if the test concentration(s) are to be measured, the precision and bias of the analytical method at the planned concentration(s) of the test material, and (6) recommended handling and disposal procedures. Addition of test material(s) to sediment may be accomplished using various methods, such as a: (1) rolling mill, (2) feed mixer, or (3) hand mixing (Guide E 1391; USEPA (64)). Modifications of the mixing techniques might be necessary to allow time for a test material to equilibrate with the sediment. Mixing time of spiked sediment should be clean sea water prepared as limited from minutes to a few hours and temperature should be kept low to minimize potential changes in Section 9. The entire contents the physico-chemical and microbial characteristics of the collecting basin, including water sediment (Guide E 1391, USEPA 2000, (134)). Duration of contact between the chemical and suspended particles; sediment can affect partitioning and bioavailability Word et al. 1987, (86). Care should be taken to ensure that the chemical is thoroughly and evenly distributed in the sediment. Analyses of sediment subsamples is advisable to determine the degree of mixing homogeneity Ditsworth, et al. 1990 (149). Moreover, results from sediment-spiking studies should be compared with the response of test organisms to chemical concentrations in natural sediments (Lamberson and Swartz 1992 (150)).

10.3.2.4 Organic chemicals have been added: (1) directly in a dry (crystalline) form; (2) coated on the inside walls of the container (Ditsworth et al. 1990, (149)); or (3) coated onto silica sand (for example, through 5 % w/w of sediment) which is added

to the sediment (Test Method E 1706). In techniques 2 and 3, the chemical is dissolved in solvent, placed in a 0.5-mm screen glass spiking container (with or without allowing overflow from sand), then the sieving container. After solvent is slowly evaporated. The advantage of these three approaches is that no solvent is introduced to the fir sediment, only the chemical being spiked. When testing spiked sediments, procedural blanks (sediments that have been handled in the same way, including solvent addition and evaporation, but contain no added chemical) should be left undisturbed for a sufficient time tested in addition to regular negative controls.

10.3.2.5 Metals are generally added in an aqueous solution (Guide E 1391; Di Toro et al. 1990, (88)). Ammonia has also been successfully spiked using aqueous solutions (Besser et al. 1998, (151)). Inclusion of fine particles (usually at least overnight). Overlying water spiking blanks is recommended.

10.3.2.6 Sufficient time should then be decanted and allowed after spiking for the spiked chemical to equilibrate with sediment components. For organic chemicals, it is recommended that the sediment be aged at least one month before starting a test. Two months or more may be necessary for chemicals with a high log Kow (for example, through a 0.5-mm screen) into water >6; Test Method E 1706). For metals, shorter aging times (1 to 2 weeks) may be sufficient. Periodic monitoring of chemical concentrations in pore water during sediment aging is highly recommended as a salinity calculated means to bring assess the equilibration of ther spiked sediments. Monitoring of pore water during spiked sediment testing is also recommended.

10.3.2.7 Organic solvents such as triethylene glycol, methanol, ethanol, or acetone may be used, but they might affect TOC levels, introduce toxicity, alter the test level, taking into account geochemical properties of the sediment, or stimulated q undesirable growths of microorganisms (Guide E 1391). Acetone is highly volatile and might leave the system more readily than triethylene glycol, methanol, or ethanol. A surfactant should not be used in the preparation of a stock solution because it might affect the interstitial water. Again, bioavailability, form, or toxicity of the test material.

10.3.2.8 If the test contains both a negative control and a solvent control, the survival, growth, or reproduction of the organisms tested should be allowed to settle, overlying water should compared in the two controls. If a statistically significant difference is detected between the two controls, only the solvent control may be decanted, used for meeting the acceptability of the test and as the basis for calculation of results. The negative control might provide additional information on the general health of the organisms tested. If no statistically significant difference is detected, the data from both controls should be thoroughly mixed to evenly distribute fine particles used for meeting the acceptability of the test and as the basis for calculation of results (Guide E 1241). If performance in the solvent control is markedly different from that settle on in the negative control, it is possible that the data are compromised by experimental artifacts and may not accurately reflect the toxicity of the chemical in natural sediments.

10.3.3 *Storage*—The control sediment Test Concentration(s) for Laboratory-spiked Sediments:

10.3.3.1 If a test is intended to generate an LC50, a toxicant concentration series (0.5 or higher) should be stored in glass or rigid plastic containers selected that will provide partial mortality at $4 \pm 3^\circ\text{C}$ until two or more concentrations of the test chambers are prepared: chemical. The LC50 of a particular compound may vary depending on physical and chemical sediment should characteristics. It may be desirable to conduct a range-finding test in which the organisms are exposed to a control and must not be frozen three or more concentrations of the test material that differ by a factor of ten. Results from water-only tests could be used to establish concentrations to be tested in a whole-sediment test based on predicted pore-water concentrations (Di Toro et al. 1991, (70)). See Section 13 for a description of procedures to analyze data generated from these studies.

10.3.3.2 Whole-sediment chemical concentrations might be normalized to factors other than dry weight. For example, concentrations of nonpolar organic compounds might be normalized to sediment organic-carbon content and simultaneously extracted metals might be normalized to acid-volatile sulfides (DiToro 1990, 1991, (88, 70)).

10.3.3.3 In some situations it might be necessary to simply determine whether a specific concentration of test material is toxic to the test organism, or whether adverse effects occur above or below a specific concentration. When there is interest in a particular concentration, it might only be necessary to test that concentration and not to determine an LC50.

10.4 *Field-Collected Test Sediment Characterization:*

10.4.1 *Collection*—The spatial or temporal

10.4.1 All sediments should be characterized for at least: salinity, pH, and ammonia of the pore water, organic carbon content (total organic carbon, TOC), particle size distribution (percent sand, silt, clay), and percent water content (Plumb, 1981, (152)). See section 10.4.5 for a description of procedures for isolating interstitial water.

10.4.2 Other analyses on sediments might include: biological oxygen demand, chemical oxygen demand, cation exchange capacity, Eh, total inorganic carbon, total volatile solids, acid volatile sulfides, dissolved organic carbon, organic nitrogen, metals, synthetic organic compounds, oil and grease, and petroleum hydrocarbons in sediment toxicity can and interstitial water.

10.4.3 Macrobenthos may be determined evaluated by collecting potentially contaminated sediment from field sites. A benthic grab or core should be used rather than a dredge to minimize disruption of subsampling the sample. field-collected sediment. If the sediment is collected with a grab, glass cores direct comparisons are to be made, subsamples for toxicity testing should be used to collect a sample collected from the upper 2 cm, or desired layer, same sample for analysis of sediment physical and chemical characterizations. Qualitative descriptions of the test sediment. This operation is facilitated if sediment may include color, texture, and presence of macrophytes or animals. Monitoring the grab can odor of sediment samples should be avoided because of potential hazardous volatile chemicals.

10.4.4 *Analytical Methodology:*

10.4.4.1 Chemical and physical data should be obtained using appropriate standard methods whenever possible. For those measurements for which standard methods do not exist or are not sensitive enough, methods should be obtained from other reliable sources.

10.4.4.2 The precision, accuracy, and bias of each analytical method used should be determined in the top so appropriate matrix: that the undisturbed sediment surface is exposed. The sample is, sediment, water, and tissue. Reagent blanks and analytical standards should be transferred to a clean (see 7.4) glass, high-density polyethylene or fluorocarbon plastic sample container. It is desirable analyzed and recoveries should be calculated.

10.4.4.3 Concentration of spiked test material(s) in sediment, interstitial water, and overlying water should be measured as much often as possible to avoid contact practical during a test. If possible, the concentration of the sample with metals, including stainless steel, and plastics including polypropylene and low-density polyethylene as contaminant interactions might occur. The sample must be cooled test material in the field to about 4°C, overlying water, interstitial water, and sediments should be measured at 4 ± 3°C until the sediment is distributed to the start and end of a test. Measurement of test chambers. Test sediment material(s) degradation products might also be desirable.

10.4.4.4 Separate chambers should be stored in set up at the dark, no longer than two weeks before start of a test and destructively sampled during and at the initiation end of the test, and must not be frozen or allowed test to dry. Freezing and longer storage might change monitor sediment properties chemistry. Test organisms and have been shown food should be added to change the toxicity these extra chambers.

10.4.4.5 Measurement of stored sediment (18, 19). Field collected test sediments should not material(s) concentration in water can be wet sieved, but if obvious large predators or other large organisms are present, they should be removed accomplished by forceps. For some applications, it might be desirable pipeting water samples from about 1 to remove small objects by press sieving 2 cm above the sediment through a 2-mm sieve. If sediment is stored longer than two weeks, it surface in the test chamber. Overlying water samples should be retested to confirm that toxicity has not changed.

10.4.1.1 If sediment is collected contain any surface debris, any material from multiple field samples and pooled to meet technical objectives, the sediment should be thoroughly homogenized by stirring sides of the test chamber, or mixing by hand, or with any sediment.

10.4.4.6 Measurement of test material(s) concentration in sediment at the aid end of a rolling mill as in 10.8.

10.4.2 Additional samples may test can be taken from by siphoning most of the same grab for other kinds overlying water without disturbing the surface of the sediment, then removing appropriate aliquots of the sediment analyses (see 10.2). Sediment temperature, interstitial for chemical analysis.

10.4.5 *Interstitial Water*—Interstitial water salinity, pH, and Eh can be recorded in (pore water), defined as the field. Qualitative description of water occupying the spaces between sediment might include color, texture, depth of oxidized layer, and presence of plants, animals, tracks, or burrows. Monitoring soil particles, is often isolated to provide either a matrix for toxicity testing or to provide an indication of the concentration or partitioning of chemicals within the sediment samples should be avoided, especially if matrix. Draft USEPA sediment equilibrium partitioning benchmarks (ESBs) are based on the presumption that the concentration of chemicals in the interstitial water are correlated directly to their bioavailability and, therefore, their toxicity (Di Toro et al. 1991, (70)). Of additional importance is associated with potentially hazardous contaminants in interstitial waters can be transported into overlying waters through diffusion, bioturbation, and resuspension processes (Van Rees et al. 1995, (153)). The usefulness of interstitial water sampling for determining chemical contaminants. A core contamination or toxicity will depend on the remainder study objectives and nature of the sediment-ins at the grab study site.

10.4.5.1 Isolation of sediment interstitial water can be sieved to provide accomplished by a macrobenthos sample.

10.4.3 The natural geochemical properties wide variety of test sediment collected from the field must methods, which are based on either physical separation or on diffusion/equilibrium. The common physical-isolation procedures can be within categorized as: (1) centrifugation, (2) compression/squeezing, or (3) suction/vacuum. Diffusion/equilibrium procedures rely on the tolerance limits movement (diffusion) of pore-water constituents across semipermeable membranes into a collecting chamber until an equilibrium is established. A description of the test species. The limits for the test species should be determined experimentally materials and procedures used in advance. Controls for such factors as particle size, organic content, salinity, etc. should be run if the limits are exceeded isolation of pore water is included in the test sediments reviews by Bufflap and Allen 1995, (154), Guide E 1391, and USEPA 2000, (64).

10.4.5.2 When relatively large volumes are required (>20 mL) for toxicity testing or chemical analyses, appropriate quantities of sediment are generally collected with grabs or corers for subsequent isolation of the interstitial water. Several isolation procedures, such as centrifugation (Ankley and Scheubauer-Berigan, 1995 (155)), squeezing (Carr and Chapman, 1995 (138)) and suction (Winger and Lasier, 1998 (156) ; Winger et al. (155)), have been used successfully to obtain adequate volumes for testing purposes. Peepers (dialysis) generally do not produce sufficient volumes for most analyses; however, larger sized peepers (500-mL volume) have been used for collecting interstitial water *in situ* for chemical analyses and organism exposures (Burton, 1992, (157) ; Sarda and Burton, 1995, (140)).

10.4.5.3 There is no one superior method for the isolation of interstitial water used for toxicity testing and associated chemical analyses. Factors considered in the selection of an isolation procedure may include: (1) volume of pore water needed, (2) ease of isolation (materials, preparation time, and time required for isolation), and (3) artifacts in the pore water caused by the isolation

procedure. Each approach has unique strengths and limitations (Bufflap and Allen, 1995 (158, 154); Winger et al. 1998, (156)), which vary with sediment characteristics, chemicals of concern, toxicity test methods, and desired test resolution (that is, data quality objectives). For suction or compression separation which use a filter or a similar surface, there may be changes to the characteristics of the interstitial water compared to separation using centrifugation (Ankley et al. 1994, (60); Horowitz et al. 1992, (159)). For most toxicity test procedures, relatively large volumes of interstitial water (for example, liters) are frequently needed for static or renewal exposures with the associated water chemistry analyses. While centrifugation can be used to generate large volumes of interstitial water, it is difficult to use centrifugation to isolate water from coarser sediment. If smaller volumes of interstitial water are adequate and logistics allow, the use of peepers which establish an equilibrium with the pore water through a permeable membrane may be desirable. If logistics do not allow placement of peeper samplers, an alternative procedure could be to collect cores which are can be sampled using side port suctioning or centrifugation. However, if larger samples of interstitial water are needed, it would be necessary to collect multiple cores as quickly as possible using an inert environment and centrifugation at ambient temperatures. See USEPA 2001 (134) and Guide E 1391 for additional detail regarding isolation of interstitial water.

10.4.5.4 There is no one superior method for the isolation of interstitial water for toxicity testing purposes. Each approach has unique strengths and limitations which vary with the characteristics of the sediment, the chemicals of concern, the toxicity test methods to be used, and the resolution necessary (that is, the data quality objectives). For suction or compression separation which use a filter or a similar surface, there may be changes to the characteristics of the interstitial water compared to separation using centrifugation (Ankley et al. 1994, (60) . For most toxicity test procedures, relatively large volumes of interstitial water (for example, liters) are frequently needed for static or renewal exposures with the associated water chemistry analyses. While centrifugation can be used to generate large volumes of interstitial water, it is difficult to use centrifugation to isolate water from coarser sediment. If smaller volumes of interstitial water are adequate and logistics allow, the use of peepers which establish an equilibrium with the pore water through a permeable membrane may be desirable. If logistics do not allow placement of peeper samplers, an alternative procedure could be to collect cores which are can be sampled using side port suctioning or centrifugation. However, if larger samples of interstitial water are needed, it would be necessary to collect multiple cores as quickly as possible using an inert environment and centrifugation at ambient temperatures. See USEPA 2001, (134) and Guide E 1391 for additional detail regarding isolation of interstitial water.

11. Quality Assurance and Quality Control

11.1 *Reference Sediment*—A reference sediment isIntroduction:

11.1.1 Developing and maintaining a clean sediment collected from laboratory Quality Assurance (QA) program requires an ongoing commitment by laboratory management and also includes the field that represents following: (1) appointment of a laboratory quality assurance officer with the test sediments in sedimental characteristics (for example, TOC; particle size, pH, Eh, salinity). This provides responsibility and authority to develop and maintain a QA program, (2) preparation of a Quality Assurance Project Plan with Data Quality Objectives, (3) prepreparation of written descriptions of laboratory Standard Operating Procedures (SOPs) for test organism culturing, testing, instrument calibration, sample chain-of-custody, laboratory sample tracking system, and (4) provision of potentially toxic adequate, qualified technical staff and non-toxic conditions. It suitable space and equipment to ensure reliable data (USEPA).

11.1.2 Quality Assurance (QA) practices within a testing laboratory should be handled in address all activities that affect the same manner as field collected test quality of the final data, such as: (1) sediment sampling and handling, (2) the source and condition of the test organisms, (3) condition and operation of equipment, (4) test conditions, (5) instrument calibration, (6) replication, (7) use of reference toxicants, (8) record keeping, and (9) data evaluation.

121.21.3 Quality Control (QC) practices, on the other hand, consist of the more focused, routine, day-to-day activities carried out within the scope of the overall QA program. For more detailed discussion of quality assurance, and general guidance on good laboratory practices related to testing, see USEPA 1993a, 1993c, 1995, 1978, 1979a, 1980a, 1980b, 1993b, and DeWoskin 1984, (132, 161, 162, 163, 164, 165, 166, 167, 168).

11.2 *Laboratory Spiked Test Sediment Performance-based Criteria* :

10.6.1 Test sediment can also be prepared in

11.2.1 The USEPA Environmental Monitoring Management Council (EMMC) recommended the laboratory by manipulating the properties use of control sediment. This can include adding various concentrations of toxic chemicals, highly toxic sediment, or complex waste mixtures (for example, sewage sludge) to the clean sediment (21). The toxicity of substances either dissolved performance-based methods in the interstitial water or adsorbed to sediment particles can be determined experimentally.

10.6.2 *Test Chemicals*—Chemicals experimentally added to sediment should be reagent-grade developing standards for chemical analytical methods (Williams 1993, (1302) or better, unless). Performance-based methods were defined by EMMC as a test on a formulation, commercial product, or technical-grade or use-grade material is specifically needed. Before a test is begun, monitoring approach which permits the following should be known about the chemical used: identities use of m appropriate methods that meet preestablished demonstrated performance standards. Minimum required elements of performance, such as precision, reproducibility, bias, sensitivity, and impurities, solubility and stability in test water, estimated toxicity to the test species and to humans, and recommended handling and disposal procedures.

10.6.3 *Stock Solution*—Toxic chemicals to detection limits should be tested in sediment are usually dissolved in a solvent to

TABLE 4 Recommended Test Conditions for Conducting Reference-Toxicity Tests (USEPA 1994a (1))

Parameter	Conditions
1. Test Type:	Water-only test
2. Dilution series:	Control and at least 5 test concentrations (0.5 dilution factor)
3. Toxicant:	Cd, Cu, ammonia, Sodium dodecyl sulfate (SDS)
4. Temperature:	15°C <i>E. estuarius</i> and <i>R. abronius</i> 20°C <i>A. abdita</i> 25°C <i>L. plumulosus</i>
5. Light quality:	Chambers should be kept in the dark covered with opaque material (USEPA 1994a (1)). Alternatively USEPA-USACE (2001(2)) recommends a 500 to 1000 lux light intensity at a 16:8 light:dark cycle for <i>L. plumulosus</i> in long-term tests (Annex A2).
6. Photoperiod:	24 h dark
7. Salinity:	28 ‰ for <i>A. abdita</i> and <i>R. abronius</i> 20 ‰ for <i>E. estuarius</i> 5 or 20 ‰ for <i>L. plumulosus</i> . Alternatively, the salinity of the overlying water can be adjusted to the salinity of the pore-water at the site of interest in tests with <i>E. estuarius</i> or <i>L. plumulosus</i> . If tests are conducted at different salinities, additional tests are required to determine comparability of results (sections 1.4 and 1.10).
8. Renewal of water:	None
9. Age of organisms:	<i>A. abdita</i> : 3 to 5 mm (no mature males or females) <i>E. estuarius</i> and <i>R. abronius</i> : 3 to 5 mm <i>L. plumulosus</i> : 2 to 4 mm (no mature males or females; USEPA 1994a (1)). Alternatively, USEPA-USACE (2001(2)) recommend testing <i>L. plumulosus</i> in a range of 0.25 to 0.60 mm in length in long-term tests (Annex A2).
10. Test chamber:	250-ml to 1-L glass beaker or jar
11. Volume of water:	80 % of chamber volume
12. Number of organisms/ chamber:	$n = 20$ if 1 replicate; $n = 10$ (minimum) if >1 per replicate.
13. Number of replicate chambers/treatment:	1 minimum; 2 recommended
14. Aeration:	Recommended; but not necessary if >90 % dissolved oxygen saturation can be achieved without aeration (USEPA 1994a (1)). Alternatively USEPA-USACE (2001 (2)) recommends that dissolved oxygen should be maintained at >60 % saturation (>4.4 mg/L). Culture water, surface water, site water, or reconstituted water
15. Dilution water:	Salinity, pH, and dissolved oxygen, at the beginning and end of a test. Temperature daily.
16. Water quality:	96 h
17. Test duration:	Survival (LC50); Reburial (EC50) optional for <i>E. estuarius</i> and <i>R. abronius</i>
18. Endpoint:	90 % control survival
19. Test acceptability:	

form a stock solution that is then added to the sediment. The concentration specified and stability of the chemical in the stock solution method should be determined before the beginning of the test. If the chemical is subject demonstrated to photolysis, meet the stock solution should be shielded from the light both before and during the process of mixing into the sediment.

10.6.4 The preferred solvent is prepared toxicity test sea water at the test salinity. If a substance is insoluble in salt water, deionized water may be used, if salinity is adjusted accordingly if necessary. Several techniques have been specifically developed performance standards.

11.2.2 In developing guidance for preparing aqueous stock solutions of slightly soluble materials culturing (22). The minimum necessary amount of a strong acid or base may *L. plumulosus*, it was determined that no single method has to be used in the preparation to culture organisms (USEPA-USACE 2001 (2)). Success of an aqueous stock solution, but such reagents might affect the pH of stock solutions appreciably. Use of a more soluble form of the test material, such as chloride or sulfate salts of organic amines, sodium or potassium salts of phenols and organic acids, and chloride or nitrate salts of metals, might affect relies on the pH more than the use health of the necessary minimum amount culture from which organisms are taken for testing. Having healthy organisms of a strong acid or base.

10.6.5 If a solvent other than water known quality and age (that is, size) for testing is used, it should be of reagent grade. Its concentration in the sediment should be kept key consideration relative to culture methods. Therefore, a minimum, and should be low enough that it does not affect the test species. Triethylene glycol performance-based criteria approach is often a good organic solvent for preparing stock solutions because of its low toxicity to aquatic animals, low volatility, and high ability to dissolve many organic chemicals. Other water-miscible organic solvents such as methanol, ethanol, or acetone may be used, but they might affect total organic carbon levels, introduce toxicity, alter the geochemical properties of the sediment, or stimulate undesirable growths of microorganisms. Acetone is highly volatile and might leave the system more readily than methanol or ethanol. A surfactant preferred method by which individual laboratories should not be used in the preparation of a stock solution because it might affect the bioavailability, form, and toxicity of the test material.

10.6.6 If a solvent other evaluate culture health, rather than water is used, both a solvent control with control sediment and a

clean sediment control must be included in the test. The solvent control must contain the highest concentration of solvent present in sediment in any other treatment and must use solvent from the same batch used control-based criteria approach. Performance-based criteria were chosen to make the stock solution. The percentage of allow each laboratory to optimize culture methods that provide organisms that show signs of stress, such as inability to rebury at the end of the test, or death, must be 10% or less in both controls. Greater than 10% mortality in the controls or obvious sublethal stress in 10% or more of the control animals invalidates the test (see 12.2.2).

10.6.7 If the test contains both a clean sediment (negative) control produce reliable and a solvent control, the survival, reburial, or other endpoint determined in the two controls should be compared. If a statistically significant difference in any endpoint is detected between the two controls, only the solvent control may be used for meeting the acceptability of the comparable test and for calculation of results. The negative control might provide additional information on the general health of the organisms tested. If no statistically significant difference is detected, the data from both controls may be pooled Performance criteria for meeting the acceptability of the test culturing and as the basis for calculation testing *L. plumulosus* are listed in Table A2.3 of results. Annex A2.

11.3 Test Concentration(s) Facilities, Equipment, and Test Chambers:

10.7.1 If the

11.3.1 Separate test organism culturing and testing areas must be provided to calculate an LC50 or other effect level, the test concentrations avoid loss of cultures because of cross-contamination. Ventilation systems should bracket the predicted effect level. The prediction might be based on the results designed and operated to prevent recirculation or leakage of a test on the same air from chemical analysis laboratories or a similar sample storage and preparation areas into test material on the same organism culturing or a similar species. If a useful prediction is not available, it is usually desirable to conduct a range-finding test in which the organisms are exposed to a control sediment testing areas, and three or more concentrations of the test material that differ by a factor of ten.

10.7.2 If necessary, concentrations above aqueous solubility can from sediment testing laboratories and sample preparation areas into culture areas.

11.3.2 Equipment for temperature control should be adequate to maintain recommended test-water temperatures. Recommended materials should be used because in the real world organisms are sometimes exposed to concentrations above solubility and because solubility is often not well known. The toxicity fabrication of the test material equipment which comes in sediments might be quite different from contact with the toxicity in water borne exposures.

10.7.3 Bulk sediment chemical concentrations might be normalized to factors other than dry weight. For example, concentrations of non-ionic organic compounds might be normalized to organic carbon content.

10.7.4 In some (usually regulatory) situations, it is only necessary to determine (a) whether or overlying water.

11.3.3 Before a specific concentration of sediment test material is acutely toxic to the test species or (b) whether the LC50 is above or below a specific concentration. For example, the specific concentration might be the concentration occurring conducted in a particular sediment, or the concentration in new facility, a dredge material to “non-contaminant” test should be deposited at a disposal site. When there is only interest conducted in which all test chambers contain a particular concentration, it might only be necessary control sediment and overlying water. This information is used to demonstrate that concentration, and the negative facility, control sediment, water, and solvent controls handling procedures provide acceptable responses of test organisms (section 11.14).

11.4 A Test Organisms—The organisms should appear healthy, behave normally, feed well, and have low mortality ion cultures of *L. plumulosus*, during holding (for example, <20% for 48 h before the start of a test), and in test controls. Test Material organisms should be positively identified to species. Obtaining wild populations of organisms for testing should be avoided unless the ability of the wild population to cross-breed with existing laboratory populations has been determined (section 12.3.3.9).

11.5 Water—The quality of water used for organism culturing and testing is extremely important. Overlying water used in testing and water used in culturing organisms should be uniform in quality. Acceptable water should allow satisfactory survival, growth, or reproduction of the test organisms. Test organisms should not show signs of disease or apparent stress (for example, discoloration, unusual behavior). See section 7.1 for additional details.

11.6 Sample Collection and Storage— Sample holding times and temperatures should conform to conditions described in Section 10.

11.7 Test Conditions—It is desirable to measure temperature continuously in at least one chamber during each test. Temperatures should be maintained within the limits specified for each test. Dissolved oxygen, salinity, conductivity (particularly when salinity is <1 ‰), ammonia, and pH should be checked in accordance Annex A1 and Annex A2.

11.8 Quality of Test Organisms:

10.8.1 If test organisms are obtained from culture, reference-toxicity tests should be conducted on all test organisms using procedures outlined in section 11.16 (at a toxic sediment, or a solution minimum, one test every six months; Table 4). If reference-toxicity tests are not conducted monthly, the lot of a chemical can be added organisms used to sediment and evenly distributed by thorough hand mixing, by use of start a rolling mill, or by adding the sediment test material to should be evaluated using a slurry reference toxicant (USEPA 1994a (1)).

11.8.2 The quality of test organisms obtained from an outside source, regardless of whether they are from culture or collected

from the field, should be verified by conducting a reference-toxicity test concurrently with the sediment that is allowed to settle. The test material might also be added to water flowing over or through (USEPA 1994a (1)). For cultured organisms, the supplier should provide data with the shipment describing the history of the sensitivity of organisms from the same source culture. For field-collected organisms, the supplier should provide data with the shipment describing the collection location, the time and allowed to partition onto date of collection, the sediment. Other methods water salinity and temperature at the time of mixing might also be appropriate provided collection, and collection site sediment for holding and acclimation purposes. If the supplier has not conducted reference toxicity tests with the test material organism, it is shown to be evenly distributed in the sediment.

10.8.2 Modifications responsibility of the mixing technique might be necessary testing laboratory to allow time conduct these reference toxicity tests (section 11.14.1).

11.8.3 *Leptocheirus plumulosus* for chronic testing can be obtained from laboratory cultures (USEPA-USACE 2001 (2)) or from commercial sources (Table 5). It is likely to be impractical to obtain test-size neonates directly from a supplier because of their sensitivity to physical disturbances and their rapid growth. Instead, test material laboratories will likely want to establish their own cultures of *L. plumulosus* from which to harvest neonates (section 12.5). It is desirable to determine the sensitivity of *L. plumulosus* obtained from an outside source. For cultured organisms, the supplier should provide data with sediment. If tests are repeated, mixing conditions such as duration the shipment describing the history of the sensitivity of organisms from the same source culture. For field-collected organisms, the supplier should provide data with the shipment describing the collection location, the time and temperature date of mixing, collection, the water salinity and temperature at the time of mixing before collection, and collection site sediment for holding and acclimation purposes. The supplier should also certify the species identification of the test organisms and provide the taxonomic references (for example, Shoemaker, 1932 (169); Bousfield, 1973 (171)) or name(s) of the taxonomic expert(s) consulted.

11.8.4 All organisms in a test must be from the same source (Section 12). Organisms may be obtained from laboratory cultures or from commercial or government sources (section 11.8.3). The test organisms used should be kept constant, unless time after spiking is identified using an experimental variable. Care appropriate taxonomic key, and verification should be taken documented. The use of field-collected *L. plumulosus* to ensure that a test material added to sediment start cultures is discussed in section 12.3. Obtaining organisms from wild populations is useful for enhancement of genetic diversity of existing cultures or to establish new cultures. (McGee et al. 1998 (170)) found seasonal variability in sensitivity to cadmium in field-collected *L. plumulosus*. Therefore, field-collected *L. plumulosus* should not be used for toxicity testing unless organisms are cultured through several generations in the sediment. If necessary, sub-samples laboratory. In addition, the ability of the sediment within a mixing container can wild population of sexually reproducing organisms to cross-breed with the existing laboratory population should be determined (Duan et al., 1997 (173)). Sensitivity of the wild population to determine degree select contaminants should also be documented.

11.9 *Quality of mixing and homogeneity:*

11. Test Organisms

11.1 *Species*—The species Food—Problems with the nutritional suitability of infaunal amphipod to the food will be used reflected in the survival, growth, or reproduction of *L. plumulosus* in cultures (section 12.5). Additionally, survival in sediment toxicity test should tests conducted with *A. abdita* and *L. plumulosus* may be affected by the nutritional suitability of food provided during holding and acclimation (USEPA 1994a (1)).

11.10 *Test Acceptability*;—Table A1.3 in Annex A1 and Table A2.3 in Annex A2 outline requirements for acceptability of tests. An individual test materials, tolerance to ecological conditions (for example, may be conditionally acceptable if temperature,

**TABLE 5 Sources of Starter Cultures of Test Organisms for
*Leptocheirus plumulosus***

<p>Aquatic Biosystems, Inc. 1300 Blue Spruce Road, Suite C Fort Collins, Colorado 80524 Scott Kellman phone: 800/331-5916; fax: 970/484-2514 email: SRK@riverside.com</p>
<p>Chesapeake Cultures, Inc. P.O. Box 507 Hays, Virginia 23702 Elizabeth Wilkins, President phone: 804/693-4046; fax: 804/694-4703 email: growfish@c-cultures.com website: www.c-cultures.com</p>
<p>Aquatic Research Organisms P.O. Box 1271 Hampton, New Hampshire 03842-1271 Stan Sinitski or Mark Rosenqvist phone: 800/927-1650; fax: 603/926-5278 website: www.aro-central.com</p>

salinity, DO, and grain size), ecological importance, other specified conditions fall outside specifications, depending on the degree of the departure and ease the objectives of handling in the laboratory tests. The acceptability of a test will depend on the experience and type professional judgment of sediment being tested or the laboratory analyst and the reviewing staff of the regulatory authority. Any deviation from test specifications should be implemented might dictate selection of noted when reporting data from a particular species. Ideally, species or genera with wide geographical distributions test.

11.11 Analytical Methods:

11.11.1 All routine chemical and physical analyses for culture and testing water, food, and sediment should include established quality assurance practices (Van Rees et al. 1991, Bufflap and Allen 1995, Ankley and Subauer-Bergian 1995, **(153-156)**).

11.11.2 Reagent containers should be selected, so that test results can dated when received from the supplier and the shelf life of the reagent should not be compared among laboratories with similar species. Species used exceeded. Working solutions should be identified with an appropriate taxonomic key, dated when prepared and identifications the recommended shelf life should not be verified by a taxonomic authority. The annexes to this guide give guidance as to requirements exceeded.

11.12 Calibration and methods of handling Standardization :

11.12.1 Instruments used for various species routine measurements of amphipods. Use of chemical and physical characteristics such as pH, DO, temperature, salinity, and conductivity should be calibrated before use each day according to the species-listed instrument manufacturer's procedures as indicated in the annexes is encouraged to increase comparability of results.

11.12.1 *Rhepoxynius abronius* is general section on quality assurance (see USEPA Methods 150.1, 360.1, 170.1, and 120.1, (125)). Calibration data should be recorded in a free-burrowing amphipod that has been successfully used permanent log.

11.12.2 A known-quality water should be included in sediment toxicity testing since the late 1970's analyses of each batch of water samples (for example, salinity, conductivity (particularly when salinity is <1 (3)‰)). It is desirable to include certified standards in the analysis of water samples.

11.13 Replication and Test Sensitivity— The sensitivity of this species to salinities less than 25 g/kg limits its use to testing sediments from marine areas, but sediment tests will depend in part on the large data base that has been developed for number of replicates/treatment, the response significance level selected, and the type of *R. abronius* to statistical analysis. If the variability remains constant, the sensitivity of a variety test will increase as the number of replicates is increased. The minimum recommended number of replicates varies with the objectives of the test and chemicals establishes its usefulness the statistical method used for analysis of the data (Annex A1 and Annex A2).

11.14 Demonstrating Acceptable Performance :

11.14.1 Intralaboratory precision, expressed as a coefficient of variation, of the range for each type of test species as well as to be used in a laboratory can be determined by performing five or more tests with different batches of test organisms, using the same reference species for comparing toxicant, at the same concentrations, with the same test conditions (for example, the same test duration, type of other species. Species water, age of test organisms, feeding), and same data analysis methods. A reference-toxicant concentration series (0.5 or higher) should be selected that will provide partial mortalities at two or more concentrations of the genus *Rhepoxynius* are widely distributed on test chemical (section 10.3.3). Information from previous tests can be used to improve the West Coast design of N subsequent tests to optimize the dilution series selected for testing.

11.14.2 Before conducting tests with a potentially contaminated sediment, it is strongly recommended that the laboratory conduct the tests with control sediment(s) alone. Results of these preliminary studies should be used to determine if the use of the control sediment and other test conditions (that is, water quality) result in acceptable performance in the tests as outlined in Annex A1 and Annex A2.

11.14.3 Laboratories should demonstrate that their personnel are present on able to recover an average of at least 90 % of the E organisms from whole sediment. For example, test organisms could be added to control or test sediments, and recovery could be determined after 1 h Tomasovic et al. 1995, **(23)(174)**.

11.15 Documenting Ongoing Laboratory Performance:

11.15.1 Outliers, which are data falling outside the control limits and trends of increasing or decreasing sensitivity are readily identified. If the data from a given test falls outside the "expected" range (for example, ± 2 SD), the sensitivity of the organisms and the credibility of the test results may be suspect. In this case, the test procedure should be examined for defects and should be repeated with a different batch of test organisms.

11.15.2 A sediment test may be acceptable if specified conditions of a reference-toxicity test fall outside the expected ranges (section 11.10). Specifically, a sediment test should not be judged unacceptable if the LC50 for a given reference-toxicity test falls outside the expected range or if control survival in the reference-toxicity test is <90 %. All the performance criteria outlined in Annex A1 and Annex A2) must be considered when determining the acceptability of a sediment test. The acceptability of the sediment test would depend on the experience and judgment of the investigator and the regulatory authority.

11.15.3 Performance should improve with experience, and the control limits should gradually narrow, as the statistics stabilize. However, control limits of ± 2 SD, by definition, will be exceeded 5 % of the time, regardless of how well a laboratory performs. For this reason, good laboratories that develop very narrow control limits may be penalized if a test result which falls just outside the control limits is rejected de facto. The width of the control limits should be considered in decisions regarding rejection of data (Section 15).

11.16 Reference-toxicity Testing :

11.16.1 Reference-toxicity tests should be conducted in conjunction with sediment tests to determine possible changes in condition of a test organism (Lee, 1980 (175)). Water-only reference-toxicity tests on cultured organisms should be conducted least every 6 months with laboratory-cultured organisms and should be performed on each batch of field-collected organisms used for testing. Deviations outside an established normal range may indicate a change in the condition of the test organism population. Results of reference-toxicity tests also enable interlaboratory comparisons of test organism sensitivity (USEPA 1994a (1)).

11.16.2 Reference toxicants such as cadmium (available as cadmium chloride (CdCl₂), copper (available as copper sulfate (CuSO₄), ammonia, and sodium dodecyl sulfide (SDS) are suitable for use. No one reference toxicant can be used to measure the condition of test organisms in respect to another toxicant with a different mode of action (Lee, 1980 (175)). However, it may be unrealistic to test more than one or two reference toxicants routinely.

11.16.3 Test conditions for conducting reference-toxicity tests with *A. abdita*, *E. sinuatus estuarius*, *L. plumulosus*, and *E. washingtonianus* have been successfully used. R. abronius are outlined in Table 4.

11.16.4 Basedime on 96-h, water-only reference-toxicity tests at 20 ‰ with neonate (24, 25) *L. plumulosus*, one should expect a mean LC50 value for cadmium of about 0.5 mg/L (range: 0.2 mg/L to 0.7 mg/L) and LC50 values for total ammonia between 25 mg/L and 60 mg/L (DeWitt et al., 1997a and haustoriids in general are more abundant than phoxocephalids on the East(7)). At 5 ‰, one should expect a mean LC50 value for cadmium of about 0.05 mg/L (range: 0.01 mg/L to 0.09 mg/L) and Gulf Coasts LC50 values for total ammonia between 37 mg/L and 53 mg/L (Emery et al., 1997 (8); Moore et al., 1997 (26). Their sand-burrowing habits, availability, ease). Kohn et al. 1994 (24) report 96-h LC50 values for total ammonia (mg/L) of handling, tolerance to a wide range of salinity 79 for *R. abronius*, 126 (estimated) for *E. estuarius*, and temperature, and ecological importance as probable prey 50 for *A. abdita*. Northwestern Aquatic Sciences (NAS) reports an average total ammonia (mg/L) 96-h LC50 of shorebirds 138 for *E. estuarius* (n=5), and fishes mean (n=20) 96-h LC50 values for cadmium of 1.1 for (27) *R. abronius*, 2.4 for make them good candidates *E. estuarius*, and 0.55 for test species, especially *A. abdita*. The NAS laboratory also reports a mean (n=20) sodium dodecyl sulfate (SDS; mg/L) 48-h LC50 of 12.6 for estuarine areas:

11.1.3 A variety *A. abdita*. All of other benthic amphipod species have been used successfully to test the toxicity of marine and estuarine sediments using similar methods or NAS tests were conducted at 28 ‰ except for the same method described here. These species include: *E. estuarius* cadmium test, which was conducted at 12 ‰ (Michele Redmond, Northwestern Aquatic Sciences, Newport, OR, personal communication). MEC Analytical has observed 96-h LC50 values for cadmium (mg/L) of 0.75 for *R. abronius*, 7.1 for *E. spinicorne* *E. estuarius*, 0.56 for *A. abdita*, and the freshwater amphipods 3.9 for *Hyalella azteca* *L. plumulosus* (David Moore, MEC Analytical, Carlsbad, CA, personal communication). DeWitt et al. (1992a (6)) provides additional data on cadmium reference toxicity tests for *Pontoporeia hoyi* *R. abronius*, *E. estuarius*, *A. abdita*, and *L. plumulosus*.

11.1.4 The environmental requirements and sensitivity of a prospective test species of amphipod to test materials and to various sediment characteristics should be established before it is widely used in toxicity tests. The tolerance of a test species to variations in sediment characteristics such as particle size distribution, organic enrichment, and interstitial water salinity should be established before responses can be ascribed to contaminant effects. Choice of the scale of the test chamber, density of test organisms, temperature, salinity, and control sediment might have to be modified to accommodate the requirements of the test species. Required modifications should be based on conditions at the natural habitat of the species.

11.1.5 The sensitivity of a prospective new test species of amphipod should be compared with a reference species such as *R. abronius* before the new species is used in routine toxicity testing. A 96-h reference toxicity test using water only could eliminate the relative effects of sediment particle size and other sediment characteristics (see 11.5.4). The test should be set up as in Section 13, but without the addition of sediment. A non-ionic organic compound whose binding properties are not affected by salinity could be used to compare species at different salinity levels (example: polynuclear aromatic hydrocarbons such as fluoranthene). It might be desirable to also test a metal such as cadmium. Any factor (such as salinity, pH, redox state, carbonates, or sulfides) that might affect the toxicity or bioavailability of the reference toxicant should be held constant.

11.1.6 If tube-building amphipods are used in sediment toxicity testing, it should be kept in mind that the amphipods might not be directly in contact with test sediment after their tubes are built, and they might pump overlying water through their tubes rather than utilizing interstitial water. They might feed on particulate materials that either are suspended in the water column or have settled on the sediment surface, while burrowing species might feed on particles or meiofauna found within the sediment. Thus tube builders and burrowing species might have different routes of exposure to adsorbed or dissolved sediment contaminants. Amphipods that emerge from the sediment and either swim in overlying water or crawl on the sediment surface might not be continually exposed to the test sediment.

11.2 Age—All organisms should be as uniform as possible in age and size. The age or size class

11.17 Record Keeping—Section 13.1 outlines recommendations for a particular species should be chosen so that sensitivity to test materials is not affected by state of maturity, reproduction, seasonality, etc. (see Annexes for species, specific requirements).

11.3 Source—All individuals in a test should be from the same source, because different populations of the same species might have different acute sensitivities to contaminants. Marine amphipods are usually obtained directly from a wild population in a clean area, although attempts have been made to culture some species. Collecting permits for field collected amphipods might be required by some local recorded keeping (that is, data files, chain-of custody).

12. Collection, Culturing, and state agencies:

11.3.1 If test organisms are cultured or held for an extended period of time in the laboratory, the response of laboratory-held organisms to test materials should be compared to that of animals freshly collected from the field to assure that laboratory stresses do not affect their sensitivity to test materials (19):

11.4 Collection and Handling:

11.4.1 Amphipods should be handled as little as possible. When handling is necessary, it should be done carefully, gently, and quickly so that organisms are not unnecessarily stressed. Amphipods that touch dry absorbent surfaces or are injured during handling should be discarded.

11.4.2 *Collection*—Amphipods can be collected intertidally with a shovel or subtidally with a small biological dredge or a grab. Sediment containing amphipods can be gently sieved to separate the amphipods. The amphipods can then be collected with a dipnet and transferred to and allowed to bury in sieved sediment from the amphipod collection site. Sieves and containers used to collect and transport amphipods should be marked “live only” and should never be used for working with formalin or any other toxic materials. Water used for sieving should be at the same temperature and salinity as bottom water at the collection site. Infaunal amphipods should be held in sediment during transport to the laboratory, and should be kept at or near collection site temperature or below. During a long transport, it might be necessary to keep containers of sediment and amphipods in coolers and to provide aeration. Collection site sediment should be saved for control, acclimation, and reburial sediment.

11.4.3 *Holding*—Amphipods should be fully acclimated to the test temperature and salinity by holding them in the laboratory prior to their use in a toxicity test. Amphipods should be collected from the field three or four days before use, but field-collection animals should not be held in the laboratory for more than two weeks before the initiation of a test.

11.4.3.1 In the laboratory, amphipods can be counted into holding containers with clean sieved sediment to ascertain whether sufficient numbers have been collected. Amphipods should be sieved from transport sediment and gently washed into a clean dish for counting. Active, apparently healthy amphipods can be picked up and removed from detritus with a wide-mouthed bulb pipette and transferred to sieved collection site sediment, into which the amphipods should quickly bury. Enough amphipods should be collected to provide at least one third more individuals than are required for the test. During counting, the temperature of the water containing the amphipods must not exceed the amphipods’ tolerance limit, and should remain close to the holding temperature. The holding containers should be provided with flowing or aerated sea water at or near the test temperature and salinity. If changes in temperature and salinity are necessary to bring amphipods from the collection site conditions to the test conditions, adjustments should be made gradually to allow amphipods to acclimate. Healthy burrowing amphipods will usually remain in the holding sediment until the initiation of the test, and can be easily retrieved for setup. Supplementary feeding during the acclimation period might or might not be necessary, as some amphipods will find food in the holding sediment (see species-specific annexes). Any individuals that fail to bury or make tubes (if they are tube builders) in holding sediment or that appear unhealthy during holding should be discarded. The temperature and salinity of the water in the holding containers should be monitored daily. Maintaining Test Organisms

12.1 *Quality*—All amphipods used in Life History:

12.1.1 *Ampelisca abdita*—*A. abdita* is a test must be of acceptable quality. A qualified tube-building amphipod taxonomist must be consulted to ensure that the animals in the test population are all of family Ampeliscidae. It occurs on the same species.

11.5.1 Amphipods Atlantic coast from central Maine to central Florida, although it is also found in holding containers should be checked daily before the initiation eastern portion of a test. Individuals that emerge from the Gulf of Mexico (Bousfield, 1973 (171)). On the Pacific coast, it is present in San Francisco Bay, CA (Nichols et al. 1985 (176); Hopkins, 1986 (177)). They are small (adult length 4 to 8 mm), laterally compressed amphipods. Healthy animals are opalescent pink and appear dead or will remain tightly curled, whereas unhealthy should animals tend to be discarded. If greater than 5 % translucent white, and may uncurl (USEPA 1994a (1)). Often dominant members of the amphipods emerge benthic community, *A. abdita* forms thick mats of tubes with amphipod densities up to 110,000/m², and appear often a dominant food source for bottom-feeding fish (Richards, 1963 (178)). The tubes are narrow and about 2 to 3 cm in length. A filter feeder, *A. abdita* feeds on both particles in suspension and those from surficial sediment surrounding the 48 h preceding the test, the entire group should be discarded tube. *Ampelisca abdita* is euryhaline, and not used has been reported in waters that range in salinity from fully marine to 10 ‰ (Hyland, 1981 (179)). Laboratory tests have shown the test:

11.5.2 *Analysis salinity application range* of *A. abdita* in sediments is from 0 to 34 ‰ when the test organisms for the test material, if salinity of overlying water is 28 ‰ (Weisberg et al. 1992, (180)). This species generally inhabits sediments from fine sand to mud and silt without shell fragments, although it might can also be present found in relatively coarser sediments with a sizeable fine component. It is often abundant in sediments with a high organic content. Analysis of historical data shows little effect of sediment grain size on survival of *A. abdita* during 10-d sediment toxicity tests (Long and other chemicals to which exposure might have occurred; Buchman, 1989 (17) ; Weisberg et al. 1992, (180)). There is desirable. Amphipods evidence that sediments with >95 % sand may be used without analysis elicit excessive mortality (John Scott, personal communication in USEPA 1994a (1)). *Ampelisca abdita* have been collected at water temperatures ranging from -2 to 27°C (USEPA 1994a (1)). Reproduction patterns of chemical concentration if *A. abdita* vary geographically. In the amphipods are obtained from colder waters of its range, *A. abdita* produces two generations per year, an over-wintering population that is monitored for chemical contamination (see 10.2) broods in the spring, and known a second that breeds in mid- to be free late-summer (Mills, 1967 (181))

). In warmer waters south of toxicants, and they Cape Hatteras, NC, breeding might be continuous throughout the year (Nelson, 1980 (182)). Juveniles are released after about two weeks in clean, uncontaminated water the brood pouch. Juveniles take about 40 to 80 d to become breeding adults at 20°C (Mills, 1967 (181); Scott and Redmond, 1989 (12); Redmond et al., 1994 (183)).

12.1.2 *Eohaustorius estuarius*— *E. estuarius* is a free-burrowing amphipod in the family Haustoriidae. It is found on protected and semi-protected beaches from the lower intertidal to shallow subtidal waters exclusively on the Pacific coast from British Columbia south to central California (Environment Canada, 1992 (5); USEPA-USACE 2001 (2)). They are stout (adult size range 3 to >5 mm) cup- or bell-shaped, dorsally compressed amphipods that are grayish-brown or yellowish-brown in color (Environment Canada, 1992 (5)). *Eohaustorius estuarius* are thought to be deposit feeders. It is an estuarine species and has been reported in areas where pore-water salinity ranges from 1 to 35 ‰ (Environment Canada, 1992 (5), USEPA 1994a (1); Michele Redmond, Northwest Aquatic Sciences, Newport, OR; personal communication). Laboratory studies have shown a salinity application range in control sediments for *E. estuarius* from 0 to 34 ‰. *Eohaustorius estuarius* inhabits clean, medium-fine sand with some organic content. The species has exhibited acceptable (that is, >90 %) survival when exposed to clean sediments with a wide range of grain sizes, with generally little effect on survival whether coarse-grained or fine-grained (that is, predominantly silt and clay) clean sediments are used (Environment Canada, 1992 (5)). Environment Canada (1998 (184)) reported that *E. estuarius* can tolerate up to 70 % clay in sediment toxicity tests. However, some correlation between survival and grain size exists (DeWitt et al., 1989 (11)). *Eohaustorius estuarius* has been collected from water temperatures from 0 to 23°C (USEPA-USACE 2001 (2)). *Eohaustorius estuarius* apparently has an annual life cycle (Environment Canada, 1992 (5); DeWitt et al. 1989 (11)). Gravid females are abundant in intertidal sediments from February through July. However, reproduction might occur year-round because juveniles are found throughout most of the experimental design specifically requires use year (DeWitt et al., 1989 (11)).

12.1.3 *Leptocheirus plumulosus*— *L. plumulosus* is a burrow-building member of that population.

H.5.3 Survival of amphipods in control sediment during the test family Aoridae. It is an infaunal amphipod found in subtidal portions of Atlantic Coast brackish estuaries from Cape Cod, Massachusetts, to northern Florida (Bousfield, 1973 (171); DeWitt et al., 1992a (6)). It is common in protected embayments, but has been collected in channels of estuarine rivers at water depths up to 13 m (Shoemaker, 1932 (169); Holland et al., 1988 (185); Schlekot et al., 1992 (13)). In Chesapeake Bay, densities of *L. plumulosus* can reach 40,000/m² (McGee 1998, (172)). *L. plumulosus* is a relatively large amphipod (adult length up to 13 mm) with a cylindrically shaped body that is brownish-gray in color. A distinguishing feature is a series of dark bands or stripes that cross the dorsal surface of the population parents and other factors. If a mean of greater than 10 % mortality occurs in the test, the test must be repeated on the sediment surface (DeWitt et al., 1992a (6)). Studies have shown that *L. plumulosus* population abundance in Chesapeake Bay is negatively correlated with sediment contamination (Holland et al., 1988 (185); McGee and Fisher 1999 (42); McGee et al. 1999 (41)). Thus, this amphipod would appear to be considered invalid.

H.5.4 Reference toxicants might a good candidate to be useful for assessing the quality an environmental indicator. *L. plumulosus* is found in both oligohaline and sensitivity mesohaline regions of east coast estuaries; ambient water salinity at collection sites has ranged from 0 to 15 ‰ (Holland et al., 1988 (185); DeWitt et al., 1992a (6); Schlekot et al., 1992 (13), 1994(186)). Laboratory studies have demonstrated that *L. plumulosus* 28-d test organisms, and can be employed using 96-h toxicity tests without conducted at salinity values ranging from 1 to 35 ‰ (A2.4; Schlekot et al., 1992 (13); DeWitt et al., 1992a (6), 1997a (7); Emery et al., 1997 (8)). This amphipod is most often found in fine-grained sediment with a relatively high proportion of particulate organic material, although it has been collected in fine sand with low organic content (Jordan and Sutton, 1984(188); Holland et al., 1988 (185); Marsh and Tenore, 1990(189); DeWitt et al., 1992a (6); Schlekot et al., 1992 (13); 1994 (186)). Laboratory studies with *L. plumulosus* revealed no effect of sediment grain size on survival in control sediment containing 5 to generate LC50 values (see H.1.4).

H.5.4.1 Reference toxicants 100 % silt-clay content (DeWitt et al., 1997a (7)). However, Emery et al. (1997 (8)) found significantly reduced survival in sediments in which clay content exceeded 84 %. Populations of *L. plumulosus* can be useful seasonally ephemeral with major population growth in fall and spring and large population declines in the summer (Holland et al., 1988 (185); Marsh and Tenore, 1990 (189); McGee, 1998 (172)). This pattern appears to be driven by changes in temperature and food availability and subsequent effects on life history traits (Marsh and Tenore, 1990 (189); McGee, 1998 (172)). Short-term population fluctuations are also a function of different populations or species of amphipods, or seasonal variation in its relatively short generation time (DeWitt et al., 1992a (6)). At 28°C in the laboratory, the age of the first brood release is about 24 d (DeWitt et al., 1992a (6)).

12.1.4 *Rhepoxynius abronius* is a field-collected population. Such assessment free-burrowing amphipod in the family Phoxocephalidae. It occurs on the Pacific Coast from Puget Sound, WA, to central California in lower intertidal and nearshore subtidal zones to depths of 274 m offshore (Environment Canada, 1992 (5); Lamberson and Swartz, 1988 (187); Kemp et al. 1985 (190); Barnard and Barnard, 1982 (191)). Densities in the field are reported to range from 150 to 2200/m² (Lamberson and Swartz, 1988 (187); Swartz et al. 1985 (10)). It is usually collected simultaneously a medium-sized (adult length from 3 to >5 mm) amphipod with a stout, somewhat rounded body shape. Color may range from salmon pink to yellowish, grayish-brown to white with a pinkish-brown hue (Environment Canada, 1992 (5)). *Rhepoxynius abronius* is a meiofaunal predator, but it also ingests sedimentary organic material (Oakden 1984 (10)). In the toxicity test. Many chemicals field, *R. abronius* is found where pore-water salinity is no lower than 20 ‰ (Environment Canada, 1992 (5)). Laboratory tests have indicated that salinities below 25 ‰ may

be toxic to *R. abronius* (Swartz et al., 1985 (10)). *Rhepoxynius abronius* should therefore normally not be chosen as reference toxicants the test species when the sediment pore water is <25 % (Swartz et al., 1985 (10)). PSEP (1995 (192)) outlines a procedure for adjusting the pore-water salinity of samples with a salinity <25 % by adding appropriately saline overlying water to the test chamber on the day before the start of the test, mixing sediment and overlying water, and allowing the material to settle overnight under aeration. The resultant overlying water is either retained or about 75 % replaced with fresh dilution water at 28 % (PSEP 1995 (192)). While this manipulation should result in an acceptable salinity for tests with *R. abronius*, the influence of this manipulation on the bioavailability of contaminants in the sediment sample is uncertain. *Rhepoxynius abronius* naturally inhabits clean, fine, sandy sediments. A number of studies have shown some reduction in survival when this species is held in very fine-grained (predominantly silt and clay) sediment (DeWitt et al., 1988 (193); Long et al., 1990 (28); McLeay et al., 1991 (194)). Normally collected at temperatures ranging from 8 to 16°C, *R. abronius* has been proven survived at temperatures ranging from 0 to 20°C under laboratory conditions. Reproduction of the overall quality of any species or test results. A reference toxicant *R. abronius* is likely to be more useful when used in conjunction annual, with tests on materials that have the same mode of action as the reference toxicant. However, frequent changing among reference toxicants can reduce the value of reference toxicant data if there is not an adequate history of use with each procedure, species and laboratory.

12. Experimental Design

12.1 Decisions concerning such aspects of experimental design as concentrations of test materials added to sediment, number of treatments, and numbers of test chambers per treatment should be based on the purpose of the test and the type of procedure to be used to calculate results (see Section 16). The amphipod sediment toxicity test can be used to test the toxicity of sediment in the field (see 12.3) or to address a great variety of sediment and water quality manipulations in the laboratory (see 12.4). Every test requires one or more control treatments (see 12.2). peak production occurring from late winter through spring (Kemp et al., 1985 (190)).

12.2 Controls—Every Species Selection—All four species have been routinely used to test requires sediments with a control treatment consisting range of sediment from the amphipod collection site grain size characteristics and pore-water salinities. Selection of one or more of the four specimens for a particular test/investigation should take into consideration the geographic location of the testing facility and within study area, the geochemical requirements pore-water salinity regime of the test species (see 10.3). The same water, conditions, procedures, study area, and organisms are used as in the other test treatments, except that none grain size characteristics of the test material sediment being tested. The species that is added used should exhibit tolerance to the control sediment or water. At least five laboratory replicates physicochemical properties of the control every sediment should be included in all tests regardless a particular study. Pore-water ammonia concentrations may also enter into selection of whether test sediments are replicated. This allows comparisons one species over others because the four species exhibit differential sensitivity to aqueous ammonia. Most often it will not be necessary to discriminate among experiments the four species, and among laboratories of the validity of procedures used in individual tests.

12.2.1 In addition decision to test one species above the standard control, if a field sediment has properties such as grain size rest may be driven by practical or logistical concerns. For example, a testing facility may choose to primarily test one species with a suitable local population in order to prevent potential complications associated with shipping. However, sediments may be encountered with characteristics that might exceed are outside of the tolerance range of one or more of the test species, it is desirable to include nontoxic reference sediment controls species. For example, grain size limitations for *A. abdita* and *R. abronius* are <10 % and >90 % fines, respectively. If these characteristics. The design of field surveys should include an additional field control involving five replicate samples from an area species are exposed to sediments that is free from sediment contamination. This provides a site-specific basis for comparison exhibit textural characteristics outside of potentially toxic and nontoxic conditions, and can account for these extremes, any mortality associated exclusively with subjecting the organisms that is observed could be due to nonnative sediments. The concentrations effects of chemical grain size independent of contaminants should be measured in these field control sediments in order to justify associated with the assumption that they are contaminant-free (see 10.3).

12.2.2 If any solvent other than water is present sediment. Ambiguity in interpretation may be avoided by careful consideration of the test chambers, a solvent control is also required. The solvent control must be identical to species given the regular control, except that the highest amount of solvent present in any other treatment is added sediment to this treatment. If the test material be tested. Comparative information is a mixture, formulation, or commercial product, none of available for the ingredients is considered a solvent unless an extra amount is used four species on sediment grain size sensitivity, salinity application ranges, and sensitivity to prepare the stock solution (see 10.6.5). aqueous ammonia (section A2.4).

12.3 Field Survey Design—Field surveys can be designed to provide either a qualitative reconnaissance of Collection:

12.3.1 Field collection is presently the distribution of sediment toxicity most common method for obtaining estuarine or a quantitative statistical comparison marine amphipods for sediment testing. All four species are commonly collected, shipped, and held in the laboratory; However, (USEPA-USACE 2001(2)) recommends establishing laboratory cultures of toxicity among stations:

12.3.1 The object of a qualitative reconnaissance survey is to identify sites of potential toxic conditions that warrant further study. It is often conducted in areas where little is known about contamination patterns. To allow *L. plumulosus* (section 12.5). Commercial vendors are available for maximum spatial coverage, the survey design might include only one sample from each

station, all four species. The lack availability of replication precludes statistical comparisons, but samples from sites where mortality exceeds the control range can appropriate size class for each species may vary seasonally. The collection site chosen should be identified one for further study.

12.3.2 The object which the presence of a quantitative statistical comparison is to test for statistically significant differences in effects among negative control or reference sediments abundant organisms of the correct size and test sediments from several sites. Replicates (that is, separate samples from different grabs taken at age has been demonstrated previously, and identification of the species has been confirmed taxonomically (for example, Bousfield, 1973 (171); Barnard and Barnard, 1982 (191)). Collection areas should be taken at each station in the survey. The number relatively free of replicates needed per station is contamination. All individuals in a function of the need for sensitivity or power (see 12.3.6). Separate subsamples test must be from the same grab source, because different populations may be used exhibit different sensitivities to test for within-grab variability, or split samples of composited sediment from one or more grabs may be used for comparisons of test procedures (such as comparative sensitivity among test species), but these subsamples should not be considered to be true replicates for statistical comparisons among stations contaminants. The four species are found in distinctly different habitats (Table 6).

12.3.2 *Species-specific Habitat Characteristics (29, 30)*

12.3.3 Station locations might be distributed among a known pollution gradient;

12.3.2.1 *Ampelisca abdita* is found mainly in relation to protected areas from the boundary low intertidal zone to depths of a disposal site, or at sites identified as being potentially toxic in a reconnaissance survey. Comparisons 60 m. This species generally inhabits sediments from fine sand to mud and silt without shell fragments, although it can also be made found in both space and time. In pre-dredging studies, relatively coarser sediments with a sampling design can be prepared to assess the toxicity sizeable fine component. This species is often abundant in sediments with a high organic content Aggregations of samples representative *A. abdita* are indicated by an abundance of tubes on the project area to sediment surface, location of which can be dredged. Such facilitated by looking through a design must include subsampling cores taken to the project depth.

12.3.4 If no amphipods survive in sediment from a particular field location, it might glass-bottom bucket. Although populations may be useful to conduct seasonally ephemeral, *A. abdita* is routinely collected year-round for toxicity tests with dilutions testing from subestuaries of Narragansett Bay, RI and from San Francisco Bay, CA.

12.3.2.2 *Eohaustorius estuarius* is found on protected and semi-protected beaches from mid-water level to shallow subtidal, within the field sediment mixed with control sediment. Concentrations should upper 10 cm (Environment Canada, 1992 (5)). *Eohaustorius estuarius* can be expressed as percent dilutions found on a wet weight basis, that is, wet weight open coasts in beds of field sediment/total wet weight of field and control sediment mixture. Experimental designs for sediment dilution experiments are freshwater streams flowing into the same as those described ocean, and in 12.4 for other laboratory experiments.

12.3.5 Sediment toxicity surveys are usually part sand banks in estuaries, above the level of more comprehensive analyses of biological, chemical, geological, other regional eohaustorids (*E. sawyeri* and hydrographic conditions. A useful summary of field sampling design is presented by Green E. *washingtonianus*; Environment Canada, 1992 (30). Statistical correlation can be increased and costs reduced if subsamples (5)). *Eohaustorius estuarius* inhabits clean, medium-fine sand with some organic content It is routinely collected for sediment toxicity tests, geochemical analyses, and benthic community structure are taken simultaneously tests from Yaquina Bay OR and Beaver Creek near Newport, OR, and on the same grab or at the same station.

12.3.6 The power west coast of the toxicity test Vancouver Island, BC, Canada.

12.3.2.3 *Leptocheirus plumulosus* is a found in subtidal portions of Atlantic Coast brackish e-estuaries. It is common in protected embayments, but has been collected in channels of replicates and the number estuarine rivers up to depths of individuals and variability 13 m. It is most often found in the response measure. On the basis fine-grained sediment with a high proportion

TABLE 6 Comparison of Habitat Characteristics and Other Life History Parameters of four Estuarine or marine Amphipod Species Used in Sediment Toxicity Tests (USEPA 1994a (1))

Criterion	<i>Ampelisca abdita</i>	<i>Eohaustorius estuarius</i>	<i>Leptocheirus plumulosus</i>	<i>Rhepoxynius abronius</i>
Substrate Relation	Tube dwelling, closed and well developed ^A	Free burrowing ^B	Tube dwelling, open and less developed ^A	Free burrowing ^C
Zoogeography	Atlantic-Gulf ^A San Francisco ^{D,E}	Pacific ^{B,F}	Atlantic ^A	Pacific ^C
Habitat	Poly-upper mesohaline ^A	Oligo-mesohaline ^{B,F}	Oligo-mesohaline ^A	Polyhaline ^{C,G}
Life cycle	40 to 80 days ^H	Annual ^B	30 to 40 days ^{I,J,K}	Annual ^L
Availability	Field or potential laboratory culture ^A	Field ^B	Field and laboratory culture ^{I,J,K}	Field ^C
Ecological importance	High	High	High ^I	High

^A Bousfield, 1973 (171)

^B DeWitt et al., 1989 (11)

^C Barnard and Barnard, 1982 (191)

^D Nichols et al., 1985 (176)

^E Hopkins, 1986 (177)

^F Environment Canada, 1992 (5)

^G Swartz et al., 1985 (10)

^H Scott and Redmond, 1989 (12)

^I DeWitt et al., 1992a (6)

^J Schlekot et al., 1992 (13)

^K McGee et al., 1993 (19)

^L Kemp et al., 1985 (190)

of historical control data particulate organic material, although it has been collected in fine silty sand with some organic content.

12.3.2.4 Primary habitats of *R. abronius* include nearshore subtidal zones on the Pacific Ocean coastline, and sub- and intertidal zones within polyhaline portions of estuaries in the Pacific Northwest. *Rhepoxynius abronius*, with five independent replicates per treatment and 20 amphipods per replicate, there is a 75 % probability of detecting a significant difference ($P < 0.05$) if the difference in mean survival between control and test sediment is 2.8 (see Table 2). For control survival of 18.0 (90 %), this corresponds to a test sediment mean survival of 15.2, about a 15 % reduction. Since the number of survivors in test sediments is often much less than 15, this is a reasonable level of precision naturally inhabits clean, fine sand. It has been collected for most applications:

12.4 Laboratory Experiments—Sediment use in toxicity tests from Lower Yaquina Bay, OR (Swartz et al. 1985 (10)), and West Beach, Whidbey Island, WA (Ramsdell et al. 1989 (195); Word et al. 1989 (16)).

12.3.3 Collection Methods:

12.3.3.1 Subtidal amphipods can be applied in the laboratory to provide information on collected with a variety of problems related to the action of contaminants in sediment. The test small dredge or grab (for example, PONAR, Smith-McIntyre, or Van Veen). Intertidal populations can be used to determine natural limits such as salinity, temperature, etc., to estimate the LC50 of collected using a contaminant in a particular sediment type, to study the interaction among contaminants in sediment, and to assess the effect of complex waste mixtures on the test species in sediment.

12.4.1 An acute test used to calculate an LC50 shovel or an EC50 usually consists of one or more control treatments and by skimming the sediment surface with a geometric series of at long-handled, fine-mesh net. At least five concentrations of test material. Except one-third more amphipods should be collected than are required for the control(s) test.

12.3.3.2 All apparatus used for collecting, sieving, and the highest concentration, each concentration transporting amphipods and control-site sediment should be at least 60 % clean and made of the next higher one, unless information concerning the concentration-effect curve indicates that a different dilution factor is more appropriate. At least one concentration non-toxic material. They should give a partial response below the LC50 or EC50 be marked “live only” and one above the LC50 or EC50. If the estimate of acute toxicity is particularly uncertain, six or more concentrations might should never be desirable to increase the likelihood of covering the appropriate range.

12.4.2 If it is only necessary to determine (a) whether a specific concentration is acutely toxic to the test species used for working with formalin or (b) whether the LC50 or EC50 is above or below a specific concentration (see 10.7.4), only that concentration any other toxic materials and should be stored separately from the controls are necessary. Two additional concentrations at about one-half aforementioned. The containers and two times the specific concentration of concern are desirable to increase confidence in the results.

12.4.3 An LC other collection apparatus should be cleaned and rinsed with deionized water, dechlorinated laboratory water, reconstituted seawater, or EC near natural seawater from the extremes of toxicity, such as an LC5 collection site or an LC95 uncontaminated source before use.

12.3.3.3 To minimize stress, amphipods should not be calculated unless at least one concentration of test material killed or affected a percentage of test organisms, other than 0 or 100 %, near the percentage for which the LC or EC is to handled carefully, gently, and quickly, and only when necessary. Amphipods can be calculated. This requirement might be met isolated from collection-site sediment using gentle sieving. *Ampelisca abdita* is exceptional in a test requiring vigorous sieving to determine an LC50 or EC50, but special tests with appropriate test concentrations and possibly more replicates per treatment might be necessary. Other ways of providing information concerning induce the extremes of toxicity are animals to report the highest concentration of test material that actually killed or affected no greater a percentage of the test organisms than did the control treatment(s), or leave their tubes. Once sieved, attempts should be made to keep amphipods submersed in collection site sea water at the lowest concentration of test material that actually killed or affected ambient collection temperature at all test organisms exposed to it. These alternatives times. Amphipods that are normally more reliable than reporting a calculated result such as reporting an LC5 dropped, or LC95 unless two or more concentrations resulted in percent killed or affected close to 5 or 95 %.

12.4.4 The primary focus of the physical and experimental design of the test and the statistical analysis of the data is the experimental unit, that is defined as the smallest physical entity to which treatments can injured should be independently assigned (see Guide E 729). Thus, discarded. Once separated from the test chamber is the experimental unit. With respect sediment, amphipods should not be exposed to factors that might affect results within test chambers and, therefore, the results direct sunlight.

12.3.3.4 The mesh size of the test, all chambers in sieve will depend on the species collected. Sieves with 0.5-mm mesh should be treated as similarly as possible. For example, the temperature in all test chambers used for sediment containing *A. abdita* and *L. plumulosus*. Larger *A. abdita*, which should not be as similar as possible unless used in the purpose test, should be excluded by sieving first with a 1.0-mm screen. When sieving *A. abdita*, only about half of the test is to study the effect of temperature. Test chambers are usually arranged in one or more rows. Treatments must amphipods will be randomly assigned to individual test chamber locations. A randomized block design (with each treatment being present in each block, that might extracted from their tubes. The tube mat should be a row or a rectangle) is preferable placed undisturbed for 20 to a completely randomized design 30 min to reduce coax the probability of chance segregation of treatments remaining animals out. Sieves with 1.0-mm mesh should be used for (27) *E. estuarius* and *R. abronius*.

12.4.5 The minimum desirable number

12.3.3.5 Collection-site water should be used to sieve sediment in the field. A 2-cm thick layer of sieved collection site sediment should be placed in transport containers, and organisms per treatment this sediment covered with collection-site water. Detritus and predators recovered by sieving should be calculated from (a) removed, and the expected variance within test chambers, (b) collected amphipods should be gently washed into the expected variance between test chambers within a treatment, transport containers with collection site water.

12.3.3.6 The salinity and temperature of surface and bottom sea water at the collection site should be measured and recorded. An adequate portion of collection site sediment should be returned with the amphipods to serve as both laboratory holding sediment or for use as control sediment in the toxicity test.

12.3.3.7 During transport to the laboratory, amphipods should be kept in sieved collection-site sediment at or below the collection site temperature. Containers of amphipods and sediment should be transported to the laboratory in coolers with icepacks, and the water in the containers of amphipods should be aerated if transport time exceeds 1 h.

12.3.3.8 An alternate collection method for either *A. abdita* involves transporting intact field-collected tubes to the maximum acceptable width laboratory for isolation of amphipods. This method is advantageous because separation of *A. abdita* from its tubes may be time-consuming when attempted in the confidence interval field, a practice which may be impractical in cold winter months. Amphipod tubes are collected and placed on a 0.5-mm sieve. The sieve should be shaken vigorously to remove most of the sediment, leaving the intact tubes. The tubes should be placed into a covered bucket that contains a sufficient quantity of collection site water to cover the collected material, and transported to the laboratory. In the laboratory, the tubes should be removed from the collection buckets and placed on a sieve series consisting of a 2-mm mesh sieve over a 0.5-mm mesh sieve. Amphipods should be forced from their tubes by spraying sea water on the material present on the 2-mm sieve. When all the tube material has been sprayed, the 0.5-mm sieve should be shaken vigorously to separate amphipods from any material that is present. The 0.5-mm sieve should then be completely submersed, at which point the amphipods will float on the water surface. The amphipods should then be skimmed from the surface with a small aquarium net and transported to a container with sea water at the appropriate temperature. The shaking process should be continued until only a few amphipods remain in the sieve.

12.3.3.9 *Leptocheirus plumulosus*—Although established cultures of *L. plumulosus* are the recommended source of organisms for new cultures, it is recognized that field collection of amphipods might be necessary to enhance genetic diversity of existing cultures or to establish new cultures at a laboratory (USEPA-USACE 2001 (2)). The taxonomy of the organisms should be confirmed before they are introduced into existing laboratory populations. New organisms should be carefully inspected, and all other species of amphipods should be removed. The ability of a wild population of sexually reproducing organisms to crossbreed with existing laboratory populations of *L. plumulosus* should be confirmed through long-term culture maintenance (Duan et al., 1997 (173)). Collection areas should be relatively free of contamination. Field collection of *L. plumulosus* neonates for immediate use in a chronic toxicity test is not recommended.

12.3.3.10 *L. plumulosus* is subtidal and can be collected with a small dredge or grab (for example, LC50 Ponar, Smith-McIntyre, or EC50 Van Veen). In very shallow water, sediment containing *L. plumulosus* can be collected with a shovel or scoop, or using a suction dredge (DeWitt et al., 1992a (6)). *L. plumulosus* can be isolated easily from collection-site sediment by gentle sieving. Ideally, amphipods will be separated into adults, subadult, and neonates. To reduce field processing time, 1.0-mm and 0.6-mm mesh sieves can be used to isolate adults and subadults with which to start a culture. Sediment passing through the minimum difference that 0.6-mm sieve could be temporarily used for holding until further processing of the sediment is practical. The final sieving of collection-site sediment through 0.25-mm mesh can be deferred until materials are returned to the laboratory.

12.3.4 Life Stage and Size:

12.3.4.1 The life stage for amphipods used in sediment toxicity tests will depend on the species tested. For *A. abdita* and *L. plumulosus*, sub-adult (immature) individuals should always be selected for testing. The life cycle of these species is relatively short, so the likelihood of senescence and any effects that could be associated with reproductive development or maturation are minimized if young individuals are selected. **(32) *Eohaustorius estuarius* and *R. abronius* are annual species with longer life spans than *A. abdita* and *L. plumulosus*. As Mature individuals can be used providing they are within the number recommended size range.**

12.3.4.2 The size range of test animals should be kept to a minimum regardless of the chosen species. For all species, mature female amphipods, which are distinguishable by the presence of embryos in the brood pouch or oviduct, should not be selected for testing. Additionally, mature male *A. abdita* and *L. plumulosus* should not be used. Recommended size ranges for the four species are as follows:

(1) *Ampelisca abdita*—3 to 5 mm; or those amphipods retained on a 0.71-mm sieve after passing through a 1.0-mm sieve. Adult male animals should not be tested; they are active swimmers and die shortly after mating.

(2) *Eohaustorius estuarius*—3 to 5 mm; or those amphipods retained on a 1.0-mm sieve. Large individuals (that is, >5 mm) should not be tested because they might be senescent.

(3) *Leptocheirus plumulosus*—For 10-d toxicity testing: 2 to 4 mm; or those amphipods retained on a 0.5-mm sieve after passing through a 0.71-mm sieve. See section 12.5 for methods to obtain *L. plumulosus* from cultures to start a 28-d sediment exposure.

(4) *Rhepoxynius abronius*—3 to 5 mm; or those amphipods retained on a 1.0-mm sieve. Large individuals (that is, >5 mm) should not be tested because they might be senescent.

12.3.5 Shipping Methods:

12.3.5.1 All four species have been routinely shipped from the collection site to the laboratory for sediment toxicity testing. Currently, shipping from the collection site is necessary for many testing laboratories because culture methods are not available for all four species. It is important that shipping methods ensure that consistently healthy animals are used in successive toxicity tests. Additionally, the amphipods that are received by a laboratory should meet the shipping acceptance criteria recommended for each species. Shipping methods and acceptance criteria will vary depending on the species used.

12.3.5.2 *Ampelisca abdita*—Collected amphipods should be shipped within 24 h of collection. Acceptable methods are available for shipping *A. abdita* in sediment and in water. For shipping in sediment, small plastic “sandwich” containers (about 500 mL) with sealable lids should be used. The containers are filled three-quarters full with a minimum depth of freedom-increases, 2 cm of sieved fine-grain collection-site sediment and therefore, then to the top with well-aerated seawater. No more than 200 amphipods should be added to each container. Amphipods should be allowed to burrow into the sediment and build tubes before the containers are sealed. Containers should be sealed with lids under water to eliminate any air pockets. For shipping in water-only, scalable plastic bags 60 (about 1 L) should be used. Amphipods in their tubes should be placed in bags and a sufficient amount of collection site water should be added to keep the confidence interval on tubes moist. The air in the bag should be replaced with pure oxygen before sealing, and then placed into a second bag. Bags should be placed in a container that has a layer of material (that is, styrofoam or newspaper) sufficiently thick to prevent excessive movement over a layer of ice-packs. The shipping container should be marked to prevent it from being inverted.

12.3.5.3 *Eohaustorius estuarius* and *Rhepoxynius abronius* —Shipping methods for these organisms are essentially the same. Small plastic “sandwich” containers (about 500 mL) with sealable lids should be used. The containers are filled three-quarters full with sieved collection site sediment (fine sand) and then with a 1-cm layer of collection site sea water. Not more than 100 amphipods should then be added and allowed to burrow. After the animals have burrowed, the overlying water should be poured off, but the sediment should be moist. The containers are then sealed and ready for shipment.

12.3.5.4 *Leptocheirus plumulosus*— *L. plumulosus* should be shipped in water only (USEPA-USACE 2001 (2)). Care should be taken to select containers with a significance test increases:

12.4.6 Mean survival firm seal that is not easily broken in-control sediment must shipment. The containers are filled to the top with well-aerated water. No more than 100 amphipods/L should be 90% added to each container. For shipping, sealable plastic bags, cubitainers, and other sealable plastic containers can be used. The containers should be filled with well-aerated collection-site water or greater. A difference of about 15% between mean survival culture water before they are sealed. The double packing bags should be placed in a container that has a protective layer of material (that is, Styrofoam or newspaper) sufficiently thick to prevent excessive movement with an under layer of ice packs. The shipping container should be marked to prevent it from being inverted.

12.3.6 Performance Criteria for Shipped Amphipods:

12.3.6.1 The process of ensuring the availability of healthy amphipods on the day that the test-sediments is usually significant set up begins when the animals arrive in the laboratory from the supplier. Although the ultimate performance criterion for amphipods are included utilized in sediment toxicity tests is achievement of five replicate test chambers of >90% survival in control and test sediment (see 16.5):

12.4.7 It is (Table A1.3 in Annex A1), it would be desirable to repeat assess the test at a later time to obtain information concerning the reproducibility quality and acceptability each batch of shipped amphipods using the results:

13. Procedure

13.1 *Dissolved Oxygen*—The concentration criteria that follow. For all four species, biological criteria should include an exhibition of dissolved oxygen (DO) active swimming behavior upon placement in water, full digestive tracts, and an acceptable color. *Ampelisca abdita* should be opalescent pink, *E. estuarius* should be grayish- or yellowish white, *L. plumulosus* should be brown or orange-gray, and *R. abronius* should be salmon pink, grayish- or yellowish-brown, or white with a pinkish-brown hue. Mortality among the shipped animals should not exceed 5%. No sexually mature animals should be included in shipments of *A. abdita* or *L. plyumulosus*. The shipping containers should arrive intact, and the temperature of water or sediment in the test chambers shipping containers should be between 4 and 10°C. Information on physical parameters of the collection site, including at or near saturation least temperature and salinity, should be provided by gently aerating the water (see annexes). Air supplier. Finally, a quantity of collection site sediment should be bubbled into included as substratum for amphipods during the acclimation period or for use as control sediment in the test. It may be desirable for the testing facility to stipulate these criteria to the supplier when the animals are ordered. If these criteria are not met, the animals may have experienced stress during shipment, and >90% survival in control sediment may not be achieved.

12.4 Holding and Acclimation:

12.4.1 *Density*—Amphipods should be held and acclimated (if necessary) in containers that contain a 2- to 4-cm layer of collection site sediment that maintains has been sieved through a ≥90% dissolved oxygen concentration, but does 0.5-mm mesh screen. Amphipod density should not cause turbulence or disturb exceed 1 amphipod/cm². *Ampelisca abdita* that have been shipped in their tube material in bags can be held under those conditions as long as aeration and food are supplied and the temperature and salinity holding and acclimation procedures are followed.

12.4.2 *Duration*—Depending on temperature and salinity at the collection site, amphipods may have to be acclimated to test conditions. If air flow necessary, changes in temperature or salinity to bring amphipods from the collection site conditions to the

test conditions should be made gradually (for example, ~~s~~ should not exceed 3°C and 5 ‰ per 24 h). Once test conditions are achieved, amphipods should be maintained at these conditions for at least two days before testing to allow for acclimation. Amphipods held for more than ten days should not be used for testing because they may not satisfy performance control criteria. Temperature and salinity should be measured at least daily during the period when amphipods are being adjusted to the conditions of the test water. Thereafter, temperature, salinity, pH, and DO should be measured in the beakers to determine whether dissolved oxygen concentrations have dropped to less than 60 % holding containers at least at the start and end of saturation (see 15.2.7).

~~13.2~~ the acclimation period, and preferably daily.

~~12.4.3 Temperature~~—The Overlying water temperature selected should be within the natural range of temperatures in the area from which the amphipods occur in the field. Within an experiment, individual temperature readings should not vary by more than 3°C from per day during acclimation to the selected test temperature, a. Once the time-weighted average measured temperature at the end of the test temperature is reached, amphipods should be within 1°C maintained at that temperature for a minimum of the selected test temperature. When 2 d. A water bath, an incubator, or temperature-regulated room can be used for temperature is measured concurrently in more than one test chamber, the highest and lowest temperatures should not differ by more than 2°C.

~~13.3~~ acclimation.

~~12.4.4 Salinity~~—The water overlying—It is unlikely that either *A. abdita* or *R. abronius* will require salinity acclimation because the test sediment in sediment toxicity tests must collection site salinity for these two species will likely be within 3 ‰ of the tolerance range test salinity of 28 ‰. Salinity of water used for temperature acclimation for these species, if necessary, should be the selected test species (see annexes): salinity, or 28 ‰. The target test salinity of for *E. estuarius* and *L. plumulosus* is 20 ‰, and it is likely that the collection site salinity will be considerably lower than this for both species. Upon arrival in the laboratory, the water of test sediments from used to hold *E. estuarius* and *L. plumulosus* should be adjusted to 20 ‰ by adjusting the salinity in the holding container at a rate that should not exceed 5 ‰ per 24 h. The amphipods should be adjusted, because such an operation might change the toxicological properties maintained at 20 ‰ for 2 d before testing. A salinity of 5 or 20 ‰ is recommended for routine application of 28-d test with *L. plumulosus* (Annex A2; USEPA-USACE 2001 (2)) and a salinity of 20 ‰ is recommended for routine application of the sediment. The 10-d test with *E. estuarius* or *L. plumulosus* (Annex A1). However, the salinity of the interstitial overlying water for tests with these two species can be adjusted to a specific salinity of sediments experimentally spiked in interest (for example, salinity representative of site of interest or the laboratory with contaminants objective of the study may be adjusted prior to spiking.

~~13.3.1~~ If test sediments are collected from low-salinity areas, evaluate the influence of salinity on the bioavailability of chemicals sediment). Importantly, the overlying water in salinity tested must be within the tolerance range of the test organisms (as outlined in Annex A1 and Annex A2). If tests are conducted with procedures different from those described section 1.3 or in Table A1.1 (for example, different salinity, lighting, temperature, feeding conditions), additional tests are required to determine comparability of results (section 1.10).

~~12.4.5 Lighting~~—Lighting should be approximately constant and continuous throughout the holding and acclimation period for 10-d tests, all species except *A. abdita* require a 16:8 L:D photoperiod to promote feeding. Fluorescent lights should be used, and they should provide from 500 to 1000 lux at the surface of the sediment in holding containers.

~~12.4.6 Water:~~

~~12.4.6.1~~ Provided that it is acceptable to the test organisms, either an uncontaminated supply of natural sea water or reconstituted sea water can be used for holding and acclimation (section 7.1). At a minimum, healthy amphipods should exhibit acceptable survival in holding water, and should not exhibit signs of stress, such as the unusual behavior or changes in appearance.

~~12.4.6.2~~ If natural sea water above the sediment at the collection site. Depending upon experimental design, is used, it might should be desirable to use water obtained from an uncontaminated area known to support a healthy, reproducing population of the sediment collection site, test species or comparatively sensitive species. Reconstituted sea water is prepared by adding commercially available sea salts to adjust water from a suitable source, in quantities sufficient to provide the desired salinity. Pre-formulated brine (for example, 60 to 90 %) prepared with dry ocean salts or heat-concentrated natural sea water can also be used. To obtain the desired holding or acclimation salinity, sea salts or brine can be added to a suitable fresh water, natural estuarine water, or the collection site salinity (see 7.3).

~~13.4 Light~~—For sediment toxicity laboratory's sea water supply. The suitability and consistency of a particular salt formulation for use in holding and acclimation should be verified by laboratory tests involving because some formulations can produce unwanted toxic effects or sequester contaminants (Environment Canada, 1992 (5)). Reconstituted water should be intensively aerated for two weeks before use (Environment Canada, 1992 (5) ; section 7.1). Suitable sources of water used for preparing reconstituted sea water include deionized water or an uncontaminated natural surface water or ground water. Chlorinated water must never be used because residual chlorine and chlorine-produced oxidants are usually left on continuously. The constant light increases highly toxic to many aquatic animals. Dechlorinated municipal drinking water should be used only as a last resort because dechlorination is often incomplete.

~~12.4.6.3~~ Assessments of the tendency quality of the water used for holding and acclimation and for preparing reconstituted sea water should be performed as frequently as required to re document acceptability (section 7.1). Analyses of variables including salinity, temperature, suspended solids, pH, DO, total dissolved gasses, ammonia, nitrite, pesticides, and metals are

recommended. Sea water used for holding and acclimating amphipods should be filtered (<5 µm) shortly before use to remain exposed to the test material. For other species a different photoperiod might remove suspended particles and organisms and should be desired (see annexes):

13.5- aerated for a two weeks before use (section 7.1).

12.4.6.4 *Feeding*—Infaunal amphipods do not—*Ampelisca abdita* and *L. plumulosus* require supplementary feeding during the 10-day holding or acclimation for 10-d toxicity test. Supplemental feeding might tests conducted with field-collected organisms. *Ampelisca abdita* should have food available daily, whereas *L. plumulosus* should be fed every other day. *Ampelisca abdita* can be supplied with an algal ration consisting of *Pseudoisochrysis paradoxa* or *Phaeodactylum tricomutum* that is provided in conjunction with sea water renewal. Some laboratories have reported success in providing *A. abdita* enriched dried algal material (Docosa Gold and Golden Shell algal-based natural feed supplements (Sanders Brine Shrimp Company, Ogden, Utah)) slurried in seawater (Michele Redmond, Northwestern Aquatic Sciences, Newport, OR; personal communication). Other diets can be used provided the diet has been demonstrated to support acceptable organisms for longer tests (see testing (Table A2.3 in Annex A3).

13.6—*Beginning A2*). See Stein (1973 (196)) for procedures to culture algae. After 75 % of the overlying water has been removed, each holding container should be renewed with sea water at the appropriate salinity that contains algae at a concentration of at least 1×10^6 cells/mL. *Leptocheirus plumulosus* should also be provided with dry food ration, consisting at a minimum of finely powdered Tetramarine® (formerly called TetraMin®). It may be desirable to grind the dry food in a blender. Each container should receive about 0.4 g dry food/350 amphipods. *Eohaustorius estuarius* and *R. abronius* will utilize organic material in the holding sediment as food and do not require supplemental feeding.

12.4.6.5 *Acceptability of Animals*—Amphipods counted into the holding or acclimation chambers should be active and appear healthy. Any individuals that fail to burrow or fail to make tubes (that is, *A. abdita*) in holding sediment or that appear unhealthy during the holding or acclimation period should be discarded. Apparently dead individuals should also be discarded. If greater than 10 % of the amphipods emerge or appear unhealthy during the 48 h preceding the test, the entire group should be discarded and not used in the test. Additionally, the group should be discarded if more than 10 % of the amphipods die or become inactive during the holding period before testing.

12.5 *Culture Procedure for Leptocheirus plumulosus*:

13.6.1 The toxicity

12.5.1 *General Culturing Procedures* :

12.5.1.1 Acceptability of a culturing procedure is based in part on performance of organisms in culture and in the sediment test (Table A2.3 in Annex A2). No s-wingle technique for culturing test organisms is required. What may work well for one laboratory may not work as well for another laboratory. Although a variety of culturing procedures are first placed outlined below for *L. plumulosus*, organisms must meet the test acceptability requirements listed in Table A2.3 of Annex A2.

12.5.1.2 All organisms in a test chambers containing test material:

13.6.2—On must be from the-d same source. Organisms may be obtained from laboratory cultures or from commercial or government sources; a partial list sources is provided in Table 5. The test begins, each test sediment sample organism used should be thoroughly homogenized within its storage container, and identified using an aliquot added to a test chamber to a depth specific for appropriate taxonomic key, and verification should be documented.

12.5.1.3 Obtaining organisms from wild populations should be avoided unless organisms are cultured through several generations in the test species (see annexes): laboratory before use in testing. In addition, the ease ability of replicate sediment samples, it might be desirable to calculate the net weight wild population of sexually reproducing organisms to crossbreed with the existing laboratory population should be determined (Duan et al., 1997 (173)).

12.5.1.4 Test organisms obtained from commercial sources should be shipped in well-oxygenated water without sediment necessary in insulated containers to make a layer maintain temperature during shipment. Temperature, salinity and DO of the desired depth water in the first chamber, shipping containers should be measured at the time of shipment and then add on arrival to determine if the-s organisms might have been subject-wed to low DO, salinity change, or temperature and salinity fluctuations. The temperature and salinity of sediment the shipped water should be gradually adjusted to the desired culture temperature and salinity at rates noth exceeding 3°C or 3 ‰ per 24 h.

12.5.2 The culturing method below is based on procedures described in DeWitt et al. (1997a (7)). A periodic-renewal culture s-wystem is used. It consists of culture bins that contain aerated water over a-treatment. The same procedure might thin (about 1 cm) layer of clean, fine-grained sediment in which the amphipods burrow. Culturing areas must be applied separate from testing areas to control sediments, measuring avoid exposing the required weight cultures to contaminants. Before *L. plumulosus* are received at a testing facility, appropriate permits or approvals for replicates import of each treatment separately, because different sediments might have different densities. Treatments live organisms should be randomly assigned to prenumbered test chambers. It obtained, if necessary. If culturing is desirable to take subsamples of the test sediment for geochemical analyses as the test chambers occur in an area where *L. plumulosus* are loaded. For some experimental designs it might not indigenous to local waters, precautions should be desirable taken to test intact cores.

13.6.3 The sediment within prevent release of living organisms to the test chamber outside environment (section 10.5.15). Test animals should be settled by tapping destroyed at the end of toxicity test-e.

12.5.3 *Starting a Culture*—Amphipods for starting a laboratory culture of *L. plumulosus* should be obtained from aga source

with an established culture in which the species has been verified (see Table 5 for commercial sources of *L. plumulosus*). Alternatively, *L. plumulosus* can be obtained from field populations (section 12.3). Upon receipt of amphipods, the temperature and salinity of the water in shipping container(s) should be gradually adjusted to 20°C and desired culture salinity, at rates not exceeding 3°C or by smoothing 3‰ per 24-h period. Feeding and regular maintenance should begin once the acclimation period is complete. Separate organisms into three size classes by gentle sieving: adults (retained on 1.0-mm mesh), subadults (pass through 1.0-mm mesh and retained on 0.6-mm mesh), and neonates (pass through 0.6-mm mesh and retained on 0.25-mm mesh). Seed each culture bin with about equal numbers of adults, subadults, and neonates to achieve a nylon, fluorocarbon, or polyethylene spatula. A disk cut from 6 mil nylon, TFE-fluorocarbon, or polyethylene sheeting population density between 0.25/cm² to 0.35/cm² (2500 to 3500/m²). Select only actively moving, healthy-looking organisms. Cultures should not be stocked at densities greater than 0.5/cm² (5000/m²). See section 12.5.13.3 for guidance on maintaining culture densities. Field-collected organisms should be added periodically to the inside diameter culture population to maintain genetic diversity of the cultured amphipods.

12.5.4 Culture Bins—Culture bins should be easy to maintain. Plastic wash tubs (about 35 by 30 by 15 cm) have been used successfully by several laboratories (DeWitt et al., 1992a (6)). They are relatively light when filled with water and attached sediment, broad enough to a length allow for easy viewing of nylon monofilament amphipod burrows, easily cleaned, inexpensive, and readily available. A wide variety of containers and materials may work just as well for removal, can culturing this species. New plasticware should be placed on soaked in running water for several days before use in the sediment surface cultures to minimize sediment disruption as prepared toxicity test sea leach out potentially toxic compounds. Previously used culture bins usually can be satisfactorily cleaned using hot water is added up to and a scrub brush or pad, without the desired level use of a chemical cleanser. Culture bins should not be washed with soap or detergent except in the test chambers (see annexes). The disk extreme conditions. If such a cleaning is deemed necessary, culture bins should be removed and rinsed and soaked thoroughly after cleaning to remove residual cleanser.

12.5.5 Culture Sediment—Cultures should be established with sea water between replicates a thin layer (1 to 1.5 cm) of sediment spread on the bottom of a treatment, and a separate disk should culture bin. Sediment used for culture purposes can be the same as the control sediment used in sediment toxicity tests. Suitable sources for culture sediment include the amphipod collection site or an area adjacent to salt marsh vegetation. Culture sediment should be uncontaminated, organic-rich, fine-grained marine or estuarine sediment that is not anoxic. The test chambers organic carbon content (% TOC) should then be covered, put in numerical order into a temperature-controlled water bath, range between 1.5 % and aerated overnight. 4 %. The system sediment should be left overnight press-sieved through a 0.25-mm screen before use to allow suspended particles to settle facilitate the harvesting of neonates and an equilibrium to removal of indigenous macroinvertebrates. Culture sediment can also be established between wet sieved. Wet-sieving involves agitating or swirling the sieve containing sediment and overlying in water before so that particles smaller than the amphipods selected mesh size are added.

13.6.4 If washed through the experimental design requires monitoring sieve into a container. The sieve may be placed on a mechanical shaker, or the sediments on the screen can be stirred with a nylon brush to facilitate the process. Alternatively, the particles may be washed through the sieve with a small volume of running water. Culture sediment chemistry (for example, metals, total volatile solids, pH, Eh, etc.); can also be frozen (>48 h) to provide additional test chambers with assurance that viable macroinvertebrates are not present. Frozen sediment and amphipods should be homogenized after thawing and before use. Culture sediment can be stored frozen for this purpose. Monitoring the about 1 year.

12.5.6 Culture Water—Culture water used for holding and acclimating test organisms and for conducting toxicity tests should be of uniform quality and from the same source. See section 7.1 for acceptable sources of water. Cultures of *L. plumulosus* are usually maintained at a salinity of either 5‰ or 20‰. Culture salinity will depend on the anticipated pore-water salinity of test sediment and desired overlying water (for DO, pH, or for certain chemicals) salinity to be used in the test chambers (Table A2.1 in Annex A2). Alternatively, the salinity of the overlying water can be accomplished without disturbing adjusted to a selected target salinity (for example, one representative of the salinity regime at the site of interest; section 1.4). To obtain these salinity values, natural or reconstituted seawater should be diluted with nonchlorinated well water, deionized water, or reverse-osmosis water. Seawater and may dilution water should be done in the test chambers containing the test amphipods. Temperature can filtered (<5 µm). Water that might be measured contaminated with pathogens should be treated shortly before use by filtration (<0.45 µm), either alone or in a simulated test beaker containing water combination with ultraviolet sterilization. DO, salinity, and control sediment but no amphipods. If more than one pH should be checked before the water bath is used to contain the test chambers, in cultures. Batches of salinity-adjusted culture water can be held for about 1 week; a separate lower holding temperature (<20°C) helps maintain acceptable water quality. Water depth in culture bins should be included in each at least 10 cm. Aeration, provided through an air stone or pipette, should be moderate and constant, but not so vigorous as to resuspend sediment. Overlying water bath (see 13.9.3).

13.6.5 The toxicity test should be replaced the day after a new culture is initiated (Day 0) when amphipods are distributed established; thereafter, it should be renewed two or three times per week.

12.5.7 Temperature and Photoperiod—Cultures should be maintained at 20 to 25°C. The reproductive rate of *L. plumulosus* increases at temperatures greater than 20°C, necessitating more frequent culture thinning. Higher temperatures also can promote unwanted growth of nuisance organisms (such as nematodes, small worms, copepods). Temperatures below 20°C may not

to foster sufficiently prolific reproductive rates. Fluorescent lights should be on a 16 h light : 8 h dark photoperiod at a light intensity of 500 to 1000 lux. An efficient procedure is to allow test chambers maintain long-term cultures at 20°C, and increase culture temperature to about 25°C a few weeks in advance of testing.

12.5.8 Food and Feeding—This method recommends the simplest effective diet for routine use for *L. plumulosus* culture by providing finely milled Tetramarine® two or three times per week. Tetramarine® is a dry fish food (flake or powder) widely available in retail pet stores. The food is prepared by milling, grinding, or chopping the flakes to select a fine powder. A small flour mill, blender, or coffee grinder is useful for this. Ground powder is then sifted through a 0.25-mm mesh screen, retaining and using only the material that passes through the sieve. Use of test chambers (usually 10 a respirator or fume hood will minimize aspiration of dust. When establishing a new culture bin, do not add food for 3 to 15) to be processed together. If treatments 4 days after amphipods are replicated, each treatment, including controls, should be represented placed in each set of test chambers new sediment. This will encourage the organisms to be processed together. If treatments are not replicated, selection consume labile organic matter in sediment and to turn over the sediment by burrowing and feeding.

12.5.8.1 Culture bins should be random.

12.5.8.2 A sufficient number provided with food in conjunction with water renewal. Two or three times a week, about 60 % of amphipods culture water should be removed from each culture bin (by decanting, siphoning, or pumping) and replaced with the holding facility at one time to provide same volume of renewal water. Each culture bin is provided with about one third more amphipods than are needed for one set 0.4 g of test chambers. This allows selection dry food sprinkled evenly over the water surface, or as a slurry in culture water two or three times per week (for example, Monday-Wednesday-Friday or Monday-Thursday). The amount of dry food added will depend on the density of each culture bin. Newly started culture bins should receive slightly less food (for example, 0.3 g) than bins containing mature cultures. Excess food can decompose encouraging microbial and fungal growth on the sediment surface deteriorating water quality.

12.5.8.2 Some laboratories have experienced success in culturing *L. plumulosus* when other food is provided (that is, live microalgae or a mixed dried food; DeWitt et al., 1992a (6)). Modifications to the diet can be used by laboratories in order to optimize culture practices as long as performance criteria are satisfied (Table A2.3 in Annex A2).

12.5.8.3 One feeding alternative is to supply renewal water consisting of seawater, cultured phytoplankton, and deionized water combined to the temperature proper salinity and adjusted to an algal density of about 10^6 cells/mL (DeWitt et al. 1992a (6)). Proportions will vary depending upon the salinity of the seawater and the density of the cultured phytoplankton. Live algae also can be used periodically to supplement a routine supply of dry food. The algae used can include a single or multiple species (for example, *Pseudoisochrysis paradoxa*, *Phaeodactylum tricornutum*, *Isochrysis galbana*, *Chaetoceros calcitrans*, *Skeletonema sp.*, *Dunaliella tertiolecta*, and/or *Thalassiosira spp.*). Other algal species might be used if it can be demonstrated that they foster amphipod growth and reproductive rates equal to those of the aforementioned food alternatives. A mixture of algal species is recommended, if live algae is included in the holding containers diet.

12.5.9 Culture Maintenance—Cultures should be observed daily to ensure that temperature is acceptable and aeration is adequate in all culture bins. An inspection for the presence of oligochaetes, polychaetes, copepods, infaunal sea anemones, or chironomids should be conducted weekly. The presence of excessive densities of these or other competing or predatory organisms should prompt renewal of culture sediment after separating *L. plumulosus* from the holding sediment invasive organisms. During routine maintenance, cultures should be inspected for the presence of microbial and transferred to fungal build-up on the sediment surface. This build-up appears as a sorting tray containing water white or gray growth that may originate near uneaten food. Presence of microbial build-up may indicate that more food is being provided than is required by the amphipods. No additional food should be provided to culture bins with surficial microbial build-up until the build-up is no longer present. Sieving of sediment and renewal of culture bins can expedite removal of decaying material.

12.5.9.1 Healthy cultures are characterized by an abundance of burrow openings on the sediment surface and returned turbid water from amphipod activity. Although amphipods may leave their burrows to the holding containers search for use as reburial sediment at food or mates, they will ordinarily remain in their burrows during the termination illuminated portion of the photoperiod. Amphipod density should therefore only be estimated by examining the number of burrow openings. An abundance of organisms on the sediment surface (for example, >15 per culture bin) could indicate inadequate sediment quality, low DO concentrations, or overcrowding. A culture bin with an abundance of amphipods or unhealthy individuals on the sediment surface should be impartially selected from examined closely, and the DO concentration should be measured in the overlying water. If the DO concentration is below 60 % saturation (<4.4 mg/L), the culture bin should be sieved, and sequentially distributed among dishes containing approximately 150 mL the population and culture sediment examined. If the population is too dense (that is, >1.5/cm²), the culture should be thinned. If the sediment becomes an unacceptable habitat because it is anaerobic or black and sulphidic below the sediment surface, or contains an excess of competitive or predatory organisms, the healthy surviving amphipods should be placed in a new culture bin with newly prepared culture sediment.

12.5.9.2 Water temperature and DO should be measured in culture bins on a regular basis, about every week. Cultures should be continuously aerated. Salinity should be measured after water until renewal. Ammonia and pH in overlying water should be measured with each new batch of sediment before organisms are added.

12.5.9.3 Renewal of Cultures—*L. plumulosus* can be prolific, and care should be taken to ensure that culture bins do not get overcrowded. Amphipods in overcrowded culture bins can be stressed because of food and space limitations, causing the required

number (usually 20; see annexes) fecundity of individuals. The number females to drop below five eggs/brood/female and the abundance of neonates and subadults to decline dramatically. Culture density should not exceed 1.5 amphipods/cm² and should ideally be maintained at about 0.5 amphipods/cm². A typical indication of overcrowding is a fairly uniform size distribution of amphipods (mostly small adults) and the presence of only two to four eggs in the brood pouches of gravid females. If sediment is not replaced occasionally, the cultures may become infested with undesirable species, such as worms or copepods. These “pests” may compete for food, bind sediment as fecal pellets, or produce mucus, thereby reducing culture productivity or increasing the effort required to harvest amphipods. Field-collected organisms should be added to the culture population periodically (about annually) in order to maintain genetic diversity of the culture organisms.

12.5.9.4 To avoid overcrowding, cultures should be thinned every 6 to 8 weeks by recounting them into sieving through a separate dish containing toxicity test water.

13.6.7 Amphipods 0.25-mm mesh screen to remove sediment. Sediment can be used for a total of 2 to 4 months before it should be replaced. Discard old sediment, prepare new culture bins and sediment, and restock each bin.

12.5.10 Obtaining *Leptocheirus plumulosus* for a 10- and 28-d Test:

12.5.10.1 The cultures usually can be harvested about 4 to test chambers without disruption 5 weeks after initiation or up until the cultures are thinned and renewed (6 to 8 weeks after initiation). Neonates used for testing may be selected on the basis of size or age. For size-selected neonates, the contents of cultured bins are gently sieved through 0.60-mm and 0.25-mm screens. Neonates used for testing in 28-d tests may be selected on the basis of size or age. For size selected organisms, animals passing through a 6-mil nylon, TFE-fluorocarbon, 0.6 or 0.5 mm sieve onto a 0.25 mm mesh are used for testing and individually neonates typically have a dry weight of about 0.03 mg to 0.06 mg and body length of about 1.3 mm to 1.7 mm. In contrast, *L. plumulosus* used in the 10-d test are those that pass through a 0.71 mm sieve on to a 0.5 mm sieve. Culture bins of about 35 by 30 cm typically produce at least 300 to 400 neonates with a healthy culture. Selecting neonates for testing based on size is the preferred option for method comparability. For age-selected neonates, gravid females are isolated from cultures 5 d before test initiation. Gravid females are placed in separate culture bins with sediment and are fed. Two days before test initiation, these females are then transferred to bins containing only water surface, (for example, at 25°C and 5 ‰ or 20 ‰). On the day of test initiation, the contents of these bins are gently passed through a 1-mm screen on which adults are retained. Neonates that pass through this screen are transferred to a shallow glass container for sorting. Special care should be taken to ensure that the neonates are handled gently, selecting and transferring them with wide-bore pipets only, and maintaining the water temperature and salinity within recommended test conditions.

12.5.10.2 About one-third more amphipods than are needed for the test should be sieved from the sediment and transferred to a sorting dish over tray. The additional organisms allow for the selection of healthy, active individuals. Organisms not used in toxicity tests can be used to establish new cultures.

12.5.10.3 Amphipods placed in the test chamber. Any holding bins should be active and healthy. Sluggish or apparently dead individuals should be discarded. If greater than 10 % of the amphipods remaining in the holding bins appear unhealthy or are dead during 48 h preceding the test, the entire group should be gently washed into the discarded and not used in tests.

12.5.11 Minimization of Risk of Release of Nonindigenous Organisms:

12.5.11.1 If test chamber. The amphipods are not endemic to the local estuarine environment, containment and water-level treatment procedures should be brought up implemented to minimize the chance of accidental release of organisms or pathogens to local waters. The same precautions might also be required if the culture population of *L. plumulosus* is not derived from local sources. Some local or state authorities might require special permits and procedures to allow receipt and culturing of nonindigenous species. All test chamber, animals should be destroyed at the disk removed, end of toxicity tests. Culturing and holding of the chamber replaced amphipods should only occur in specially designated laboratory areas, separate from those used to hold, culture, or experiment with native species. These areas should have no access to drains leading directly to local surface waters. Handling of nonindigenous species should be limited to trained and authorized personnel. The amphipods should be cultured in a static-renewal manner to minimize the amount of water bath, covered, that needs to be treated. Any seawater removed from culture bins should be treated with chlorine bleach or ozonation to kill any escaping organisms and pathogens. All equipment used to culture or handle the amphipods should be cleaned thoroughly. Any excess or dead amphipods that do not bury within the time specified for the species (see annexes) should be removed placed in bleach or treated by ozonation or heat killed (boiling water) to ensure they are killed before disposal as sanitary waste.

13. Calculation

13.1 Data Recording:

13.1.1 Quality assurance project plans with data quality objectives and standard operating procedures should be developed before starting a test. Procedures should be developed by each laboratory to verify and archive data (USEPA 1994e, (160)).

13.7.1.2 DA file should be maintained for each sediment test or group of Test—The tests on closely related samples (Section 11). This file should contain a record of the sample chain-of-custody; a copy of the sample log sheet; the original bench sheets for the test organism responses during the sediment test(s); chemical analysis data on the sample(s); control data sheets for reference toxicants; detailed records of the test organisms used in the test(s), such as species, source, age, date of receipt, and other pertinent information relating to their history and health; information on the calibration of equipment and instruments; test chambers containing test sediment. Amphipods conditions used; and results of reference-toxicant tests. Original data sheets should

be signed and dated by the laboratory personnel performing the tests. A record of the electronic files of data should also be included in the file.

13.2 Data Analysis:

13.2.1 Statistical methods are used to make inferences about populations, based on samples from those populations. In most sediment tests, test organisms are exposed to chemicals in sediment to estimate the response of the population of laboratory organisms. The organism response to these sediments is usually compared with the response to a control or reference sediment. In any sediment test, summary statistics such as means and standard errors for ten days. There are no observed substantial effects response variables (for example, survival, chemical concentrations in tissue) should be provided for each treatment (for example, pore-water concentration, sediment concentration). See Section 14 of Test Method E 1706 and Guide E 1847 provide specific guidance on statistical analyses of data from sediment tests. Specifically, Test Method E 1706 provides guidance on the following: (1) experimental design (including replication, minimum detectable differences, randomization, pseudoreplication, compositing of samples) and (2) Statistical analysis of data (including hypothesis testing (for example, Analysis of variance) and regression analysis (for example, Effect concentrations (EC_x) and Inhibition concentrations (IC_x)).

13.2.2 Types of Data—Two types of data can be obtained from sediment tests. The most common endpoint in toxicity testing is mortality, which is a dichotomous or categorical type of data. Other endpoints might include growth and reproduction. These types of endpoints are representative of continuous data.

13.2.3 Sediment Testing Scenarios—Sediment tests are conducted to determine whether contaminants in sediment are harmful to benthic organisms. Sediment tests are commonly used in studies designed to: (1) evaluate hazards of time dredged material, (2) assess site contamination in the environment (for example, to rank areas for cleanup), and (3) An exposure period determine effects of less than ten days is not generally recommended. Experiments with cadmium and field sediments have shown that many amphipods emerge from specific contaminants, or combinations of contaminants, through the use of sediment spiking techniques. Each of these broad study designs has specific statistical design and analytical considerations, which are alive but unable described as follows.

13.2.3.1 Dredged Material Hazard Evaluation— In these studies, n (number) sites are compared individually to re-bury after four days, but most of a reference sediment. The statistical procedures appropriate for these amphipods studies are dead after ten days generally pairwise comparisons. Additional information on toxicity testing of exposure dredged material and analysis of data from dredged material hazard evaluations is available in (3114, 115, 116). For some experimental designs, such as comparison

13.2.3.2 Site Assessment of a 96-h LC50 between species Field Contamination—Surveys of sediment toxicity are often included in the presence or absence more comprehensive analyses of sediment, other exposure periods biological, chemical, geological, and hydrographic data. Statistical correlation can be improved and costs may be used:

13.8 Biological Data—Response criteria indicating toxicity of test reduced if subsamples are taken simultaneously for sediment include mortality tests, chemical analyses, and sublethal effects. Sublethal effects include (a) emergence from highly toxic sediment during benthic community structure determinations. There are several statistical approaches to field assessments, each with a specific purpose. If the course of the test, and (b) inability of surviving but affected amphipods objective is to re-bury in clean, collection site sediment at compare the termination of the test. Response criteria must be monitored in response or residue level at all sites individually to a “blind” fashion, that is, control sediment, then the observed must have no knowledge of pairwise comparison approach described as follows is appropriate. If the treatment of the sediment in the test chambers. This objective is accomplished through randomization of sample numbers:

13.8.1 Emergence—Since most infaunal amphipods remain buried during sediment toxicity tests, there is little opportunity to monitor temporal changes compare among all sites in mortality or sublethal effects. An exception the study area, then a multiple comparison procedure that employs an experiment-wise error rate is appropriate. If the temporal pattern objective is to compare among groups of emergence from highly toxic sediment. The test should be monitored at least daily (including the day sites, then orthogonal contrasts are a useful data analysis technique.

13.2.3.3 Sediment Spiking Experiments—Sediments spiked with known concentrations of initiation and the day of termination) for temperature, aeration, lights, and emergence of the amphipods from the test sediment. Each test chamber should chemicals can be observed by temporarily turning off the air used to the test chambers, establish cause and gently removing the cover from individual chambers with minimal disturbance effect relationships between chemicals and biological responses. Results of the chamber. The number of amphipods observed completely or partially out of the sediment, either on the sediment surface, swimming in the overlying water, or floating toxicity tests with test materials spiked into sediments at the water surface, should different concentrations may be recorded. Amphipods that are caught reported in the surface film should be gently pushed down into the water. Any pertinent observations on the appearance terms of the sediment (such as color, presence of non-test organisms, growth of mold an LC50, EC50, IC50, NOEC, or algae, or depth of oxidized layer) should be recorded.

13.8.2 Mortality—The primary effect of LOEC. The statistical approach for spiked sediment toxicity is mortality tests also applies to the analysis of data from water-only reference-toxicity tests (Test Method E 1706).

13.2.4 Experimental Design—The guidance outlined below on the test amphipods, which analysis of data is determined at the end adapted from a variety of the exposure period. After daily observations have been made sources including Test Method E 1706, Guide E 1688, Guide E 1847, USEPA 1979b, 1993c, 1993b, (132, 161, 167), USEPA-USACE 1977, 1991, 1998 (114, 115, 116), Practices E 29, E 105, E 122, E 178, E 141, and any necessary samples have been taken, the contents Terminologies E 456, E 1325,

and E 1402. The objectives of the a sediment test chambers should be sieved are to remove the quantify contaminant effects on or accumulation in test species. Use of a larger screen size sieve for initiation organisms exposed to natural or spiked sediments or dredged materials and to determine whether these effects are statistically different from those occurring in a smaller screen size sieve for termination reduces the possibility control or reference sediment. Each experiment consists of losing small amphipods through the screen at termination. Screen sizes are specific for various test species (see annexes). Material retained on least two treatments: the screen should be washed into a sorting tray with clean sea water. The total numbers of live control and dead amphipods of the one or more test species should be recorded. treatment(s). The sum test treatment(s) consist(s) of these numbers might be less than the number of amphipods at T_0 because of decomposition. If the test species contaminated or potentially contaminated sediment(s). A control sediment is naturally present in always required to ensure that no contamination is introduced during the test sediment, the total number of live experimental setup and dead amphipods might exceed the number at T_0 . Amphipods that test organisms are inactive but not obviously dead should be observed under a lowpower microscope and should be counted as alive if there healthy. A control sediment is any sign of movement, such as a neuromuscular pleopod twitch. Gentle prodding may be used in an attempt to elicit movement.

13.8.3 *Reburial*—Data on judge the acceptability of the amphipods to rebury in clean test. Some designs will also require a reference sediment a t that represents an environmental condition or potential treatment effect of the sediment toxicity test can be interest. Controls are used to detect biologically important sublethal effects. Amphipods that survive evaluate the test should be transferred to dishes containing a layer acceptability of clean, 0.5 mm sieved control sediment. Sediment saved from the pretest holding containers test (Annex A1 and kept either in flowing sea water or at 4°C Annex A2) and might b include app control sediment or a formulated sediment (section 7.2). Testing a reference sediment provides a site-specific basis for use as reburial sediment. The numbers evaluating toxicity of amphipods able to bury within the time period specified for the species should be recorded. These data are used to document sublethal effects on behavior, and can be used to calculate an EC50. Infaunal amphipods unable to rebury are very unlikely to survive in nature. Toxicity data can therefore be analyzed in relation to *effective mortality*, that is, the sum test sediments. Comparisons of dead individuals plus those survivors that are not able test sediments to rebury. EC50 calculations can be made on multiple reference or control sediments representative of the b physical characteristics of effective mortality. In most cases, amphipods that survive in a ten-day the test are able to rebury.

13.9 *Other Measurements sediment (that is, grain size, organic carbon) may be useful in these evaluations.*

13.3 Data Calculations:

13.9.1 *Field Sediment*—If the sediment to be tested is collected from a potentially contaminated site in the field, sediment samples should be collected from the same grab for analysis

13.3.1 Sediments spiked with known concentrations of various geochemical properties (see 10.2). A separate sample for faunal analyses is also desirable. These samples may chemicals can be stored under appropriate conditions for possible future analysis, after the results used to establish cause and effect relationships between chemicals and biological responses. Results of the sediment toxicity tests with test are known. Sediment Eh and pH should be measured both in the field and in the test chambers materials spiked into sediments at the beginning and at the end of the test. This is especially desirable for field sediments, that might contain high different concentrations of organic materials. All measurements should also may be taken reported in control samples.

13.9.2 *Laboratory Spiked Sediments*—In experiments in which terms of an LC50 (median lethal concentration), an EC50 (median effect concentration), an IC50 (inhibition concentration), or as an NOEC (no observed effect concentration) or LOEC (lowest observed effect concentration). Most studies with spiked sediment are often started only a known test material is few days after the chemical has been added to sediment, the concentration of the test material sediment. Consistent spiking procedures should be determined followed in stock solutions or mixtures added order to sediment, and make interlaboratory comparisons (section 10.3).

13.3.2 Evaluating effect concentrations for chemicals in test chambers at the beginning and at the end sediment requires knowledge of factors controlling the test. Sea water and sediment samples can be taken as test chambers are loaded, and small water samples can be taken from the test chambers containing amphipods. To monitor changes bioavailability. Similar concentrations of a chemical in sediment or interstitial water chemistry during the course units of the experiment, separate sediment chemistry beakers should be set up and sampled at the initiation and at the termination mass of the experiment. It is not necessary to add amphipods to chemistry chambers sampled at the initiation chemical per mass of the experiment, but amphipods should be added to those sampled later. Some sediment and water quality characteristics, such as pH, Eh, and dissolved oxygen, can be measured by inserting analytical probes into the test chambers containing amphipods. If radiolabeled test compounds are used, separate chemistry beakers might not be necessary.

13.9.2.1 The concentration dry weight often exhibit a range in toxicity in different sediments (DiToro et al. 1991, USEPA 1992c (70, 95)). Effect concentrations of test material chemicals in sediment have been correlated to interstitial water concentrations, and sediment should be measured at several effect concentrations and as in interstitial water are often as practicable during similar to effect concentrations in water-only exposures. The bioavailability of nonionic organic compounds are often inversely correlated with the test. At a minimum, the organic carbon concentration of the test material should be measured at sediment. Whatever the beginning and at the end route of exposure, the test in the control and at low, medium, and high concentrations. Measurement correlations of degradation products of the test material might also be desirable.

13.9.2.2 Measurement of test material effect concentrations to interstitial water concentrations indicate predicted or measured

concentrations in interstitial water can be accomplished by pipetting water samples through glass or fluorocarbon plastic tubing from a point midway useful for quantifying the exposure concentration to an organism. Therefore, information on partitioning of chemicals between top, bottom, solid and liquid phases of sediment may be useful for establishing effect concentrations.

13.3.3 Toxic units can be used to help interpret the test chamber. Water samples should not contain any surface scum, any material from the sides response of the test chamber, or any sediment.

13.9.2.3 Samples for measurement organisms to multiple chemicals in sediment. A toxic unit is the concentration of a chemical divided by an effect concentrations. For example, a toxic unit of test material in sediment exposure can be taken calculated by siphoning off dividing the overlying measured concentration of a chemical in pore water without disturbing by the surface of the sediment, and then taking appropriate aliquots of the sediment water-only LC50 for the same chemical analysis.

13.9.2.4 Interstitial water can (Ankley et al. 1991a, (197)). Toxic units could also be sampled calculated by using dividing the water that comes to the surface concentration in a rolling mill jar or in a sample container as the whole sediment settles, by centrifuging a sediment sample to separate the sediment particles from the interstitial water, or by using a filter apparatus to extract interstitial water from a threshold concentration in whole sediment sample. Care should (Kemble et al. 1994, Long and Morgan 1991, (82, 96)). Toxicity expressed as toxic units may be taken to ensure that test materials do not undergo transformation, degradation, or volatilization during sample preparation. It should be kept in mind that filtering summed and this may provide information on the toxicity of chemical mixtures (Ankley et al. 1991a,(197)).

13.3.4 Field surveys can remove certain test materials from solution.

13.9.3 All Tests—Temperature should be recorded in designed to provide either a separate temperature beaker throughout qualitative reconnaissance of the test. If test chambers distribution of sediment contamination or a quantitative statistical comparison of contamination among sites (Burton and Ingersoll 1994, (198). Surveys of sediment toxicity are in usually part of more than one temperature controlled water bath, a temperature beaker should be set up in each water bath. Temperature should be monitored at least hourly using a recording thermometer or the daily maximum comprehensive analyses of biological, chemical, geological, and minimum temperatures should hydrographic data. Statistical correlation can be monitored (see Guide E-729). Individual temperature measurements should not vary by more than 3°C improved and the time-weighted average costs reduced if subsamples are taken simultaneously for sediment tests, chemical analyses, and benthic community structure.

13.3.5 Descriptive methods such as toxicity tests with field-collected sediment should not differ by more than 1°C from be used alone to evaluate sediment contamination. An integration of several methods using the designated test temperature (see 13.2).

14. Analytical Methodology

14.1 If samples weight of sediments or overlying water from test chambers, stock solutions, test sediment, or interstitial water are not evidence is needed to be analyzed immediately, they should be handled and stored appropriately (33) to minimize loss assess the effects of test material or contaminants through such processes as microbial degradation, hydrolysis, oxidation, photolysis, reduction, sorption and volatilization (see Practice D 3976).

14.2 Chemical and physical associated with sediment (Long et al. 1990; Ingersoll et al.1996; Ingersoll et al. 1997; MacDonald et al. 1996, (199, 98, 99, 97)). Hazard evaluations integrating data should be obtained using appropriate ASTM standards whenever possible. For those measurements for which ASTM standards do not exist or are not sensitive enough, methods should be obtained from laboratory exposures, chemical analyses, and benthic community assessments (the Sediment Quality Triad) provide strong complementary evidence of the degree of pollution-induced degradation in aquatic communities (Burton 1991, (369), Chapman et al. 1992, 1997, (100, 101); Canfield et al.1994, 1996, 1998, (200, 118, 201) .

14.3 The analytical method

13.3.6 Toxicity Identification Evaluation (TIE) procedures can be used to help provide insights as to specific contaminants responsible for toxicity in sediment (Ankley and Thomas 1992 (58), Ankley et al. 1991a,(197)). For example, the concentration toxicity of contaminants such as metals, ammonia, hydrogen sulfide, and nonionic organic compounds can be identified using TIE procedures.

14. Report

14.1 The record of the results of an acceptable sediment test chambers should be validated before beginning include the following information esit. The pr directly or by referencing available documents:

14.1.1 Name of test and investigator(s), name and location of laboratory, and dates of start and end of test.

14.1.2 Source of the method should be checked using reference control or split samples, interlaboratory comparisons, or alternative (preferably reference or corroborative) methods test sediment, method for collection, handling, shipping, storage, and disposal of analysis:

14.4 Concentrations sediment.

14.1.3 Source of test materials in interstitial water should be measured as well as material, lot number if applicable, composition (identities and concentrations of major ingredients and impurities if known), known chemical and physical properties, and the bulk sed identity and concentration(s-I) of any solvent used.

14.1.4 Source and characteristics of overlying water, description of any pretreatment, and results of any demonstrations of the total concentration ability of test material in interstitial water an organism to survive or grow in the overlying water from water.

14.1.5 Source, history, and age of test chambers, measurement organisms; source, history, and age of brood stock, culture

procedures; and source and date of collection of the apparent dissolved or free form test organisms, scientific name, name of person who identified the organisms and the taxonomic key used, age or life stage, means and ranges of weight or length, observed diseases or unusual appearance, treatments, holding, and acclimation procedures.

14.1.6 Source and composition of food, concentrations of test material is desirable. The free form for organic contaminants is that which is not bound and other contaminants, procedure used to prepare food, feeding methods, frequency, and ration.

14.1.7 Description of the experimental design and test chambers, the depth and volume of sediment and overlying water in the chambers, lighting, number of test chambers and number of test organisms/treatment, date and time test starts and ends, temperature measurements, dissolved organic carbon, oxygen concentration (as percent saturation), and for metals it is any aeration used before starting a test and during the ionic form conduct of a test.

14.1.8 Methods used for physical and chemical characterization of sediment.

14.1.9 Definition(s) of the element. The “apparent dissolved” fraction is usually defined effects used to calculate LC50 or EC50s, biological endpoints for tests, and determined as that which passes through a 0.45-µm membrane filter. However, passing solutions through membrane filters can result summary of general observations of other effects.

14.1.10 Methods used for statistical analyses of data: (1) summary statistics of the transformed or raw data as applicable (for example, mean, standard deviation, coefficient of variation, precision and bias); (2) hypothesis testing (raw data, transformed data, null hypothesis, alternate hypothesis, target Type I and II error rates, statistics used (including calculation of test statistic)), decision rule used (for example, *W* statistic >0.65 results in the rejection of the null hypothesis), calculated test statistic and decision rule result, achieved Type I and II error rates (for some discrete tests, achieved error rates only approximate the target rates); (3) results of regression analyses (parameters of regression fit, uncertainty limits on the regression parameters, correlation coefficient).

14.1.11 Summary of general observations on other effects or symptoms.

14.1.12 Anything unusual about the test, any deviation from these procedures, and any other relevant information.

14.2 Published reports should contain enough information to clearly identify the methodology used and the quality of the results.

15. Precision and Bias

15.1 Determining Precision and Bias :

15.1.1 Precision is a term that describes the degree to which data generated from replicate measurements differ and reflects the closeness of Test

15.1 A 10-day agreement between randomly selected test results. Bias is the difference between the value of the measured data and the true value and is the closeness of agreement between an observed value and an accepted reference value (Practices E 177 and E 691). Quantitative determination of precision and bias in sediment testing of aquatic organisms is difficult or may be impossible in some cases, as compared to analytical (chemical) determinations. This is due, in part, to the many unknown variables which affect organism response. Determining the bias of a sediment test using field samples is not possible since the true values are not known. Since there is no acceptable reference material suitable for determining the bias of sediment tests, bias of the procedures described in this standard has not been determined (section 15.2).

15.1.2 Sediment tests exhibit variability due to several factors. Test variability can be described in terms of two types of precision, either single laboratory (intralaboratory or repeatability; section 15.6.1) precision or multilaboratory (interlaboratory or reproducibility; sections 15.5.2 and 15.6) precision (also referred to as round-robin or ring tests). Intralaboratory precision reflects the ability of trained laboratory personnel to obtain consistent results repeatedly when performing the same test on the same organism using the same toxicant. Interlaboratory precision is a measure of how reproducible a method is when conducted by a large number of laboratories using the same method, organism, and toxic sample. Generally, intralaboratory results are less variable than interlaboratory results (USEPA 1993a, Swartz 1989, USEPA 1993b, Marcus and Holtzman 1988, Grothe and Kimerle 1985, Pittinger et al. 1989, (132, 120, 167, 202, 203, 204)).

15.1.3 A measure of precision can be calculated using the mean and relative standard deviation (percent coefficient of variation, or CV % = standard deviation/mean × 100) of the calculated endpoints from the replicated endpoints of a test. However, precision reported as the CV should not be the only approach used for evaluating precision of 10% tests and should not be used for the NOEC effect levels derived from statistical analyses of hypothesis testing. The CVs may be very high when testing extremely toxic or nontoxic samples. For example, if there are multiple replicates with no survival and one with low survival the CV may exceed 100 %, yet the range of response is actually quite consistent. Therefore, additional estimates of precision should be used, such as range of responses and minimum detectable differences (MDD) compared to control survival or growth. Several factors can affect the precision of the test, including test organism age, condition, sensitivity, handling, and feeding of the test organisms, overlying water quality, and the experience in conducting tests. For these reasons, it is recommended that trained laboratory personnel conduct the tests in accordance with the procedures outlined in Annex A1 and in Annex A2. Quality assurance practices should include: (1) single laboratory precision determinations that are used to evaluate the ability of the laboratory personnel to obtain precise results using reference toxicants for each of the test organisms and (2) preparation of control charts (Figure 16 in Test Method E 1706) for each reference toxicant and test organism. The single laboratory precision determinations should be made before conducting a sediment test and should be periodically performed as long as whole-sediment tests are being conducted at the laboratory.

15.1.4 Intralaboratory precision data are routinely calculated for test organisms using water-only 96-h exposures to a reference

toxicant such as Cd Cl₂. Intralaboratory precision data should be tracked using a control chart. Each laboratory's reference-toxicant data will reflect conditions unique to that facility, including dilution water, culturing, and other variables (Section 11). However, each laboratory's reference toxicant CVs should reflect good repeatability.

15.1.5 Two interlaboratory precision (round-robin) tests have been completed using 10-d whole sediment tests, one with *Rhepoxynius abronius* (Mearns et al. 1986 (205)), and the other with *Ampelisca abdita*, *Eohaustorius estuarius*, and *Leptocheirus plumulosus* (Schlekat et al. 1995 (21)). The results of these round-robin studies are described in section 15.5.

15.1.6 One interlaboratory precision test has been completed on the 28-d chronic test with *Leptocheirus plumulosus* (DeWitt et al., 1997b (25)). Ten laboratories participated in the round-robin study, which used a dilution series of highly contaminated Black Rock Harbor sediment from a Superfund site in Connecticut mixed with uncontaminated, diluent sediment from Sequim Bay, Washington. The results of this round-robin study are described in section 15.6.

15.2 *Bias*—The bias of toxicity tests cannot be determined since there is no acceptable reference material. The bias of the reference-toxicity tests can only be evaluated by comparing test responses to control charts.

15.3 *Replication and Test Sensitivity*— The sensitivity of sediment tests will depend in part on the number of replicates per concentration, the probability levels (alpha and beta), and the type of statistical analysis. For a specific level of variability, the sensitivity of the test will increase as the number of replicates is increased. The minimum recommended number of replicates varies with the objectives of the test and the statistical method used for analysis of the data (Section 13).

15.4 *Demonstrating Acceptable Laboratory Performance:*

15.4.1 Intralaboratory precision, expressed as a coefficient of variation (CV), of the range for each type of test to be used in a laboratory can be determined by performing five or ~~show signs~~ more tests with different batches of test organisms, using the same reference toxicant, at the same concentrations, with the same test conditions (for example, the same test duration, type of water, age of test organisms, feeding), and same data analysis methods. A reference-toxicant concentration series (0.5 or ~~stress~~; higher) should be selected that will consistently provide partial mortalities at two or more concentrations of the test chemical (section 11.14 and Table 4). See section 11.16 for additional detail regarding reference-toxicity testing.

15.4.2 Before conducting tests with potentially contaminated sediment, it is strongly recommended that the laboratory conduct the tests with control sediment(s) alone. Results of these preliminary studies should be used to determine if the use of the control sediment and other test conditions (that is, water quality) result in ~~an individual~~ acceptable performance in the tests as outlined in Annex A1 and Annex A2.

15.4.3 A control chart can be prepared for each combination of reference toxicant and test organism. Each control chart should include the most current data. Endpoints from five tests are adequate for establishing the control charts. In this technique, a running plot is maintained for the values (X_i) from successive tests with a given reference toxicant (See Figure 16 in Test Method E 1706), and the endpoint (LC50, NOEC, ICp) are examined to determine if they are within prescribed limits. Control charts as described in USEPA 1993a, 1993b, (1320-%, 167) are used to evaluate the cumulative trend of results from a series of samples. The mean and upper and lower control limits (± 2 SD) are recalculated with each successive test result.

15.24.4 A ~~+~~The outliers, which are values falling outside the upper and lower control limits, and trends of increasing or decreasing sensitivity, are readily identified using control charts. With an alpha of 0.05, one in 20 tests would be expected to fall outside of the control limits by chance alone. If 2 of 20 reference-toxicity tests fall outside the control limits, the sediment toxicity tests conducted during the time in which the second reference-toxicity test failed are suspect, and should ~~usually~~ be considered as provisional and subject to careful review.

15.4.5 A sediment test may be acceptable if specified conditions of a reference-toxicity test fall outside the expected ranges (section 11.10). Specifically, a sediment test should not be judged unacceptable if the LC50 for a given reference-toxicity test falls outside the expected range or if control survival in the reference-toxicity test is <90 %. All the performance criteria outlined in Annex A1 and Annex A2 should be considered when determining the acceptability of a sediment test. The acceptability of the sediment test would depend on the experience and judgment of the investigator and the regulatory authority.

15.4.6 If the value from a given test with the reference toxicant falls more than two standard deviation (SD) outside the expected range, the sensitivity of the ~~following~~ occurred:

15.2.1 ~~AH~~ organisms and the overall credibility of the test system may be suspect USEPA 1993a, (132) . In this case, the test procedure should be examined for defects and should be repeated with a different batch of test organisms.

15.4.7 Performance should improve with experience, and the control limits for point estimates should gradually narrow. However, control limits of ± 2 SD, by definition, will be exceeded 5 % of the time, regardless of how well a laboratory performs. Highly proficient laboratories which develop a very narrow control limit may be unfairly penalized if a test which falls just outside the control limits is rejected *de facto*. For this reason, the width of the control limits should be considered in determining whether or not identical.

15.2.2 Treatments were not randomly assigned an outlier is to be rejected. This determination should may be made by the regulatory authority evaluating the data.

15.4.8 The recommended reference-toxicity test ~~chambers~~.

15.2.3 Test organisms were not randomly consists of a control and five or ~~impartially~~ distributed more concentrations in which the endpoint is an estimate of the toxicant concentration which is lethal to 50 % of the test ~~chambers~~.

15.2.4 Required negative, reference sediment, positive or solvent controls were not included organisms in the time period

prescribed by the test. The LC50 is determined by an appropriate procedure, such as the trimmed Spearman-Kärber Method, Probit Method, Graphical Method, or the Linear Interpolation Method (Section 13 and Test Method E 1706).

15.24.59 ~~The point estimation analysis methods recommended in this test method have been chosen primarily because they are well-tested, well-documented, and are applicable to most types of test data. Many other methods were considered in the selection process, and it is recognized that the methods selected are not from the same population, were only possible methods of analysis for toxicity data.~~

15.5 Precision of 10-d Amphipod Sediment Toxicity Test Methods:

15.5.1 Intralaboratory Precision—~~Intralaboratory precision has not~~ been evaluated for any of the four amphipod species described in Annex A1.

15.5.2 Interlaboratory Precision:

15.5.2.1 Interlaboratory precision for *R. abronius* using 10-d whole sediment toxicity tests using the methods described in this standard (Table A1.1 in Annex A1) is described by Mearns et al. 1986 (205). Five laboratories participated in the study, including federal and state government laboratories, a contract laboratory, and an academic laboratory. The laboratories were chosen because each had demonstrated experience in sediment toxicity tests with *R. abronius*. The experimental design required each laboratory to conduct 10-d whole sediment tests on a total of acceptable quality.

15.2.6 ~~Amphipods~~ 7 sediment treatments. One control sediment was tested. Three sediment treatments consisted of control sediment that was amended with CdCl₂ to result in the following measured concentrations: 4, 8, and 12 mg Cd/kg dry weight. Three field-collected sediments were also used. They were collected from a wild population the following locations in Puget Sound, WA: Central Basin (Metro Seattle Station A600E), inner Sinclair Inlet, and Slip No. 1 in City Waterway, Commencement Bay.

15.5.2.2 Amphipods were collected from a depth of 6 m off West Beach, Whidbey Island, WA, and distributed to each participating laboratory. Each laboratory used its own source of clean seawater.

15.5.2.3 All five laboratories had >90 % survival in control sediment, and thereby met the laboratory performance criteria for more than two weeks, unless the test. Mean survival in control sediment was 96.4 %, the CV was 3.7 %, and the range in mean survival was from 92 to 100 % (Table 7). Of the cadmium-spiked sediments, survival was the least variable in the 4 mg/kg Cd treatment. Mean survival was 96.2 %, the CV was 4.2 %, and the range was from 89 to 98 %. The most variable response was in the 12 mg/kg Cd sediment. Mean survival was 19 %, the CV was 79.1 %, and the range was from 6 to 41 %. City Waterway showed the least variability among the field-collected sediments, with a mean survival of 83 %, a CV of 6.4 %, and a range from 74 to 87 %. Sinclair Inlet showed the greatest variability among the field-collected sediments, with a mean survival of 78.8 %, a CV of 11.3 %, and a range from 67 to 88 %.

15.5.2.4 Interlaboratory precision for *A. abdita*, *E. estuarius*, and *L. plumulosus* using 10-d whole sediment toxicity tests is described in Schlekot et al. (1995 (21)). Details of this study are described below. The number of participating laboratories varied with the test species: six for *A. abdita*, eight for *E. estuarius*, and seven for *L. plumulosus*. Laboratories were chosen on the basis of demonstrated experience with the particular test species. Each laboratory has been shown to have no significant effect conducted 10-d sediment toxicity tests on sensitivity.

15.2.7 The 4 sediment treatments. Sediment treatments were selected for each species to include one negative control sediment and three contaminated sediments. Highly contaminated sediment from Black Rock Harbor, CT, was diluted with species-specific, non-contaminated control sediment, creating test- θ sediments that ranged in relative contamination from low to high.

15.5.2.5 Independent suppliers distributed amphipods to each laboratory. *Ampelisca abdita* and *E. estuarius* were field-collected from locations in Narragansett, RI, and Newport, OR, respectively. *Leptocheirus plumulosus* were obtained from cultures located at the University of Maryland, Queenstown, MD. Each laboratory used its own supply of clean seawater.

15.5.2.6 Mean survival of *A. abdita* in control sediment ranged from 85 to 100 % (Table 8). Five of the six laboratories achieved greater than 90 % survival in control sediment, which is the minimum survival that must be obtained in control sediment in order for the test temperature to be accepted. The grand mean was 94.5 %, and the CV was 5.5. A dose response was exhibited with decreasing survival with increasing proportions of BRH sediment. Test sediments (that is, 7, 25, and 33 % BRH dilutions) exhibited a higher degree of variability than in control sediment. In 7 % BRH sediment, mean survival ranged from 20 % in Laboratory 5 to 97 % in Laboratory 6 (Table 8). Twenty-percent BRH exhibited the greatest magnitude of variability, with a range

TABLE 7 Inter-laboratory Precision for Survival of *Rhepoxynius abronius* in 10-d Whole Sediment Toxicity Tests Using Seven Sediments (Mearns et al. 1986 (205), USEPA 1994a (1))

Lab	Percent Survival (SD) in Sediment Samples						
	Control	4 mg/kg Cd	8 mg/kg Cd	12 mg/kg Cd	Central Basin	Sinclair Inlet	City Waterway
1	92 (7)	89 (7)	87 (9)	8 (3)	83 (11.5)	78 (13)	74 (11.5)
2	96 (4)	98 (3)	90 (10)	41 (11)	69 (7.5)	67 (11)	87 (12)
3	100 (0)	97 (3)	78 (10.5)	12 (7.5)	90 (8)	87 (7.5)	83 (12.5)
4	94 (7)	99 (2)	50 (15)	6 (5.5)	92 (5.5)	88 (3)	84 (11)
5	100 (0)	98 (4.5)	77 (3)	28 (11.5)	80 (3.5)	74 (9)	87 (3)
Mean	96.4 (3.6)	96.2 (4.1)	76.4 (15.8)	19 (15.5)	82.8 (9.1)	78.8 (8.9)	83 (5.3)
CV (%)	3.7	4.2	20.7	79.1	11.0	11.3	6.4

TABLE 8 Inter-laboratory Precision for Survival of *Ampelisca abdita* in 10-d Whole Sediment Toxicity Tests Using Four Sediments (USEPA 1994a (1))

Percent Survival (SD) in Sediment Samples				
Lab	Control	7 % Black Rock Harbor	20 % Black Rock Harbor	33 % Black Rock Harbor
1	97.0 (4.5)	63.0 (19.6)	10.0 (7.9)	6.0 (4.2)
2	94.0 (8.9)	75.0 (6.1)	7.0 (4.5)	0.0 (0)
3	97.0 (4.5)	90.0 (3.5)	36.0 (9.6)	38.0 (14.4)
4	94.0 (8.9)	79.0 (17.8)	7.0 (4.5)	3.0 (6.7)
5	85.0 (7.1)	20.0 (12.7)	1.0 (2.2)	1.0 (2.2)
6	100.0 (0)	97.0 (4.5)	90.0 (5.0)	72.0 (13.0)
Mean	94.5 (5.2)	70.7 (13.0)	25.2 (34.0)	20.0 (29.2)
CV (%)	5.5	38.9	135.1	146.2

of 1 to 90 %. Thirty-three percent BRH also exhibited considerable variability. The overall rank of sediment toxicity as measured by absolute mortality was consistent among laboratories. One hundred percent of laboratories were in agreement for in ranking control and 7 % BRH sediments as the first and second least toxic sediments, respectively (Table 8).

15.5.2.7 Every laboratory surpassed the minimum survival criteria of 90 % survival in control sediment with *E. estuarius*. The range was from 96 to 100 %, with a grand mean of 98.2 % and a CV of 1.5 (Table 9). Grand mean survival decreased with increasing proportions of BRH. BRH sediment dilutions exhibited greater variability than control sediment, with 25 % BRH displaying the highest coefficient of variation. All eight laboratories ranked survival of *E. estuarius* for control and 9 % BRH as the least and second least toxic, respectively (Table 8). With the exception of Laboratories 1 and 8, the rank for 25 and 42 % BRH were appropriately third and fourth least toxic, respectively.

15.5.2.8 *Leptocheirus plumulosus* exhibited a range of survival in control sediment from 86 to 99 % (Table 10). The grand mean was 91.8 %, and the test chambers:

15.2.8 Temperature, dissolved oxygen, CV was 4.7. Two laboratories, 3 and 5, failed to meet the minimum control sediment survival criteria of 90 %. Grand means displayed a dose response of decreasing survival with increasing proportion of BRH sediment. Coefficients of variation were uniformly higher in BRH sediment dilutions as compared to control sediment, but did not vary greatly among BRH sediments (Table 10). Laboratory 1 appeared to be an outlier with respect to survival in BRH sediment dilutions, as survival of *L. plumulosus* was the lowest for all three BRH sediments for any laboratory. The rank of sediments according to their toxicity was generally consistent among laboratories. Agreement was 100 % for control and the highest BRH sediment; these were appropriately ranked 1 and 4, respectively (Table 10). Laboratories 4 and 5 anomalously ranked 10 and 28 % BRH as 3 and 2, respectively, whereas the remaining laboratories ranked these sediments appropriately according to the proportion of BRH.

15.5.3 These tests exhibited similar or better precision than many chemical analyses and effluent toxicity testing methods (USEPA, 1991 (127)). The success rate for test initiation and completion of this round-robin evaluation is a good indication that a well equipped and trained staff will be able to successfully conduct this test. This is an important consideration for any test performed routinely in any regulatory program.

15.6 Precision of the 28-d Sediment Toxicity Test Methods with *Leptocheirus plumulosus*:

15.6.1 Intralaboratory Performance—Studies described in DeWitt et al. (1997b (25)) provide data that can be used to characterize intralaboratory precision with the 28-day long-term toxicity test with *L. plumulosus* . These data provide an estimate of intralaboratory precision from a single laboratory from a total of 88 treatments (Table 11). To be consistent with standard statistical procedures, these data were transformed to reduce the heterogeneity of within class variance. Percent survival was

TABLE 9 Interlaboratory Precision for Survival of *Eohaustorius estuarius* in 10-d Whole Sediment Toxicity Tests Using Four Sediments (USEPA 1994a (1))

Percent Survival (SD) in Sediment Samples				
Lab	Control	9 % Black Rock Harbor	25 % Black Rock Harbor	42 % Black Rock Harbor
1	96.0 (6.5)	45.0 (19.7)	6.0 (6.5)	16.0 (9.6)
2	98.0 (2.7)	76.0 (10.8)	46.0 (13.9)	25.0 (7.1)
3	97.0 (2.7)	89.0 (4.2)	59.0 (10.8)	45.0 (10.0)
4	98.8 (2.7)	59.0 (23.0)	47.2 (23.2)	45.8 (27.0)
5	100.0 (0)	75.0 (19.7)	36.0 (12.4)	16.0 (9.6)
6	100.0 (0)	69.0 (12.9)	56.0 (18.8)	38.0 (14.4)
7	99.0 (2.2)	79.0 (6.5)	61.0 (10.8)	50.0 (7.9)
8	97.0 (6.7)	53.0 (14.4)	24.0 (14.7)	29.0 (15.6)
Mean	98.2 (1.5)	68.1 (14.7)	41.9 (19.1)	33.1 (13.5)
CV (%)	1.5	21.6	45.5	40.9

TABLE 10 Interlaboratory Precision for Survival of *Leptocheirus plumulosus* in 10-d Whole Sediment Toxicity Tests Using Four Sediments (USEPA 1994a (1))

Percent Survival (SD) in Sediment Samples				
Lab	Control	10 % Black Rock Harbor	28 % Black Rock Harbor	47 % Black Rock Harbor
1	91.3 (4.8)	6.0 (4.2)	5.0 (3.5)	2.5 (2.9)
2	91.0 (8.9)	62.0 (11.0)	51.0 (15.6)	33.0 (11.5)
3	88.0 (8.4)	34.0 (15.2)	22.0 (13.0)	7.0 (5.7)
4	92.0 (7.6)	48.0 (23.9)	59.0 (21.6)	27.0 (10.4)
5	86.0 (10.2)	20.0 (9.4)	28.0 (4.5)	12.0 (9.1)
6	95.0 (6.1)	76.0 (10.2)	65.0 (14.6)	38.0 (17.5)
7	99.0 (2.2)	78.0 (13.0)	56.0 (4.2)	26.0 (6.5)
Mean	91.8 (4.3)	46.3 (27.7)	40.9 (22.6)	20.8 (13.6)
CV (%)	4.7	59.8	55.2	65.5

TABLE 11 Intralaboratory Precision Distribution of the Coefficient of Variation for Each Test Endpoint (DeWitt et al. 1997a; USEPA-USACE 2001 (2))

Endpoint	Sample Size	Mean	Median	Minimum	Maximum	1st Quartile	3rd Quartile
% Survival (Arcsine transformed)	88	14 %	11 %	0 %	173 %	8 %	14 %
Growth rate (log transformed)	87	4 %	3 %	0 %	16 %	2 %	6 %
Reproduction (square root transformed)	88	31 %	18 %	0 %	141 %	13 %	36 %

transformed to the arcsine-square root of the value; growth rate was transformed to the natural logarithm of the value; and reproduction (offspring per survivor) was transformed to the arcsine -square root of the value. A CV was calculated on the transformed data for each treatment within an experiment. The observed distribution obtained from the resulting sample of CVs from all experiments was then characterized. This distribution of CVs then provides an appropriate range on which to base sample size calculations for future experiments. The median CVs were 11 % for survival, 3 % for growth rate, and 18 % for reproduction (Table 11). The range between the first and third quartiles provides a useful nonparametric interval bounding the distribution. This range was 8 to 14 % for survival, 2 to 6 % for growth rate, and 13 to 36 % for reproduction (Table 11). These values are similar to CVs for intralaboratory precision calculated for survival from 10-d tests with control sediment using *Hyalella azteca* and *Chironomus tentans* (7.2 % and 5.7 %, respectively; USEPA 2000 and Test Method E 1706).

15.6.2 Interlaboratory Precision:

15.6.2.1 Interlaboratory precision for *L. plumulosus* in Section 13:

~~15.2.9 Aeration~~ the 28-d whole sediment toxicity test using methods similar to those described in this standard (Table A2.1 in Annex A2) was evaluated by round-robin testing (DeWitt et al., 1997b (25)). Ten laboratories, including federal and state government laboratories, contract laboratories, and academic laboratories with demonstrated experience in chronic toxicity testing using *L. plumulosus*, participated in round-robin toxicity testing (DeWitt et al., 1997b (25)). The experimental design required each laboratory to conduct the 28-d chronic test using a dilution series of Black Rock Harbor sediment (BRH; a Superfund site in Connecticut) mixed with clean, diluent sediment from Sequim Bay, Washington. Each sediment treatment was prepared in a single batch that was subsampled and shipped to testing laboratories. A total of four concentrations of BRH sediment and one negative control sediment were tested. Across all treatments, total organic carbon averaged 2.6 % dry weight, total solids averaged 33 %, and grain size averaged 15 % sand, 42 % silt, and 43 % clay. In general, cadmium, chromium, copper, lead, nickel, and zinc, as well as total PAHs, increased along the dilution series gradient. Table 12 summarizes the concentration ranges for the inorganic contaminants.

15.6.2.2 About 4 months before the start of the round-robin study, laboratories not currently maintaining cultures of *L. plumulosus* were supplied with amphipods, sediment, food, and culturing methods by the Battelle Marine Sciences Laboratory (MSL). Each laboratory maintained cultures following the culturing method detailed in DeWitt et al. (1997a (7)). Each laboratory used its own source of clean seawater.

TABLE 12 Ranges of the BRH Sediment Dilution Series Chemical Concentrations (mg/kg dry wt; from DeWitt et al., 1997b (25) and USEPA-USACE 2001 (2))

	Low (BRH treatment)	High (BRH treatment)
Cadmium	4.09 (0.0 %)	13.5 (15.1 %)
Chromium	104 (0.0 %)	767 (15.1 %)
Copper	104 (0.0 %)	1503 (15.1 %)
Lead	31.1 (0.0 %)	209 (15.1 %)
Nickel	91.2 (0.0 %)	150 (15.1 %)
Zinc	189 (0.0 %)	736 (15.1 %)
Total PAHs	9.85 (1.4 %)	17.5 (15.1 %)

15.6.2.3 Of the ten laboratories participating in the round-robin, only five laboratories had >80 % survival in the negative control sediment, and thereby met this performance criterion for test acceptability (Top of Table 13). Analysis of the data resulting from the round-robin included only these five laboratories. Mean survival in the negative control sediment was 93.6 %, the CV was 4.2 %, and the range was from 89 to 98 % (Table 13). The CVs across laboratories from the five treatments ranged from 3.1 to 12.8 %, with a mean of 8.4 %, and increased with dose. None of the laboratories produced less than 70 % survival, even in the highest concentration of saturation.

15.2.10 The BRH sediment. Further, none of the laboratories produced a monotonic dose response for survival. This suggests that the test did not contain a wide enough series of dilutions to adequately measure the response of survival. For those laboratories that showed a statistically significant decrease in survival in the highest concentration of solvent BRH (n=4), an average of 16 % change in survival was produced between the control and the highest concentration of BRH sediment.

15.6.2.4 For the five laboratories that met the performance criterion, interlaboratory precision for this study was characterized by the maximum and minimum CV for each endpoint. The minimum interlaboratory CV averaged about 4 % for survival, 14 % for growth rate, and 35 % for reproduction (Table 14). Maximum interlaboratory CV averaged 19 % for survival, 38 % for growth

TABLE 13 Results of Round-robin Interlaboratory Precision of Endpoint Sensitivity for *L. plumulosus* in a 28-d Long-term Toxicity Test Using Black Rock Harbor Sediments (DeWitt et al., 1997b (25 and USEPA-USACE 2001 (2))

A) Results for Laboratories that met Control Performance Criteria						
Concentration of Black Rock Harbor Sediment						
Lab	0.0 %	1.4 %	4.6 %	8.3 %	15.1 %	
Mean Percent Survival (%CV)						MDD %
4	89 (11.5)	92 (3.0)	82 (17.6)	76 (16.4)	73 (13.4)	16
6	96 (6.8)	93 (2.9)	97 (4.6)	95 (7.4)	96 (5.7)	8
7	90 (6.8)	88 (9.5)	84 (12.9)	92 (6.2)	82 (11.9)	13
8	95 (6.4)	92 (6.2)	72 (42.4)	74 (42.0)	70 (18.2)	31
9	98 (2.8)	96 (2.3)	84 (15.4)	91 (10.6)	86 (14.5)	14
Mean	93.6	92.2	83.8	85.6	81.4	
%CV	4.2	3.1	10.6	11.5	12.8	
MDD %	10	7	26	24	16	
Mean Growth Rate mg/d (%CV)						MDD mg/ind/d
6	0.059 (9.8)	0.054 (6.0)	0.046 (19.0)	0.039 (11.7)	0.020 (24.1)	0.009
7	0.084 (4.4)	0.075 (4.9)	0.063 (8.5)	0.053 (7.2)	0.035 (28.0)	0.009
8	0.045 (18.3)	0.031 (12.7)	0.036 (25.1)	0.024 (27.5)	0.014 (14.1)	0.010
9	0.089 (8.7)	0.078 (13.4)	0.065 (12.7)	0.060 (12.0)	0.045 (11.6)	0.012
Mean	0.063	0.057	0.049	0.039	0.025	
%CV	35.8	35.7	29.8	45.1	59.4	
MDD	0.014	0.014	0.017	0.012	0.011	
Mean Offspring per Survivor (%CV)						MDD # offspring
4	0.27 (14.1)	2.26 (72.3)	0.65 (14.9)	0.35 (56.5)	0.33 (81.2)	1.33
6	4.37 (41.0)	2.96 (53.8)	2.58 (27.5)	1.70 (43.4)	0.18 (76.6)	1.77
7	5.22 (55.7)	3.99 (40.5)	3.61 (42.5)	2.21 (75.4)	0.48 (65.6)	2.73
8	1.66 (65.8)	1.10 (54.2)	1.52 (29.8)	0.25 (91.5)	0.10 (108)	0.92
9	7.09 (30.8)	5.43 (21.9)	3.48 (29.8)	1.65 (60.7)	0.19 (99.0)	1.96
Mean	3.72	3.15	2.37	1.23	0.25	
%CV	73.8	52.5	53.8	71.2	59.5	
MDD	2.86	2.10	1.53	1.42	0.33	
B) Results for Laboratories that did not meet the Control Performance Criteria						
Concentration of Black Rock Harbor Sediment						
Lab	0.0 %	1.4 %	4.6 %	8.3 %	15.1 %	
Mean Percent Survival (%CV)						
1	53 (31.7)	74 (13.0)	65 (38.5)	58 (18.9)	39 (64.4)	
2	0 (-)	10 (-)	27 (137.1)	15 (-)	0 (-)	
3	72 (34.6)	85 (17.1)	74 (15.4)	61 (21.2)	55 (24.9)	
5	60 (56.5)	88 (18.7)	66 (29.5)	84 (24.7)	76 (11.8)	
10	69 (29.6)	59 (49.9)	58 (44.2)	37 (70.0)	25 (58.3)	
Mean Growth Rate mg/ind/d (%CV)						
1	0.024 (81.7)	0.032 (37.7)	0.012 (74.9)	0.012 (67.9)	0.008 (71.2)	
2	0 (-)	0.027 (-)	0.028 (49.0)	0.017 (-)	0 (-)	
3	0.050 (50.2)	0.067 (21.0)	0.055(33.3)	0.034(52.4)	0.025 (32.0)	
5	0.058 (16.0)	0.062 (31.7)	0.037 (67.6)	0.036 (43.0)	0.024 (12.0)	
10	0.006 (54.5)	0.014 (139)	0.007 (47.1)	0.003 (54.0)	0.003 (80.2)	
Mean Offspring per Survivor (%CV)						
1	0.7 (45.2)	1.7 (57.0)	0.4 (206)	0.1 (163)	0 (-)	
2	0 (-)	1.3 (-)	1.2 (18.0)	0.6 (-)	0 (-)	
3	4.8 (42.5)	3.7 (51.5)	3.4 (34.9)	0.4 (92.4)	0 (138)	
5	3.1 (80.8)	2.3 (25.5)	1.1 (136)	0.8 (113)	0.6 (117)	
10	0.1 (131)	1.4 (111)	0.5 (98.3)	0.8 (157)	0.3 (144)	

TABLE 14 Summary of Interlaboratory Precision at Five Laboratories for the 28-Day *Leptocheirus plumulosus* Chronic Test Using Five Dilutions of Black Rock Harbor Sediment (DeWitt et al. 1997b (25) and USEPA-USACE 2001(2))

	Lab-4	Lab-6	Lab-7	Lab-8	Lab-9
Survival					
Min CV (%)	3	3	6	6	2
Max CV (%)	18	7	13	42	15
Growth rate					
Min CV (%)	36	6	4	13	9
Max CV (%)	96	24	28	27	13
Offspring per Survivor					
Min CV (%)	56	27	40	30	22
Max CV (%)	149	77	75	108	99

rate, and 102 % for reproduction. The interlaboratory MDD for survival ranged from 8 to 31 %, and the intralaboratory MDD for survival ranged from 10 to 26 %. The interlaboratory MDD for growth rate ranged from 0.011 to 0.017 mg/individual/d, and the intralaboratory MDD for growth rate ranged from 0.009 to 0.024 mg/individual/d. The interlaboratory MDD for reproduction ranged from 0.33 to 2.86 offspring per survivor, and the intralaboratory MDD for reproduction ranged from 0.92 to 2.73 offspring per survivor. These MDDs should be interpreted cautiously, because they are derived from one study consisting of a small number of comparisons. Although the technical staff for laboratories participating in the round-robin had extensive sediment toxicity testing experience, many had limited testing experience specifically with *L. plumulosus*. Therefore, these values for interlaboratory precision may be higher than would be expected from laboratories with routine experience testing with this species.

15.6.2.5 A cost-power analysis was conducted on round-robin data to determine the number of replicates required per treatment for the 28-d whole-sediment standard testing using *L. plumulosus* (DeWitt et al., 1997b (25)). This analysis involved evaluating both the improvement in statistical power of the test to detect a difference between treatment means and the additional expense of adding more replicates. For this analysis, the cost of a replicate was assumed to be proportionate to the time required to conduct all of the tasks associated with one treatment. If cost was not a concern, 14 replicates would be optimal and would provide 80 % power for detecting a 30 % difference in reproduction at a CV of about 36 %. This number of replicates is impractical because of costs and logistics. The cost-power analysis for the *L. plumulosus* chronic test indicated that six replicates per treatment gives the greatest statistical power at the most efficient cost. However, this conclusion was based on the assumption that every 1 % increase in improved detection equals a 1 % increase in cost. The decision to specify 5 replicates per treatment in this standard was based primarily on an effort to keep the cost of performing this test to a minimum. Based on the median CVs for growth rate, survival, and reproduction calculated from a large data set (3 %, 11 %, and 18 %, respectively), five replicates will provide high power (>0.80) to detect a 20 % decrease in survival and growth rate endpoints relative to the control (see Figure 12.5 in USEPA-USACE 2001 (2)). For the reproduction endpoint, the power to detect a 20 % decrease will be closer to 0.40 using five replicates and 0.50 using six replicates. With power fixed at 80 % and at a CV of 20 %, the median CV demonstrated for reproduction with five replicates would be suitable to detect about 18 % reduction in reproduction and with six replicates

about 156 % reduction. Thus, there is relatively little gained by increasing the number of replicates from five to six. Nevertheless, if reproduction is the assessment endpoint of most concern, then incorporation of more than five replicates should be considered. Because space and cost considerations make use of five replicates desirable, this method would benefit from additional research to measure find ways to reduce the among-replicate variability for the reproduction endpoint.

15.6.2.6 The mean growth rates across the laboratories for each dose decreased with increasing concentration of toxicant in BRH sediment (Table 14). Thus, the test chamber growth rate was not validated before beginning a more sensitive measure to the test.

15.2.12 Response criteria concentration of BRH survival. The CVs across the laboratories from the five treatments ranged from 29.8 to 59.4 %, with a mean of 41.2 %, and were on average five times greater for growth rate than for survival (Table 14). Of the five laboratories that met the performance criterion for control survival, three laboratories produced a monotonic dose response to growth rate. The percentage of change in the growth rate between control and the highest concentration of BRH sediment was on average 58 % for these three laboratories.

15.6.2.7 The mean reproduction across laboratories for each dose decreased with increasing concentration of BRH sediment. Thus, the measure of reproduction was more sensitive to the concentration of BRH than was survival; however, the CVs across laboratories are on average eight times greater for reproduction than for survival. The CVs for the five treatments ranged from 52.5 % to 73.8 %, with a blind fashion, mean of 62.2 %. Of the five laboratories that is, observers had knowledge met the performance criteria for control survival, three laboratories produced a monotonic dose response in reproduction. The percentage of change in reproduction (offspring/survivor) between the control and the highest concentration of BRH sediment was on average 95 % for these three laboratories.

15.6.2.8 USEPA (2000 (206)) and Test Method E 1706 included a review of a series of round-robin studies from which interlaboratory precision was analyzed. CVs for survival in 10-d whole-sediment tests with *H. azteca* ranged from 6 to 114 % in three test sediments. Similar tests with *C. tentans* produced CVs of 8 to 181 % in three test sediments. In 28-d whole-sediment

tests with *H. azteca*, CVs from five test sediments ranged from 7 to 28 % for survival, from 52 to 78 % for growth (dry weight), and from 66 to 193 % for reproduction.

15.6.2.9 The *Leptocheirus* round-robin study exhibited similar or better intra- and interlaboratory precision than many chemical analyses and toxicity testing methods (for example, USEPA, 1991 (127)). The cause(s) of the high failure rate among laboratories participating in the round-robin study is not known. Several of the laboratories had not conducted this toxicity test previously, and inexperience with the procedures may have contributed to some of the test failures. Some of the laboratories suggested that uneaten food might have accumulated during early days of the experiment, which might have led to lethal low-DO stress to the young amphipods (DeWitt et al. 1997b (25)). Because of this potential problem, additional experiments were conducted (Annex A2) to find the minimum food ration that would minimize the build-up of excess food, minimize mortality, and produce significant growth rate and reproduction endpoints in the 28-d *L. plumulosus* sediment toxicity test. The diet recommended in this standard (A2.3.6) is based on the results of that experiment.

16. Interpretation of Results

16.1 The calculating procedure(s) and interpretation of the results should be appropriate to the experimental design. Procedures used to calculate results of toxicity tests can be divided into two categories: those that test hypotheses and those that provide point estimates. No procedure should be used without careful consideration of (a) the advantages and disadvantages of various alternative procedures and (b) appropriate preliminary tests, such as those for outliers and for heterogeneity. Preprocessing of data might be required to meet the assumptions of the analyses.

16.2 LC50 or EC50 and their 95 % confidence limits should be calculated on the basis of (a) the measured initial concentrations of test material, if available, or the calculated initial concentrations, and (b) the mortality or “effective mortality” (see 13.8.3). If other LCs or ECs are calculated, their 95 % confidence limits should also be calculated (see Guide E 729).

16.3 Most acute toxicity tests produce quantal data, that is, counts of the number of organisms in two mutually exclusive categories, such as alive or dead. A variety of methods (35) can be used to calculate an LC50 or EC50 and its 95 % confidence limits from a set of quantal data that is binomially distributed and contains two or more concentrations at which the percent dead or affected is between 0 and 100, but the most widely used are the probit, moving average, trimmed Spearman-Kärber and Litchfield-Wilcoxon methods (35). The method used should appropriately take into account the number of test chambers per treatment and the number of test organisms per chamber. The binomial test can usually be used to obtain statistically sound information about the LC50 or EC50 even when less than two concentrations kill or affect between 0 and 100 %. The binomial test does not provide a point estimate of the LC50 or EC50, but it does provide a range within which the LC50 or EC50 should lie.

16.4 The results of toxicity tests on field samples without replication may be reported in terms of survival values. A sample should be considered to be toxic if the single sample value lies outside the 95 % tolerance limits of the survival of the controls. Alternately, the field result may be compared with the control survival data using outlier detection methods; the sample may be considered toxic if it would be rejected as an extreme value when considered as part of the control population. Another approach is to use the special case comparison of a single value against a sample, described by Sokal and Rohlf (36). It is strongly recommended that samples be replicated if comparisons among sites are desired (see 12.3.2).

16.5 If samples from field stations are replicated, the mean survival at the stations and the mean control survival should be statistically compared by a one-tailed *t*-test or analysis of variance (ANOVA) followed by a multiple comparison test. Analysis of variance is used to determine whether any of the observed differences among the concentrations (or samples) are statistically significant. This is a test of the null hypothesis of no difference among concentrations (or samples). If the *F*-test is not statistically significant ($P > 0.05$), it can be concluded that the effects observed in the toxicant treatments (or field station samples) were not large enough to be detected as statistically significant by the experimental design and hypothesis test used.

16.5.1 Following a significant *F*-test result, all exposure concentration effects (or field station samples) can be compared with the control effects by using mean separation techniques such as those explained by Chew (37) orthogonal contrasts, Fisher’s methods, Dunnett’s procedure and William’s method. The Dunnett’s procedure is a multiple comparison test specifically designed to compare several experimental samples to the concurrent control (38). A multiple comparison test is a technique that accounts for the fact that several comparisons are being made simultaneously.

16.6 Daily observations on the numbers of amphipods that have completely or partially emerged from the sediment, either lying on the sediment surface, swimming in the water column, or floating at the water surface, can be used to document an apparent avoidance response to the sediment. Emergence data plotted against time can give the observer an impression of the degree of toxicity of the sediment during the course of the toxicity test, as amphipods often emerge earlier and in greater numbers from more highly toxic sediment.

17. Report

17.1 The record of the results of an acceptable sediment toxicity test should include the following information either directly or by reference to other available documents:

17.1.1 Names of test and investigator(s), name and location of laboratory, and dates of initiation and termination of the test.

17.1.2 Source of test material, lot number if applicable, composition (identifies and concentrations of major ingredients and impurities if known), known chemical and physical properties, and the identity and concentration(s) of any solvent used.

- ~~17.1.3 Source and method of preparation of water used, its salinity, and any other pertinent chemical characteristics.~~
- ~~17.1.4 Source of the control, reference and test sediments, dates and methods of collection, method of transport and storage of field sediments, method and dates of treatment of laboratory prepared sediment, and method of distribution to test chambers.~~
- ~~17.1.5 Source and date of collection of the test organisms, scientific name, name of person who identified the organisms and the taxonomic key used, age, life stage, means and ranges of weights and lengths, observed diseases or unusual appearance, treatments, holding and acclimation procedures.~~
- ~~17.1.6 Description of the experimental design, test chambers and covers, the depth, weight, and volume of sediment and water in the chambers, the date, time, and method of beginning the test, numbers of test organisms and chambers, temperature, salinity, and lighting regime.~~
- ~~17.1.7 The average and range of holding and test temperatures, and the method(s) of measuring or monitoring, or both.~~
- ~~17.1.8 Schedule for obtaining samples of sediment and water for geochemical analyses, and methods used to obtain, prepare, and store them.~~
- ~~17.1.9 Methods used for, and results (with standard deviations or confidence limits) of, chemical analyses of water quality and concentrations of test material, sediment geochemical analyses, and concentrations of test materials in sediment, including validation studies and reagent blanks.~~
- ~~17.1.10 Definition(s) of the effects used to calculate LC50s or EC50s and a summary of general observations of other effects.~~
- ~~17.1.11 A table of the biological data for each test chamber for each treatment (including the control(s)) in sufficient detail to allow independent statistical analyses.~~
- ~~17.1.12 The 10-day LC50s or EC50s and the methods used to calculate them, and their 95 % confidence limits, or the survival or mortality data and their significance relative to the control(s); specify whether results are based on measured or nominal concentrations of the test material.~~
- ~~17.1.13 Results of any other analyses that were made on the same sediment, such as faunal analyses, field notes made while collecting the sediment, chemical measurements made in test chambers, or chemical and geological analyses of subsamples of the sediment.~~
- ~~17.1.14 Anything unusual about the test, any deviation from these procedures, and any other relevant information.~~
- ~~17.2 Published reports should contain enough information to clearly identify the procedures used and the quality of the results.~~

18. Keywords

- ~~18.1 acute toxicity tests; *ampelisca*~~
- ~~16.1 *Ampelisca abdita*; amphipods; benthic amphipods; corophium; EC50 test; eohaustorius; estuarine environments; experimental design; exposure tests; grandidierella; hyalella; LC50 test; marine environments; *pontoporeia amphipod*; *bioavailability*; *chronic*; *Eohaustorius estuarius*; *reference toxicants*; *rhepoxynius estuarine*; *invertebrates*; *Leptocheirus plumulosus*; *saline water*; *saltwater*; *marine*; *Rhepoxynius abronius*; *sediment*; *sediment toxicity testing*; *static test*; *toxicity*; *toxicology*~~

ANNEXES

(Mandatory Information)

A1. RHEPOXYNIUS ABRONIUS

A1.1

A1. PROCEDURE FOR CONDUCTING A 10-D SEDIMENT SURVIVAL TEST WITH THE AMPHIPODS

Ecological Requirements—Ampelisca abdita, Eohaustorius estuarius, Leptocheirus plumulosus, or Rhepoxynius abronius (3)

A1.1 Introduction

A1.1.1 *Ampelisca abdita*, *Eohaustorius estuarius*, *Leptocheirus plumulosus*, and *Rhepoxynius abronius* occur along have been used extensively to test the West Coast toxicity of North America from central California to Puget Sound, Washington (39). It estuarine or marine sediments. The choice of these amphipod species as test organisms is based on sensitivity to sediment-associated contaminants, availability and ease of collection, tolerance of environmental conditions (for example, temperature, salinity, grain-size), ecological importance, and ease of handling in the desired test laboratory. Additionally, the species chosen for this method are intimately associated with sediment by nature of their burrowing or tube-dwelling and feeding habits. Field validation studies have shown that amphipods are absent or have reduced abundances at sites where toxicity is in laboratory tests. Amphipod sediment toxicity tests have been successfully performed for regulatory and research purposes by numerous laboratories, including state and federal government agencies, private corporations, and academic institutions. Test guidance for

A. abdita, *E. estuarius*, *L. plumulosus*, and *R. abronius* has been described (USEPA 1994a (1), Environment Canada 1992 (5)). The four species chosen are representative of both estuarine and marine habitats and sediments that span the spectrum of particle sizes from fine-to coarse-grained sediment. Thus, either alone or in combination, they may be used to measure toxicity of any commonly encountered estuarine or marine sediments (See Section 1 for additional details).

A1.1.2 Specific test methods for conducting the 10-d sediment toxicity test for the amphipods *Ampelisca abdita*, *Eohaustorius estuarius*, *Leptocheirus plumulosus*, and *Rhepoxynius abronius* are described in section A1.2. This test method was developed based on Swartz et al. (1985 (10)); DeWitt et al. (1989 (11)); Scott and Redmond (1989 (12)); Schlekert et al. (1992 (13)); and Environment Canada (1992 (5)). Results of tests using procedures different from the procedures described in section A1.2 may not be comparable and these different procedures may alter bioavailability. Comparison of results obtained using modified versions of these procedures might provide useful information concerning new concepts and procedures for conducting sediment-is 25-g/kg tests with estuarine or marine organisms. If tests are conducted with procedures different from the procedures described in this standard, additional tests are required to determine comparability of results (sections 1.4 and 1.10).

A1.2 Recommended Test Method for Conducting a 10-d Sediment Toxicity Test with *Ampelisca abdita*, *Eohaustorius estuarius*, *Leptocheirus plumulosus*, or *Rhepoxynius abronius*

A1.2.1 Recommended conditions for conducting a 10-d sediment toxicity test with *A. abdita*, *E. estuarius*, *L. plumulosus*, and *R. abronius* are summarized in Table A1.1. A general activity schedule is outlined in Table A1.2. Decisions concerning the various aspects of experimental design, such as the number of treatments, number of test chambers/treatment, and water quality characteristics should be based on the purpose of the test and the methods of data analysis (Section 13). The number of replicates and concentrations tested depends in part on the significance level selected and the type of statistical analysis. When variability remains constant, the sensitivity of a test increases as the number of replicates increase.

A1.2.2 The recommended 10-d sediment toxicity test with *A. abdita*, *E. estuarius*, *L. plumulosus*, and *R. abronius* is conducted at the species-specific temperature and salinity with a 24 h light photoperiod at a illuminance of about 500 to 1000 lux (Table A1.1). Test chambers are 1-L glass chambers containing 175 mL of sediment and 775 mL of overlying seawater. Twenty amphipods are added to each test chamber at the start of a test. The size range of the amphipods will depend on species that is being tested (see section 12.3.4 for allowable size range for each species). The number of replicates/treatment depends on the objective of the test. Five replicates are recommended for routine testing (Section 13). Exposure is static (that is, water is not renewed), and the animals are not fed over the 10-d exposure period. Overlying water can be culture water, surface water, site water, or reconstituted water. For site-specific evaluations, the characteristics of the overlying water should be as similar as possible to the site where sediment is collected. For all other applications, the characteristics of the overlying water for each species should be chosen according to Table A1.1. Requirements for test acceptability are summarized in Table A1.3).

A1.3 General Procedures

A1.3.1 *Sediment into Test Chambers* *Rhepoxynius abronius* naturally inhabits clean, fine, sandy sediments. In areas where:

A1.3.1.1 On the day before the addition of amphipods (Day -1), each test-sediments are predominantly silts sediment (either field collected or-elays, laboratory spiked) should be homogenized by stirring in the sediment storage container or by using a rolling mill, feed mixer, or other suitable apparatus. Control and reference sediments are included. Sediment should be visually inspected to judge the extent of homogeneity. If a silt-clay control treatment quantitative measure of clean sediment homogeneity is required, replicate subsamples should be taken from an uncontaminated reference collection site near that of the sediment batch and analyze for TOC, chemical concentrations, and particle size.

A1.3.1.2 A 175-mL aliquot of thoroughly homogenized sediment is added to each test-sediments, in addition chamber. It is important that an identical volume be added to each replicate test chamber; the native sediment control.

A1.2—*Collection and Handling Techniques*—*R. abronius* inhabits clean, fine, sandy sediments from the lower-intertidal volume added should equate to a depth of at least 274 m. Amphipods can 2 cm in the test chamber. The sediment added to the test chamber should be collected from a boat using a small biological dredge settled either by tapping the side of the test chamber against the side of the hand or a grab sampler. A sieve by smoothing the sediment surface with a 1.0-mm diameter mesh size can nylon, fluorocarbon, or polyethylene spatula. Highly contaminated sediment should be used added to separate adult *R. abronius* from their native sediment. Individual amphipods can be transferred between sorting trays, acclimation dishes, and test chambers in a certified laboratory fume hood.

A1.3.2 *Addition of Overlying Water*—As test water is added, disruption of the sediment surface should be minimized. One way to accomplish this is by using use of a bulb pipette turbulence reducer. Possible designs of turbulence reducer include a suitable size (for example, one with disk cut from polyethylene, nylon, or Teflon® sheeting (4 to 6 mil), or a 5-mm diameter opening).

A1.2.1 For acclimation, *R. abronius* can be counted into 10-cm diameter specimen dishes containing glass petri dish attached (open face up) to a 2-cm deep layer of 0.5-mm sieved collection site sediment, at glass pipette. If a disk is used as the turbulence reducer, it should fit the inside diameter of the test chamber and have attached a density length of 20 amphipods per dish. These dishes can be transferred nylon monofilament (or nontoxic equivalent) line. The turbulence reducer is positioned just above the sediment surface and raised as sea water is added to holding tanks supplied the 750-mL mark on the side of the test chamber. The turbulence reducer is removed and rinsed with aerated or flowing test sea water between replicates of a treatment. A separate

TABLE A1.1 Summary of Ecological and Testing Conditions that Should Be Considered When Conducting Ten-Day 10-d Sediment Toxicity Tests with *Ampelisca abdita*, *Eohaustoria estuarii*—See Annex for further estuarine, L explanation.

NOTE: 2—N/A = NOT APPLICABLE; N.D. = NO DATA AVAILABLE, or *Rhepoxynius t. abronimus* (USEPA 1994a (1))

A1a <i>Rhepoxynius abronius</i> (Family Phoxocephalidae)		
Field	Laboratory	
Parameter	Conditions	
Geographic range	Puget Sound to Southern California (23, 39)	N/A
Geographic range	1. Test type:	Whole sediment toxicity test, static.
Habitat	Free-burrowing sand dweller, low intertidal to 274 m (21, 23)	clean, finesand, 2 cm (3) Life-cycle
Habitat	2. Temperature:	15°C: <i>E. estuarii</i> Life-cycle
Life-stage tested	N/A	Mature 3 to 5-mm amphipods, mixed sexes (3)
Life-stage tested		20°C: <i>A. abdita</i>
Temperature	Annual range at collecting site = 8 to 16°C (3)	Standard temperature is 15°C; (3) survives 0 to at least 20°C
Temperature		25°C: <i>L. plumulosus</i>
Salinity	Annual range at collecting site = near 0 to 35 g/kg (40)	Standard salinity is 28 g/kg, salinity effects noted below 25 g/kg Well-sorted-f
Salinity	3. Salinity:	(3) Sediment type
Sediment depth	Usually upper 2 cm, to 6 cm (36, 39)	28‰ : <i>A. abdita</i> Sediment type
Sediment depth		Well-sorted-f
Light cycle	Natural light	Test sediment depth 2 cm (3) Nutrition
Light cycle		20‰ : <i>E. estuarii</i> Nutrition
Chronic-test	N/A	Continuous light (3) Control mortality
Chronic-test	4. Light quality:	Alternatively, the salinity of the overlying water can be adjusted to the salinity of the pore water at the site of interest for tests with <i>E. estuarii</i> or <i>L. plumulosus</i>
96 h LC50, cadmium, water-only exposure	N/A	Not developed
96 h LC50, cadmium, water-only exposure	5. Illuminance:	Wide-spectrum fluorescent lights
		0.92 (0.68–1.25) mg/L (42)
		500.68–1.25) mg/L (42)

A1b *Eohaustoria estuarii* (Family 1000 Ly Haustoriidae)

Field	Laboratory	
Geographic range	Central British Columbia to Central California (24, 27)	N/A
Geographic range	6. Photoperiod:	24 light
Habitat	Free-burrowing sand dweller, upper to mid-intertidal, shallow subtidal (24, 27)	Clean, fine sand, 2 cm (24)
Habitat	7. Test chamber:	1-L glass beaker or jar with 10-cm inner diameter.
Life-cycle	Probably annual (24)	N/A—Field collected
Life-cycle	8. Sediment volume:	175 mL (about 2-cm depth)
Life-stage tested	N/A	Mature amphipods 3–5 mm, mixed sexes (24)
Life-stage tested	9. Overlying water volume:	775 mL
Temperature	Approximately 0 to 21°C (24, 27)	Standard temperature is 15°C (3, 24); tolerates 5 to at least 21°C (27)
Temperature	10. Renewal of overlying water:	None
Salinity	Annual range at collecting site = near 0 to 35 g/kg	Standard salinity is 2 to ≤28 g/kg (24, 27)
Salinity	11. Size and life stage of amphipods:	<i>A. abdita</i> : 3 to 5 mm (no mature males or females)
Sediment type	Clean fine to medium sand (27, 43)	92% mature survival in sediments with ≥80% silt-clay, 97% in sandy sediments (25)
Sediment type		<i>E. estuarii</i> : 3 to 5 mm
Sediment depth	Approximately top 5 cm	2 cm (3, 24)
Sediment depth		<i>L. plumulosus</i> : 2 to 4 mm (no mature males or females)
Nutrition	Probable deposit feeder (24)	Amphipods are not fed in the laboratory (3, 24)
Nutrition		<i>R. abronius</i> : 3 to 5 mm
Light cycle	Natural light	Continuous light (3, 24)
Light cycle	12. Number of organisms/chamber:	20 per test chamber
Control mortality	N/A	≤10% (3, 24)
Control mortality	13. Number of replicate chambers/treatment:	Depends on objectives of test. At a minimum, four replicates should be used.
Chronic-test	N/A	Not developed
Chronic-test	14. Feeding:	None
96 h LC50, cadmium, water-only exposure	N/A	9.33 (7.20–12.00) mg/L (24)
96 h LC50, cadmium, water-only exposure	15. Aeration:	9.33 (7.20–12.00) mg/L (24)

A4 Water in each test chamber should be aerated overnight before set up of test, and throughout the test; aeration at rate that maintains >90% saturation of dissolved oxygen concentration.

Field	Laboratory	
Geographic range	Central Main to Northern Florida, eastern Gulf of Mexico (44), San Francisco Bay (45)	N/A
Geographic range	16. Overlying water:	Clean sea water, natural or reconstituted water.
Habitat	Infaunal tube dweller, low intertidal to 60 m (44, 46)	Collection site sediment, 4 cm (33)
Habitat	17. Overlying water quality:	Temperature daily, pH, ammonia, salinity, and DO of overlying water at least at test start and end. Salinity, ammonia, and pH of pore water.
Life-cycle	Two to several generations per year, temperature dependent, probably one brood per female (46)	Life cycle approximately 6 weeks at 20°C (48)
Life-cycle	18. Test duration:	10 d
Life-stage tested	N/A	Immature amphipods, or mature females only
Life-stage tested	19. Endpoints:	Survival (reburial only)
Temperature	Collected in water temperatures from –2 to 27°C (44) Standard temperature is 20°C, has been tested in 4	

TABLE A1.2 General Activity Schedule for Conducting a Sediment Toxicity Test with *Ampelisca abdita*, *Eohaustorius estuarius*, *Leptocheirus plumulosus*, or *Rhepoxynius abronius* (USEPA 1994a (1))

TABLE A1.1—Continued		
A1d— <i>Grandidierella japonica</i> (Family Corophiidae)		
Field	Laboratory	
Day	Activity	
Geographic range -10 to -3	Japan, San Francisco Bay (50), Southern California bays (51)	N/A
Habitat -9 to -2	Intertidal to subtidal (51)	Collection site sediment, 1 cm (51)
Life-cycle -1	N.D.	4 to 5 life cycles per year at 20°C
Life stage tested 0	N/A	4 Observe amphipods, monitor water quality. Add sediment to each test chamber, place chambers into exposure system, and start aeration.
Temperature 1	Collected in California from water temperatures ranging from 9 to 26°C (51)	Standards test temperatures 15 to 19°C; satisfactory survival at 15 to 23°C (51)
Salinity 2	Full ocean salinity to hypersaline waters of unknown salinity (51)	Standards test salinity is 30 to 35 g/kg; survival at 16 to 34 g/kg, 15% mortality at 4 g/kg (51)
Sediment type 3 to 7 and 9	Mud sand; occurs in sands, silts, clay (50, 52)	Standards test silty clay (51)
Sediment depth 8	Sand; occurs in sands, silts, clay (50, 52)	Standards test silty clay (51)
Nutrition 10	Upper 2–4 cm (51)	Standards test 2 cm (51)
Light cycle Natural light	Same as Day 2–4 cm (51)	Standards test 2 cm (51)
Light test by collect light	Algae, detritus, sediment	Suspension of finely ground Tetramin and <i>En teromorpha</i> (51)
Control mortality	Measure ment	Superature. End the (51)
Control mortality	Continuous light (51)	
Chronic test	Control mortality	≤40% (51)
Competition test	Control mortality	≤40% (51)
96h LC50, c admium, water only exposure	Control mortality	Not developed
96s with a sieve exposure	Control mortality	Not developed
		4.17 (0.94–1.46) mg/L (53)
		±

turbulence reducer is used for each treatment. The test chambers should be covered, placed in a temperature controlled water bath (or other acceptable equivalent) and gently aerated. A test begins when the organisms are added to three test chambers (Day 0).

A1.3.3 Addition of Amphipods—On the following day (Day 0), amphipods are added to the test conditions. A sieve with a 1.0-mm diameter mesh size can be used to separate *R. abronius* from the acclimation culture or control sediment immediately prior in the holding container(s), and transferred to a sorting tray. The additional animals allow for the selection of healthy, active individuals. The sieve size for isolating amphipods from the culture or control sediment will depend upon the selected species. *Ampelisca abdita* and *L. plumulosus* should be isolated using a toxicity test.

A1.3 0.5-mm sieve, whereas Toxicity Test Specifications—The toxicity test *E. estuarius* and *R. abronius* should be run at 15 ± 3°C isolated using 28 g/kg overlying water in the test chambers. The test chamber is usually a standard 1-L glass beaker with a 10-cm internal diameter. Beakers should be covered with an 11.4-cm diameter watch glass to reduce contamination conducted with sea water of the same temperature and evaporation of salinity as the water holding and test material. Aeration can be provided to each test chamber through an impartially selected using a 1-mL glass transfer pipette that extends between the beaker spout or other suitable tool (not forceps), and distributed among dishes or cups containing test sea water until each container has twenty amphipods. The number of amphipods in each dish should be verified by recounting before adding to a depth not closer than 2 cm from the sediment surface. Sediment test chambers. To facilitate recounting, amphipods may be distributed to test chambers in batches of 5 or 10 instead of the full complement of 20. The distribution of amphipods to the test chambers should be 2-cm deep, and toxicity test water impartial.

A1.3.3.1 Amphipods should be added up to test chambers without disruption of the 700-mL mark on sediment. Any amphipods remaining in the beakers. Sediment and sorting container should be gently washed into the test chamber using test sea water. The

TABLE A1.3 Test Acceptability Requirements for a 10-d Sediment Toxicity Test with *Ampelisca abdita*, *Eohaustorius estuarius*, *Leptocheirus plumulosus*, or *Rhepoxynius abronius* (USEPA 1994a (1))

TABLE A1.3e <i>Leptocheirus plumulosus</i> (Family Aoridae)	
A. It is recommended for conducting a 10-d test with <i>A. abdita</i> , <i>E. estuarius</i> , <i>L. plumulosus</i> , or <i>R. abronius</i> that the following performance criteria are met:	
Field Laboratory	
1. Size, life stage, and reproductive stage of amphipods must be within the prescribed species-specific ranges at the end of the test (Section 12).	
Geographic range	Cape Cod, Massachusetts to Northern Florida (54)
Geographic range	2. Average survival of amphipods in the control sediment must be greater than or equal to (54)
N/A	
3. Salinity, pH, and ammonia in the overlying water and sediment grain size should be within tolerance limits of test species.	
Habitat	U-shaped burrows in fine sand to muddy sediments; shallow subtidal (54, 55)
Collection site or culture sediment, 2 cm (56, 59-62)	
B. Performance-based criteria for culturing <i>L. plumulosus</i> include:	
Life cycle	Annual; reproduction spring through fall in Chesapeake Bay, at least two broods per female (54, 56).
Life cycle	1. Laboratories should perform periodic 96-h water-only reference-toxicity tests (at a minimum, one test every six months) to assess the sensitivity of culture organisms (section 11.16).
Multiple broods per year, lifespan \geq 7 weeks (62)	
2. Records should be kept on the frequency of restarting cultures.	
Life stage tested	N/A
Immature or mature 3 to 5 mm amphipods; mixed sexes (56, 59-62)	
4. Laboratories should characterize and monitor background contamination and nutrient quality of food if problems are observed in culturing or testing organisms.	
Temperature	Range at collecting site = 0 to 29°C (57)
Routinely tested at 20°C, has been tested at 25°C (56, 59-61)	
C. Performance-based criteria for field-collected amphipods include:	
Salinity	Collected in water ranging from near 0 to 33 g/kg (55, 58).
Salinity	1. Laboratories should perform reference-toxicant tests on each batch of field-collected amphipods received used in a sediment test (section 11.16).
\geq 90% mean survival in salinities 2 to 32 g/kg (56, 59-61) ∞	
2. Acclimation rates to test salinity and temperature should not exceed 3°C and 5‰ per 24 h.	
Sediment type	Fine sand to silty clay (55-58)
Sediment type	3. Amphipods used in a toxicity test should exhibit active swimming behavior upon placement in water, have full digestive tracts, and display (55-58)
Up to 100% survival with $>$ 90% silt-clay; 85% mean survival with $>$ 95% sand-gravel (56, 59-61)	
Sediment depth	Usually in upper 2 cm; rarely deeper than 5 cm
Sediment depth	D. Additional requirements - 5 cm
2 cm (56, 59-61)	
1. All organisms in a test must be from the same source.	
Nutrition	Surface deposit and suspension-feeder (57)
Nutrition	2. Storage of sediments collected from the field should follow guidance outlined in section-feeder (57)
Combination of "amphipod group" and micro-algae (62)	
3. All test chambers (and compartments) should be identical and should contain the same amount of sediment and overlying water.	
Light cycle	Natural light 16h:8h light:dark (56, 59-61)
Light cycle	4. Negative-control sediment and appropriate solvent controls must be included in a test. The solvent control used must not adversely affect test organisms.
Control mortality	N/A
Control mortality	N/A
Chronic test	N/A
Chronic test	N/A
96h LC50, cadmium water only exposure	N/A
96h LC50, cadmium water only exposure	N/A

water level should be added brought up to beakers the day before 950-mL mark, the amphipods are added, test chamber covered, and aeration may be discontinued for up to allow suspended sediment particles 1 to settle, and 2 h to allow the amphipods to burrow into the sediment. Aeration should then continue for equilibration the remainder of temperature and the sediment-water interface.

A1.3.1 After test.

A1.3.3.2 After the overnight equilibration time, 20 amphipods are distributed to each addition of the animals, the test chambers, with additional toxicity test water to bring chambers should be examined for animals that may have been injured or stressed during the water level up to the 950-mL level: isolation, counting, or addition processes. Injured or stressed animals will not burrow into sediments, and should be removed. The period of time allowed for healthy amphipods to bury into test sediments will depend upon the species used. *Eohaustorius estuarius*, *L. plumulosus*, and *R. abronius* should be allowed 5 to 10 min to bury into the test substrate. Any amphipods sediment. *Ampelisca abdita*, which may take longer to build tubes, should be allowed 1 h. Amphipods that have not burrowed within that the prescribed time or appear damaged should be replaced, unless replaced with animals from the amphipods same sieved population, unless they are repeatedly burrowing into the sediment and immediately emerging in an apparent avoidance response to the test substrate. response. In that case, the amphipods are not replaced. Amphipods are not removed from the surface The number of test sediments during the course of the test even if they appear dead, since some amphipods that seem dead might actually are removed should be alive and might later rebury recorded.

A1.3.4 Test Conditions:

A1.3.4.1 Aeration—The overlying sea water in the each test sediments:

A1.3.2 The toxicity test chamber should be aerated continuously after the water is terminated when amphipods are separated from added (that is, Days -1 through 10) except during introduction of the test-substrates using a 0.5-mm mesh-diameter screen. Amphipods are transferred to a sorting tray organisms. Compressed air, previously filtered and numbers free of oil, should be bubbled through a glass or plastic pipette and dead amphipods are counted. Survivors are transferred attached plastic tubing. The tip of the pipette should be suspended 2 to dishes containing a 2-cm deep layer 3 cm above the surface of clean, native the sediment and allowed 1 h layer so as to rebury: not disturb the sediment surface. The numbers concentration of survivors unable to rebury dissolved oxygen (DO) in clean the water overlying the sediment can be used to calculate an EC50 for this sublethal effect.

A1.4 Life Cycle and Age Class—*Rhepoxynius abronius* has an annual life cycle (40), with recruitment occurring primarily in the 1 test chambers is maintained at or near saturation by gently aerating the water. Air is bubbled through the spring months. Large immature and adult amphipods, 3 test chamber at a rate that maintains a >90 % DO concentration, but does not cause turbulence or disturb the sediment surface. If air flow to 5 mm total length, one or more test chambers is interrupted for more than one hour, DO should be measured in the toxicity those test because they are available year round, and their sensitivity chambers to contaminated sediments has been shown to be not greatly different from that determine whether DO concentrations have fallen below 60 % of juveniles (19). They are also large enough to saturation. Results may be easily handled and counted unacceptable for test chambers in the toxicity test. Mature males and females, even those carrying eggs, have been found which aeration was interrupted and DO concentrations fell to be equally sensitive to test materials, so it is possible to use a mixed population of both sexes, although very large mature individuals below 60 % saturation.

A1.3.4.2 Lighting—Lights should not be used because they might be senescent. It is necessary left on continuously at an intensity of 500 to change year classes sometime 1000 lux during the summer, as old amphipods die out and are replaced by 10-d exposure period. The constant light increases the maturing juveniles.

A1.5 Control Survival—Control survival using *Rhepoxynius* is generally 95 % or greater, and must be at least 90 % for the toxicity test to be considered valid.

A1.6 Sensitivity—*Rhepoxynius abronius* has been shown to be among the most sensitive tendency of sediment toxicity test the organisms to test materials, but is fairly tolerant of handling remain buried in the sediment, and thus to a variety of physical characteristics of sediment (3). The genus *Rhepoxynius* is one of the first remain exposed to disappear from benthic communities impacted by pollution (3, 39).

A1.7 Interpretation and Interferences—In interpreting the data from 10-day sediment toxicity tests with adult *Rhepoxynius*, it should be kept in mind that the very early life stages, the reproductive ability test material.

A1.3.4.3 Feeding—The four species of amphipods, or their long-term survival might be affected by contaminants at lower concentrations than those that produce a lethal or sublethal effect in mature amphipods used in a short-term test. *Rhepoxynius* has been shown to this method are not be somewhat adversely affected by very fine-grained sediments (20). Despite these limitations, fed during the toxicity 10-d exposure period.

A1.3.4.4 Water Temperature—The test using *Rhepoxynius abronius* has been demonstrated temperature will depend on the species that is tested. Test temperatures were selected to be very useful in detecting sediment toxicity, and can near the summertime thermal maximum that each species would be used expected to encounter in a variety of research and regulatory applications.

A2. *EOHAUSTORIUS ESTUARIUS*

A2.1— the environment. *Ecological Requirements*— *Eohaustorius estuarius* (24) and lives in intertidal sands along *R. abronius*, the North American west coast from British Columbia south to Pacific Coast amphipods, should be tested at least central California 15°C. *Ampelisca abdita* (26, 43). It is a desirable test species for sediments which have interstitial salinities ranging between 2 should be tested at 20°C and 28 g/kg. Since *E. estuarius*—*L. plumulosus* normally inhabits sandy sediments, at 25°C.

A1.3.4.5 *Salinity*—The salinity of the experimental design should include a fine sediment control (in addition to water overlying the native sediment control) if test sediments are predominantly silts or clays. This control sediment should consist of clean sediment from an uncontaminated reference collection site near that of will vary depending on the selected test sediment(s) and have a similar grain size distribution.

A2.2— species. For routine testing, *Collection and Handling Techniques* *A. abdita* —*E. estuarius* and *R. abronius* can be found in the upper 10 cm of fine, intertidal, estuarine sands +0.5 to +2.0 m above mean low low water (MLLW). The amphipods can be collected by shovel tested at low tide and sieved from their native sediments with a 1.0-mm mesh diameter screen. They can be transferred between sorting trays, acclimation dishes, and test chambers with a 5-mm diameter bulb pipette.

A2.2.1 For acclimation, up to 20 an overlying water salinity of 28 ‰, whereas *E. estuarius* can be held in 10-cm diameter specimen dishes containing a 2-cm deep layer of native sediment served to ≤0.5 mm. These dishes should be transferred to holding tanks supplied with aerated or flowing sea water tested at the 20 ‰. The target test temperature and salinity. Two to three days are sufficient salinity for acclimation to test conditions. The amphipods should be separated from the acclimation sediments using a 1.0-mm mesh diameter sieve immediately prior to initiating the toxicity test.

A2.3— *Toxicity Test Specifications*—The toxicity test should be run at 15 ± 3°C with *L. plumulosus* is 5 or 20 ‰ depending on the pore-water salinity. The recommended overlying water composed of toxicity test seawater diluted to the same salinity as the interstitial water of the test substrate. The test chamber is usually a standard 1-L glass beaker with a 10-cm internal diameter. Beakers should be covered with an 11.4-cm diameter watch glass to reduce contamination of the contents and evaporation of the water and 5 ‰ for test material. Aeration can be provided to each test chamber through a 1-mL glass pipette that extends between the beaker spout and the watchglass cover to a depth not closer than 2 cm from the with sediment surface. Sediment in the test chambers should be 2 cm deep, and toxicity test pore water should be added up from 1 to the 700-mL mark on the beakers. Sediment and water should be added to beakers the day before the amphipods are added, to allow suspended sediment particles to settle, and to allow time 10 ‰ or 20 ‰ for equilibration of temperature and the sediment-water interface.

A2.3.1 After the overnight equilibration time, 20 amphipods are distributed to each of the test chambers, sediments with additional toxicity test pore water to bring >10 ‰. Alternatively, the water level up to the 950-mL level. The amphipods should be allowed 5 to 10 min to bury into the test substrate. Any amphipods that have not buried within that time should be replaced, unless the amphipods are repeatedly burrowing into the sediment and immediately emerging in an apparent avoidance response to the test substrate. In that case, amphipods are not replaced. Amphipods are not removed from the surface salinity of test sediments during the course of the toxicity test even if they appear dead, since some amphipods that seem dead might actually be alive and might later rebury into the test substrate.

A2.3.2 The toxicity test terminates when the amphipods are sieved from the test substrate using a 0.5-mm mesh diameter screen and the animals are transferred to a sorting tray. After survivors are counted, the ability of surviving amphipods to rebury into clean native sediments may be used to determine an EC50 overlying water for this sublethal effect. Surviving toxicity tests with *E. estuarius* should or *L. plumulosus* can be transferred adjusted to specimen dishes containing a 2-cm deep layer of native sediment sieved to ≤0.5 mm, and should be allowed selected target salinity (for example, one hour to rebury).

A2.4— representative of the salinity regime at the site of interest). Depending on the objectives of the study, *Life Cycle and Age Classes*—*Eohaustorius* *E. estuarius* and *L. plumulosus* can be tested with overlying water salinity ranging from 1 to have an annual life cycle, 32 ‰. However, if tests are conducted with reproduction occurring procedures different from February through July (27). Large immature and adult amphipods, 3 those described in Table A1.1 (for example, different salinity, lighting, temperature, feeding conditions), additional tests are required to 5-mm total length, determine comparability of results (sections 1.4 and 1.10). Pore-water salinity of each test sediment should be used in measured before the toxicity test because they are available year round and are easily handled and counted. Larger individuals initiation of a test. Sediment pore water should not be used as they might be senescent. Age, size, and sex-specific sensitivity obtained by centrifugation. Alternatively, salinity can be measured before homogenization in the water that comes to the surface in the sample container as the sediment settles. The pore-water salinity of *E. estuarius* to contaminants has not been examined, but mixed-sex populations of animals the test sediment should be within their recommended size the salinity application range show highly replicable responses to laboratory-spiked and field-collected contaminated sediments of the chosen amphipod species (Table A1.4). (24). This strongly suggests that both sexes *Rhepoxynius abronius* cannot be tested when sediment pore-water salinities are comparably susceptible to contaminated sediments.

A2.5— <25 ‰. Another species should be used for such sediments. *Control Survival*—Control survival using *Ampelisca abdita*,

TABLE A1.4 Limits on Environmental Conditions Under Which to Conduct the 10-d Sediment Toxicity Tests for 10-d Sediment Toxicity Tests with *Ampelisca abdita*, *Eohaustorius estuarius*, *Leptocheirus plumulosus*, or *Rhepoxynius abronius* (USEPA 1994a (1))

Parameter	<i>Ampelisca abdita</i>	<i>Eohaustorius estuarius</i>	<i>Leptocheirus plumulosus</i>	<i>Rhepoxynius abronius</i>
Temperature (°C)	20	15	25	15
Overlying water salinity (‰)	>10	1 to 35	1.5 to 32	>25
Grain size (% silt/clay)	>10	<70 clay ^A	Full range	<90
Ammonia (total mg/L, pH 7.7)	<30	<60	<60	<30
Ammonia (unionized, mg/L, pH 7.7)	<0.4	<0.8	<0.8	<0.4

^A Environment Canada (1998 (221))

Eohaustorius E. estuarius, and *L. plumulosus* must be tested over the entire pore-water salinity range (that is, 1 to be considered valid.

A2.6– 34 ‰) when the recommended species-specific overlying salinity is used. PSEP (1995 *Sensitivity—Eohaustorius estuarius* (192) *is only slightly less sensitive than Rhepoxynius to contaminants, and is fairly tolerant*) outlines a procedure for adjusting the pore-water salinity of samples with a salinity <25 ‰ by adding appropriately saline overlying water to the test chamber on the day before the start of the test, mixing sediment and overlying water, and allowing the material to settle overnight under aeration. The species resultant overlying water is less sensitive than either retained or about 75 % replaced with fresh dilution water at 28 ‰ (PSEP 1995 *Rhepoxynius* (192)). While this manipulation should result in an acceptable salinity for tests with *R. abronius*, the influence of physical characteristics this manipulation on the bioavailability of contaminants in the sediment and sample is tolerant of salinity levels ranging from about 2 g/kg to uncertain. See section 12.1.4.

A1.3.5 Measurements and Observations :

A1.3.5.1 Temperature should be measured at least 35 g/kg.

A2.7 *Interpretation*—When interpreting daily in at least one test chamber from each treatment The temperature of the water bath or the exposure chamber should be continuously monitored. The time-weighted average of daily temperature readings must be within $\pm 2^{\circ}\text{C}$ of the desired temperature. The instantaneous temperature must always be within $\pm 3^{\circ}\text{C}$ of the desired temperature.

A1.3.5.2 Salinity, DO, and pH of the overlying water should be kept measured daily in at least one test chamber per treatment. Care should be taken not to disturb the sediment when sampling overlying water quality.

A1.3.5.3 Ammonia should be measured in overlying water towards; the reproductive ability, or beginning (for example, Day 2) and towards the long-term survival end of *E. estuarius* might the test (for example, Day 8). Measurement of overlying water pH and temperature should accompany each ammonia measurement. Simultaneous measurements of ammonia, pH, and temperature in sediment pore water should be measured at the beginning of the test. Pore water should be extracted after the sediment has been press-sieved and homogenized. Samples of pore water should be obtained by centrifugation.

A1.3.5.4 Each test chamber should be examined at concentrations lower than those that produce a lethal or sublethal response. Despite these limitations, least daily during the toxicity 10-d test using adult *E. estuarius* has been demonstrated period to be useful in quickly detecting ensure that airflow to the overlying sea water is acceptable. The number of amphipods emerged from the sediment toxicity including those swimming in estuarine sediments of widely varying interstitial salinity; the water column and can trapped in the air-water interface should be used noted. Amphipods caught in the air-water interface should be gently pushed down into the water using a glass rod or pipette.

A1.3.6 Ending a Test:

A1.3.6.1 Laboratories should demonstrate the ability of their personnel to recover an average of at least 90 % of the organisms from control sediment. For example, test organisms could be added to control sediment and regulatory applications:

A3. AMPELISCA ABDITA

A3.1– recovery could be determined after 1 h (Tomasovic et al., 1994 *Ecology—Ampelisca abdita* is a tube-dwelling amphipod belonging (207)).

A1.3.6.2 The contents of the test chambers are sieved to isolate the family Ampeliseidae, found mainly in protected areas test animals. The mesh size for sieving the contents of the test chambers should be no larger than 0.5 mm. Test water should be used for sieving. Material retained on the sieve should be washed into a sorting tray with clean test sea water. *Ampelisca abdita* are tube-builders and can be forced from their tubes for enumeration by slapping the low intertidal zone to depths sieve forceably against the surface of 60 m. It ranges the water or by using a stream of water (for example, from central Maine a spray bottle or from a tube connected to south-central Florida and the eastern Gulf a source of Mexico running water). (44, 63) *Eohaustorius estuarius*, and has also been introduced into San Francisco Bay (45). It is euryhaline, *L. plumulosus*, and has been reported in waters that range from fully marine to 10 parts per thousand salinity (44). This species generally inhabits sediments R.

abronius are easily removed from fine sand to mud and silt without shell, although it can also the sediment by the sieving process. The sieve should be carefully examined for any animals remaining, and in relatively coarser sediments with a sizable fine component (46). the case of *A. abdita* is often abundant in sediments with a high organic content (47).

A3.1.1 In the colder waters of its range, *A. abdita* produces two generations per year, an overwintering generation, also for any tubes (which may contain animals) remaining.

A1.3.6.3 Material that breeds in has been washed from the spring and a second that reproduces in mid to late summer (45, 46). In New England, breeding of sieve into the overwintering generation begins when sorting tray should be carefully examined for the water temperature is about 8°C, but in warmer waters south presence of Cape Hatteras, breeding might be continuous throughout amphipods. A small portion of the year. Adults mate in the water column, and intense breeding activity is correlated with the full moon and spring tides. Juveniles are released after approximately two weeks in the brood pouch, material should be sorted through at about 1.5 mm in length. It then takes 40 to 80 days for newly released juveniles to become breeding adults (46). When *A. abdita* are present, a time, removing amphipods as they are often dominant members of the benthic community found. Material from tests conducted with densities up to 110 000 m⁻² (45, 47, 64). *Ampelisca A. abdita* is a particle feeder, feeding both on particles in suspension and on those from the surface of the sediment surrounding its tube. Gut contents of field-collected specimens have been found to include algal material, sediment grains, and organic detritus (44, 45).

A3.2 Collection and Handling Techniques—*Ampelisca* should be sieved from their native (collection site) sediment as soon as possible after collection. A 2-mm mesh sieve nesting over a 0.5-mm mesh sieve is useful for this procedure. It is desirable for will include tubes built by the sediment containing the amphipods to be rinsed first through during the upper, 2-mm sieve with a forceful stream of seawater at the collection temperature test. The tubes should be carefully examined and salinity. This teased apart under a dissecting microscope or magnifying glass because *A. abdita* will break up often remain in the sediment material and also force most tubes even after vigorous sieving. Numbers of the live, missing, and dead amphipods out of their tubes. The material thus retained on the 0.5-mm sieve should be vigorously shaken determined and swirled so the fine sediments pass through and the amphipods recorded for each test chamber. Missing animals are separated from tubes, sediment, and detrital material. If the sieve is then lifted from the water, allowed assumed to drain, have died and then slowly lowered into a shallow tray of seawater, decomposed during the *Ampelisca* will be caught on the water's surface tension test and can disintegrated; they should be easily collected with a fine mesh dip net. The amphipods can be held temporarily included in large culture dishes in a constant temperature bath, and then separated into two size classes with the use of nested 1.0 and 0.5-mm sieves.

A3.2.1 During acclimation, *Ampelisca* can be held number dead in 1-gal glass jars, each containing approximately a 4-cm deep layer calculations of sieved collection site sediment. If seawater is flowing through the holding containers, a screened overflow must percent survival for each replicate treatment. Amphipods that are inactive but not obviously dead should be observed using a low-power dissecting microscope or a hand-held magnifying glass. Any animal that fails to prevent loss exhibit movement (that is, neuromuscular twitch of swimming amphipods. Amphipods should have food available on pleopods or antennae) upon gentle prodding with a daily basis during acclimation. Research probe should be considered dead.

A1.3.7 Test Data—Survival is currently being conducted to determine optimal food sources for culturing this amphipod. Reasonable growth and reproduction have been obtained when the primary endpoint recorded at the end of the 10-d sediment toxicity test with *A. abdita* has been fed the diatom, *Phaeodactylum tricornerutum* *E. estuarius* daily in excess (a suggested amount is 0.5 to 1 L of algae per gallon jar, or 3×10^{-4} cells/mL), *Skeletonema costatum* L. *plumulosus*, and has also been used successfully. Amphipod exposure *R. abronius*. The ability of surviving amphipods to the food source will be increased if, during the feeding period (for example, overnight), the holding system is static, with aeration to circulate the algae. Sloping upper sides on the holding containers will aid rebury in movement of algae across the clean control sediment surface. Care should can be taken used to calculate effective mortality, that ins, the temperature with a water bath when seawater is not flowing through the jars. Approximate density in the holding jars should not exceed 300 amphipods. Acclimation sum of dead animals plus those survivors that fail to the test temperature should not exceed 3°C per day, rebury. This endpoint has been used for *E. estuarius*, *L. plumulosus*, and *R. abronius*. If it is desired to determine rebury, surviving amphipods should be used within 2 weeks after collection.

A3.2.2 *Ampelisca abdita* may be shipped if this is done within one day of collection. Small plastic “sandwich” containers (approximately 500 mL) can be used transferred to hold the amphipods. The containers are filled three quarters full with holding a minimum depth 2-cm layer of 2 cm of 0.5-mm sieved collection site control sediment and an overlying layer (22 cm) of test sea water. Salinity of the top with well-aerated seawater. No more than 200 amphipods test sea water for rebury should be added to each container. Amphipods should be allowed to burrow into the sediment and build tubes before the containers are capped. The capping must be done underwater to eliminate any air pockets same as that measured in the containers. Containers should be shipped by means test chamber. The number of overnight delivery in coolers with a few ice packs surviving amphipods unable to prevent extreme temperature changes during transit.

A3.3 Ten-Day Sediment Toxicity Test—The variation of *Ampelisca*'s sensitivity to toxic materials under different physical conditions rebury in control sediment after 1 h is still being examined. This species is routinely tested at 20°C, but has been tested from 8 to 25°C. In nature, feeding and somatic growth occur at temperatures as low as 3 to 5°C (44). For comparison with other *Ampelisca abdita* recorded for each test results, 20°C chamber and is recommended. Similarly, *A. abdita* is tolerant used to calculate effective mortality.

A1.4 Interpretation of a wide salinity range, but most tests have been conducted at salinities Results

A1.4.1 Section 13.3 describes general information for interpretation of 28 to 35 g/kg. This amphipod inhabits fine-grained sediments, and as with other physical conditions, if it is suspected test results. The following sections describe species-specific information that a coarse grain size is useful in helping to interpret the results of a test sediment will stress the animals, a grain size control should be included.

A3.3.1 The exposure chamber routinely used to test toxicity tests with *A. abdita*, is a quart-sized glass canning jar with a narrow mouth. This container was selected because it is inexpensive, easily available, easily drilled if a screened overflow is needed for flow-through tests; *E. estuarius*, *L. plumulosus*, and has sloping upper sides to improve circulation *R. abronius*.

A1.4.2 Influence of algal material in experiments where growth or reproductive endpoints are measured and feeding is necessary: Indigenous Organisms *Ampelisca abdita* has not been tested—Indigenous organisms may be present in the 1-L beaker exposure chamber used field-collected sediments. An abundance in other amphipod tests, but it is not anticipated that use of beakers would create any problems. With either exposure chamber, the water column should be gently aerated with a glass pipette inserted above the sediment surface. Sediment in the exposure chamber should be 4 cm deep.

A3.3.2 *A. abdita* can be collected throughout the year. However, during certain times sample of the year, juvenile amphipods might be difficult to obtain. If mature animals are used, adult males must not be tested; they are very active swimmers and they die shortly after mating. test organism (McGee et al. 1999 *Ampelisca* should be sieved from the holding containers using a 0.5-mm sieve. Twenty(41)), or organisms taxonomically similar to thirty amphipods should be tested per replicate. For each replicate, the contents test organism, may make interpretation of a sorting cup treatment effects difficult. The presence of predatory organisms can be rinsed into a plastic cup with a 400 or 500-micron screened base also adversely affect test organism survival. For example, Redmond and from there into Scott (1989 (104)) showed that the exposure container. Any animals caught on the water's surface polychaete *Nephtys incisa* can be gently pushed consume *Ampelisca abdita* under using a glass rod. Amphipods should be given 1 h to burrow into the sediment. If the lack toxicity test conditions.

A1.4.3 Effect of ability to burrow does not Sediment Grain Size —All four species show a dose-response, then the animals not burrowed can be replaced tolerance to most sediment types, with others from the same sieved population.

A3.3.3 The endpoint for the 10-day test is mortality, generally little effect on survival whether coarse-grained or fine-grained (that is, predominantly silt and dead animals should be counted and removed daily. An amphipod is considered dead if it does not respond clay) clean sediments are used. However, adverse effects due to gentle probing. It is also useful to note any animals out of their tubes on the grain-size distribution of test sediment or water surface, amphipods may occur when sediments that are nearly dead and only exhibit a muscular pleopod twitch, either extremely sandy or fine depending on the presence species of molts, and the condition amphipod used. In order to separate effects of the tubes built. Emergence sediment-associated contaminants from the sediment effects of particle size, an appropriate clean control and reference sediment should be incorporated into the inability to construct a proper tube test when test sediments are sublethal behavioral responses that would ultimately result in death.

A3.3.4 After checking within the assay on the last day, the contents range of each exposure container should be rinsed through a 0.5-mm sieve. (A smaller mesh sieve can be used for the final sieving if there is concern about losing very small animals, but this will make the sieving process more time-consuming.) If the experiment is small, the material retained on the sieve can be examined that day. If time does not permit same-day examination, the retained material from each jar can be preserved in 5 % buffered formalin with Rose Bengal stain for later examination. Any amphipods each species. Alternatively, another species that are not accounted for when the sieved material is examined are presumed to have died during tolerant of the test. Amphipods that have died sediment extreme in their tubes will generally decompose during the test or break apart during sieving. Rarely, an individual that has died during the test will question might be recovered in tested o conjunction with the preserved material, and its appearance will be markedly different from those chosen species. Ranges of the amphipods that were alive when preserved. For instance, there might be little tissue left within its exoskeleton, it might be contorted, etc.

A3.4 Other Testing—Growth concern are outlined below.

A1.4.3.1 *Ampelisca abdita*—Survival of *Ampelisca abdita* in sediment that is >95 % sand may elicit excess mortality, but this has not been quantified (USEPA 1994a (1), USEPA-USACE 2001(2)). Toxicity tests conducted with *A. abdita* on sediments that are >95 % sand should be conducted with a clean control sediment characteristic of that test sediment. (John Scott, personal communication USEPA, 1994a (1).)

A1.4.3.2 *Leptocheirus plumulosus*—In short-term exposures, juvenile *L. plumulosus* have been shown to be tolerant of a wide range of grain size. *Leptocheirus plumulosus* has exhibited >90 % survival in 10-day tests. Small juveniles clean sediments ranging from about 100 % sand to about 100 % silt + clay (Schlekat et al., 1992 (13), USEPA 1994a (1)). See additional information provided in sections 12.1.3 and A2.4.3.

A1.4.3.3 *Eohaustorius estuarius*—*Eohaustorius estuarius* has exhibited acceptable (100 %) survival when exposed to clean sediments ranging from 0.6 to 100 % sand (USEPA 1994a (1)). Environment Canada (1998 (184)) reported that *E. estuarius* can tolerate up to 70 % clay in sediment toxicity tests. However, *E. estuarius* naturally inhabits sandy sediments, and a narrow correlation between survival of *E. estuarius* and grain size range should has been reported by DeWitt et al. (1989 (11)) and by USEPA (1994a (1)) with increased mortality associated with increased proportions of fine-grained sediment. Therefore, it may be

desirable to include clean control sediments with a range of particle sizes characteristic of those of the test sediment(s) in toxicity tests conducted with *E. estuarius*.

A1.4.3.4 *Rhepoxynius abronius*—*Rhepoxynius abronius* has been used to test sediments with a wide range of sediment grain sizes. However, *R. abronius* naturally inhabits clean, fine, sandy sediments, and a number of studies have shown some reduction in survival when this species is held in very fine-grained (predominantly silt and clay) sediment (DeWitt et al., 1988 (193); Long et al., 1990 (28); McLeay et al., 1991 (194); USEPA 1994a (1)). Therefore, when test sediments are predominantly silts or clays, the experimental design include at silt-clay control sediment with a range of particle sizes characteristic of the test—at least one additional group of amphipods should be sorted. This extra group represents sediment(s). Alternatively, when the initial particle size and should of test sediments are known, regression techniques can be used to evaluate protential effects of fines on *R. abronius* survival (DeWitt et al., 1988 (193)).

A1.4.4 *Effects of Pore-water Salinity*—The four amphipod species exhibit variability in 5% buffered formalin their salinity tolerance ranges. There are two options available for laboratory sediment testing regarding the choice of overlying water salinity for a given sediment. The amphipods must be fed during options are to either use the selected species-specific overlying water salinity for each test species (20 ‰ for *E. G. estuarius*, 5 or 20 ‰ for *L. plumulosus*, 28 to 32 ‰ for *A. abdita* or *R. abronius*), or to choose another target salinity (for example, to match the salinity to that of the pore water). The range of pore-water salinities in which a given species can survive for 10 d when using species-specific overlying water salinities is measured by length from the salinity application range. If tests are conducted with procedures different from those described in Table A1.1 (for example, different salinity, lighting, temperature, feeding conditions), additional tests are required to determine comparability of results (sections 1.4 and 1.10).

A1.4.5 In either scenario, the first antennae potential for a toxic response due to salinity alone exists if a species is exposed to conditions outside of its range of tolerance. For estuarine sediments, it is very important to know the base pore-water salinity of each sediment before testing is started, to choose a species that will not be affected by the telson. Measurements pore-water salinity, and to use overlying water of an appropriate salinity.

A1.4.5.1 Salinity tolerance ranges for each species are as follows: *Ampelisca abdita*: 20 to 32 ‰; (USEPA 1994a (1)); *Eohaustorius estuarius*: 2 to 34 ‰; *Leptocheirus plumulosus*: 1.5 to 32 ‰; *Rhepoxynius abronius* : 25 to 32 ‰ . While there is some evatidence of salinity-related stress for *E. estuarius* and counting *L. plumulosus* at salinity extremes, the breadth of test survivors.

A3.4.1 Chronic tests have also been conducted with this salinity tolerance exhibited by these species (DeWitt et al., 1989 (48)(11); Schlekat et al., 1992 and research(13); USEPA 1994a (1)) is underway most likely sufficient for application to determine the majority of sediments that may be encountered in an estuarine system. If it is desirable to have matching overlying and pore-water salinity from areas where pore-water salinities are 0 to 2 ‰ , an organism that has been demonstrated to tolerate this salinity range should be used, either instead of or in addition. **A3.5**— The amphipod *Hyalella azteca*— is one such species (Test Method E 1706). Likewise, sediments collected from areas of high salinity (that is, >32 ‰ for *L. plumulosus*) should probably utilize *A. abdita* , *E. estuarius*, or *R. abronius* .

A1.4.5.2 Salinity application ranges for each species are as follows: *Ampelisca abdita* with overlying water salinity of 28 to 32 ‰ : 0 to 34 ‰ , (Weisberg et al., 1992 (180) ; USEPA 1994a (1)); *Eohaustorius eestuarius* with overlying water salinity of 20 ‰ : <2 to 34 ‰ (DeWitt et al., 1989 (11); USEPA 1994a (1)); *Leptocheirus plumulosus* with overlying salinity of 20 ‰ : <1.5 to 32 ‰ (Schlekat et al., 1992 (13); USEPA 1994a (1) ; Emery et al. 1997 (8)) and *Rhepoxynius abronius* with overlying water salinity of 28 to 32 ‰ : 25 to 34 ‰ (Swartz et al., 1985 (10); Lamberson and Swartz, 1988 (187)).

A1.4.6 *Effects of Sediment-associated Ammonia*—Field-collected sediments may contain concentrations of ammonia that are toxic to amphipods. Water column no effect concentrations for the four amphipod species are presented in Table A1.4. If ammonia concentrations are above these concentrations, mortality occurring after 10 d may be sensitive due in part to a variety effects of ammonia. Depending on the application, it may be desirable to logwer the ammonia concentration by manipulating the test system before introduction of test organisms if measured ammonia in the marine environment. For example, when exposed overlying water is greater than the species specific no effect concentration (USEPA 1994a (1)). An errata sheet to USEPA (1994a (1)) suggests for dredged material from Black Rock Harbor, Connecticut, testing under the Clean Water Act or the Marine Protection, Research, and Sanctuaries Act, the following procedure can be used. This procedure was described in a December 21, 1993 guidance memorandum issued by the U.S. EPA Office of Wetlands, Oceans and Watersheds, U.S. EPA Office of Science and Technology, and U.S. Army Corps of Engineers Operations, Construction, and Readiness Division. When ammonia is present at toxicologically important concentrations and when ammonia is not a contaminant of concern, the laboratory analyst can reduce ammonia in the sediment pore water to species-specific no-effect concentrations (Table A1.4). Ammonia levels in the pore water can be reduced by sufficiently aerating the sample and replacing two volumes of water per day. The analyst should measure pore-water ammonia periodically (for example, every 1 to 3 days) until it reaches the appropriate species-specific no-effect concentration. After placing the test organism in the sediment, the analyst should ensure that ammonia concentrations remain within an acceptable range by conducting the toxicity test with continuous flow or volume replacement of overlying water not to exceed two volumes per day. The purging of ammonia using this procedure may also remove other contaminants from the pore water of the sediment. *Ampelisca abdita*

A2. PROCEDURE FOR CONDUCTING A *Leptocheirus plumulosus* w 28-D TEST FOR MEASURING SUBLETHAL EFFECTS OF SEDIMENT-ASSOCIATED CONTAMINANTS

A2.1 Introduction

A2.1.1 *Leptocheirus plumulosus* has been used extensively to test the most sensitive toxicity of H estuarine or marine sediment. The choice of this amphipod species as a test organism is based on sensitivity to sediment-associated contaminants, availability and ease of fish collection and invertebrates tested (65). This material was contaminated primarily culturing, tolerance of environmental conditions (for example, temperature, salinity, grain size), ecological importance, ease of handling in the laboratory, and ease of measuring test endpoints. Additionally, this species is intimately associated with polyaromatic hydrocarbons sediment by nature of its burrowing and heavy metals. At a concentration feeding habits. *L. plumulosus* is tolerant of 5 mg/L suspended Black Rock Harbor sediment, growth, salinity values between >1 to 35 ‰ and consequently sexual maturation, were delayed, and effects were seen sediment from fine- to coarse-grained. Field validation studies have shown that amphipods are absent or have reduced abundances at sites where toxicity has been demonstrated in the laboratory tests. Amphipod sediment toxicity tests have been successfully performed for regulatory and research purposes by numerous laboratories, including state and federal government agencies, private corporations, and academic institutions (see Section 1 for additional details).

A2.1.2 Guidance for conducting 10-d tests with *L. plumulosus* is described in Annex A1. Most standard whole sediment toxicity tests have been developed to produce a survival endpoint with potential for a sublethal endpoint (reburial) with some species. Methods that measure sublethal effects have either not been previously available or used routinely to evaluate sediment toxicity (Craig, 1984 (27); Dillon and Gibson, 1986 (208); Ingersoll and Nelson, 1990 (30); Ingersoll, 1991 (209) ; Burton et al., 1992 (210); USEPA 2000 (206); Test Method E 1706). Most assessments of contaminated sediment rely on short-term lethality testing methods (for example, 10 d; USEPA-USACE, 1991 (48)-, 1998 *A. abdita*(49) also showed sensitivity to a series). Short-term lethality tests are useful in identifying “hot spots” of sediments from New Bedford Harbor, Massachusetts, that were heavily contaminated with polychlorinated biphenyls and heavy metals.

A3.5.1 All routes of exposure have contamination, but may not been fully examined for *A. abdita*. Since it is a particle feeder, it will be exposed sensitive enough to evaluate moderately contaminated particles in suspension or on the areas. However, sediment surface. This amphipod feeds ventral side up in its tube, by quality assessments using its second antennae to pick up particles or by capturing small particles carried to the mouth in the current created by the action sublethal responses of the pleopods benthic organisms, such as growth and s reproduction, have been used to successfully evaluate moderately contaminated areas (Scott, 1989 (46). Therefore,(12); Niewolny et al. 1997 *Ampelisca*(211); DeWitt et al. 1997c (212)).

A2.1.3 The 28-d toxicity test with *L. plumulosus*'s feeding current exposes it to overlying water. Pore water also enters the tube, and research is currently underway a test with a lethality endpoint and two sublethal endpoints: growth and reproduction. These sublethal endpoints have potential to determine the extent provide a toxic response to which *A. abdita* is exposed to this interstitial water. In chemicals that might not cause acute effects or significant mortality in a flow-through system, it test. Sublethal response in 28-d exposures is also valuable for population modeling of contaminant effects. These data can be used for population-level risk assessments.

A2.1.4 A1.2 describes guidance for conducting the sensitivity 28-d test with *L. plumulosus* that this amphipod shows can be used to contaminated sediments is due primarily to exposure to pore water contaminants, since evaluate the overlying water effects of sediment contaminants are continually removed. It might on survival, growth, and reproduction. Refinement of these methods may be possible to use other species described in future editions of *Ampelisca* in toxicity tests. For information this standard, after additional laboratories have successfully used this method (section 15.6). These methods are based on congeners, see Bousfield procedures described in DeWitt et al. (1997a (44)(7) and Mills; 1997b (46, 63).

A4. GRANDIDIERELLA JAPONICA

A4.1 Ecological Requirements—*Grandidierella japonica* (52) was accidentally introduced into San Francisco Bay(25) and some other northern California bays by unknown means. It was first collected in 1971 Emery et al. (1997 (50). Later it was found in southern California where large populations are known(8)).

A2.1.5 Results of tests using procedures different from Upper Newport Bay and Shoreline Aquatic Park the procedures described in Long Beach (52). It has proved to A2.2 may not be a useful test species for environmental studies in southern California. The toxicity test should be conducted at 15 to 19°C using sea water with salinities between 30 comparable, and 35 g/kg. *Grandidierella japonica* lives in a variety these different procedures may alter contaminant bioavailability. Comparisons of results obtained using modified versions of these procedures might provide useful information concerning new concepts and procedures for conducting sediment types that makes it possible to conduct tests with a variety of sediment types (sands, silts, estuarine or elays):

A4.2 Collecting and Handling Techniques—This species can be collected intertidally and subtidally marine organisms. If tests

are conducted using procedures different from the localities listed above. The upper 2 those described in this standard, additional tests are required to determine comparability of sediment should be collected results (sections 1.4 and placed in a bucket with sea water. The contents should be gently stirred and the supernatant fluid decanted into a 1.0-mm sieve. The material retained on the screen should be transferred to a container 1.10).

A2.2 Procedure for transport to the laboratory. In the laboratory, the material should be placed in Conducting a sorting tray (white) containing sea water. Amphipods can be picked up using a bulb pipette with a 5-mm diameter. Females carrying embryos in their marsupium should not be used. This stage in the life cycle can be detected with the naked eye after some experience. These females can be set aside to establish cultures if so desired.

A4.2.1 For acclimation, *G. japonica* *Leptocheirus plumulosus* can be placed in an aquarium containing a 1-cm deep layer 28-d Test for Measuring Sublethal Effects of 0.5-mm sieved sediment from the collection site at a density of about 10 to 15 amphipods per 100 cm² of surface area. Two to three days are sufficient Sediment-associated Contaminants

A2.2.1 Recommended conditions for acclimation to the conducting a 28-d chronic sediment toxicity test environment. A sieve with a 1.0-mm diameter mesh size can be used to separate *G. japonica* from *L. plumulosus* are summarized in Table A2.1. A general activity schedule is outlined in Table A2.2. Decisions concerning the acclimation sediment at various aspects of experimental

TABLE A2.1 Test Conditions for Conducting a 28-d Sediment Toxicity Test with *Leptocheirus plumulosus* (USEPA-USACE 2001 (2))

Parameter	Conditions
1. Test type:	Whole sediment toxicity test, static-renewal
2. Test sediment grain size:	>5 % silt and clay to <85 % clay
3. Test sediment pore-water salinity:	1 to 35‰
4. Overlying water salinity:	5 ‰ if pore water is 1 to 10 ‰, 20 ‰ if pore water is >10 to 35 ‰; Alternatively, the salinity of the overlying water can be adjusted to a selected target salinity (for example, one representative of the salinity regime at the site of interest). If tests are conducted at a different salinity, additional tests are required to determine comparability of results (sections 1.4 and 1.10).
5. Test sediment pore-water ammonia:	< 60 mg/L (total mg/L, pH 7.7); < 0.8 mg/L (unionized mg/L, pH 7.7)
6. Test sediment pore-water sulfides:	Not established.
7. Temperature:	Daily limits: 25°C (±3°C); 28-d mean: 25°C (±2°C)
8. Light quality:	Wide-spectrum fluorescent lights
9. Illuminance:	500 to 1000 lux
10. Photoperiod:	16 h light; 8 h dark
11. Test chamber:	1-L glass beaker or jar with 10-cm inner diameter
12. Sediment volume:	175 mL (about 2-cm depth)
13. Sediment preparation:	Press-sieved through 0.25-mm sieve.
14. Overlying water volume:	Fill to 950 mL mark in test chamber (about 775 mL of water)
15. Renewal of overlying water:	3 times per week: siphon off and replace 400 mL
16. Source:	Laboratory cultures
17. Life stage and size:	Neonates: age-selected (<48 h old) or size-selected: retained between 0.25-mm and 0.6-mm mesh screens.
18. Number test organisms/chamber:	20
19. Number of replicate chambers/	5 for toxicity test; >2 additional replicate chambers for pore-water treatment: ammonia (Day 0 and Day 28)
20. Diet:	Days 0 to 13, 20 mg Tetramarine® per test chamber; Days 14 to 28, 40 mg Tetramarine® per test chamber.
21. Feeding schedule:	3 times per week (M-W-F) after water renewal.
22. Aeration and dissolved oxygen:	Aerate constantly with trickle flow of bubbles
Daily limits:	>3.6 mg/L (50 % saturation) 28-d mean: >4.4 mg/L (60 % saturation)
23. Overlying water:	Clean seawater, natural or reconstituted water; same source as used for culturing.
24. Overlying water quality and monitoring	Daily temperature in water bath or in an additional replicate chamber, daily frequency: minimum/maximum recommended; salinity, temperature, DO, and pH at test initiation and termination, and in one replicate per sediment treatment preceding water renewal during the test (three times per week); aeration rate daily in all containers; total ammonia on Days 0 and 28 in one replicate per treatment.
25. pH:	7.0 to 9.0 pH units
26. Pore-water quality:	Total ammonia, salinity, temperature, and pH of pore water from surrogate containers on Days 0 and 28; recommended in whole sediment before testing.
27. Test duration:	28 d
28. Test organism observations:	Observe condition and activity in each test chamber preceding water renewal (3 times per week).
29. Endpoints:	Survival, growth rate, and reproduction.
30. Test acceptability:	Minimum mean control survival of 80 %, growth and reproduction measurable in all control replicates, and satisfaction of performance-based criteria outlined in Table A2.3.

TABLE A2.2 General Activity Schedule for Conducting a 28-d Sediment Toxicity Test with *Leptocheirus plumulosus* (USEPA-USACE 2001 (2))

Day	Activity
Preparation Pretest	Start or renew cultures about 6 to 8 weeks in advance of test initiation. Increase culture water temperature to about 25°C about 2 weeks in advance of test initiation.
Pretest	Determining pore-water salinity of test sediment and acclimate <i>L. plumulosus</i> cultures to overlying water salinity to be used in testing.
Day-1	Layer sediment in test chambers, add overlying water. Measure pore-water total ammonia in whole sediment and begin purging procedures, if appropriate (section A2.4.5). Measure tare weight of weigh boats for dry weights. Set up positive control reference-toxicity test chambers if appropriate.
Initiation Day 0	Measure pore-water total ammonia, temperature, salinity, and pH in an additional replicate chamber. Measure salinity, temperature, DO, and pH in all test chambers. If water quality parameters are within test ranges, proceed with initiation; if not, correct problem and re-measure water quality. Obtain neonate test organisms, initiate test, and initiate positive control reference toxicant test if conducted. Only feed if a Monday, Wednesday, or Friday. Prepare 3 sets of 20 neonates for initial weight of growth rate endpoint; rinse in deionized water; dry overnight at 70°C, and weigh or measure length on Day 1 or later.
Positive Control Reference-toxicity Test Day 1 to 3	Measure and record water quality parameters in one replicate test chamber from each positive control treatment.
Day 4	Measure water quality parameters and record observations of amphipod activity in all positive control test chambers. Terminate the positive control reference-toxicity control test if conducted.
Maintenance of 28-d Test Daily	Check aeration in all test chambers and test temperature (water bath, environmental chamber, or in an additional replicate chamber). If aeration is interrupted in a test chamber, measure and record DO before resumption of aeration. Check photoperiod controllers.
3 Times per Week (M-W-F)	Measure water quality in one replicate test chamber per sediment treatment. Record observations of amphipod activity and condition of sediment and water in all test chambers. Siphon off and replace 400 mL of water in all test chambers. Add food to all test chambers.
Termination of 28-d Test Day 28	Measure salinity, temperature, DO, and pH in all test chambers. Measure tare weight of weight boats for dry weight measurements. Terminate 28-d test: sieve adults and offspring from sediment, count surviving adults, prepare adults for drying, and dry to constant weight at 70°C. Count offspring, or preserve and stain offspring.
Day 29 or later	Measure dry weight or length of adults. If offspring were preserved, count them.

design, such as the time number of treatments and water quality characteristics, should be based on the initiation purpose of the test and the methods of data analysis (Section 13).

A2.2.2 The 28-d chronic sediment toxicity-test.

A4.3- test with Toxicity Test Specifications *L. plumulosus*—The toxicity test should be run at 15 to 19 ± 3°C using 30 to 35 g/kg overlying water in the test chambers. The test chamber is usually conducted at 25°C and a standard 1-L glass beaker salinity of either 5 ‰ or 20 ‰ with a 10-cm internal diameter. Beakers should be covered with 16 h light : 8 h dark photoperiod at an 11.4-cm diameter watch glass to reduce contamination illuminance of about 500 to 1000 lux (Table A2.1). Alternatively, the contents and evaporation salinity of the overlying water and test material. Aeration can be provided adjusted to each test chamber through a 1-mL glass pipette that extends between selected target salinity (for example, one representative of the beaker spout and salinity regime at the watchglass cover to a depth not closer than 2 cm site of interest). If tests are conducted with procedures different from the sediment surface. Sediment those described in the test chambers should be 2 cm deep, and toxicity test water should be added up to the 700-mL mark on the beakers. Sediment and water should be added to beakers the day before the amphipods Table A2.1 (for example, different salinity, lighting, temperature, feeding conditions), additional tests are added, required to allow suspended determine comparability of results (section 1.10).

A2.2.3 Test chambers are 1-L glass chambers containing 175 mL of sediment particles to settle, and to allow time for equilibration about 775 mL of temperature and the sediment-water interface:

A4.3.1 After the overnight equilibration time, 20 overlying seawater. Twenty neonate amphipods are distributed added to each of the test chambers, with additional toxicity test water to bring chamber at the water level up to the 950-mL level. The amphipods should be allowed 5 to 10 min to bury into the test substrate. Any amphipods that have not buried within that time or appear

damaged should be replaced, unless the amphipods are repeatedly burrowing into the sediment and immediately emerging in an apparent avoidance response to the test substrate. In that case amphipods are not replaced. Amphipods are not removed from the surface start of a test. Five replicate test-sediment during the course of the test even if they appear dead since some amphipods that seem dead containers per treatment are actually alive recommended for routine testing. Exposure is static-renewal with water exchanges and might re-bury in the feeding three times per week, on Monday, Wednesday, and Friday. The test-sediment. At the termination of the test, the re-burial data organisms are fed after water renewals. Overlying water can be used to determine an EC50 for a sublethal measurement.

A4.3.2 The toxicity test should be terminated and the amphipods recovered using a 0.5-mm sieve. Surviving amphipods should be allowed 1 h to re-bury in a 2-cm deep layer of clean, collection site sediment.

A4.4 Life Cycle and Age Class—*Grandidierella japonica* has a short life cycle and is capable of completing four culture, surface, site, or five life cycles a year under laboratory conditions of 20°C. Immature amphipods, 3 reconstituted water adjusted to 6 mm in total length, should be used in the toxicity test. No females carrying embryos in their marsupium should be used in these tests. Animals can be cultured in test salinity. For site-specific evaluations, the laboratory on a diet characteristics of powdered fish flakes. It is easier to initiate such a culture with females carrying embryos.

A4.5 Interpretation—In interpreting the data from acute toxicity tests, it overlying water should be kept in mind that the reproductive ability or long-term survival might be affected by contaminants at lower concentrations than those that produce a lethal or sublethal effect in a short-term test. Despite these limitations, the toxicity test using Grandidierella japonica has been used in detecting sediment toxicity or toxic elements. Its ability as similar as possible to live in a burrow in a variety of sediment types gives broad application for the use of G. japonica in research and regulatory applications.

A5. LEPTOCHEIRUS PLUMULOSUS

A5.1 Ecological Requirements—*Leptocheirus plumulosus* (family Aoridae) is an infaunal amphipod distributed subtidally along the east coast of the United States from Cape Cod, Massachusetts to northern Florida (54). In Chesapeake Bay, *L. plumulosus* is indigenous to oligohaline and mesohaline regions (55, 57, 58), though it can tolerate an even broader salinity range, from near 0 to 33 g/kg (55, 56, 58). This species constructs U-shaped burrows in sediments ranging from fine sand to silty clay (56-58). Due to its broad salinity and site where sediment tolerances, it is a desirable test species collected. Requirements for east coast estuarine sediments and has been used successfully test acceptability are summarized in the assessment of contaminated sediments in Chesapeake Bay (59-61) Table A2.3.

A5.2—

A2.3 General Procedures

A2.3.1 Sediment into Test Chambers Collecting and Handling Techniques—*Leptocheirus plumulosus* is most abundant in:

A2.3.1.1 The day before the upper 2 cm addition of amphipods (Day-1), each test sediment, rarely penetrating to depths below 5 cm (66). Amphipods including control and reference sediment, should be homogenized among replicate beakers. This can be collected with benthic grab samplers (for example, Peterson, Ponar) from various tributaries achieved by mixing, by stirring manually, or by using a rolling mill, feed mixer, or other apparatus (section 10.3) or by serially spooning out small aliquots of Chesapeake Bay. The contents of sediment to each grab are sieved through test chamber. If a 0.5-mm mesh screen and the retained material quantitative confirmation of homogeneity is gently rinsed into polyethylene buckets containing collection site required, replicate subsamples should be taken from the sediment batch and w analyzed for: TOC, chesemical concentrationers, and particle size. The concentrations of ammonia in pore water should-t also be measured at the laboratory where they are aerated: start of the test.

A2.3.1.2 A 175-mL aliquot of sediment is added to each test chamber with five replicates per sediment treatment. It is desirable important that an identical volume be added to sort amphipods from collection site debris within 12 hours. A 0.5-mm mesh sieve each replicate test chamber; the volume added should provide a sediment depth of 2 cm in the test chamber. The sediment added to the test chamber should be settled by tapping the bottom or side of the test chamber against the palm of the hand or another soft object. Alternatively, sediment can be smoothed with a nylon, fluorocarbon, glass, or polyethylene spatula. Sediment known or suspected to be contaminated should bep added to test chambers in a certified laboratory fume hood.

A2.3.2 Addition of Overlying Water—As test water is added, disruption of the sediment surface should be minimized. One way to accomplish this is by use of a turbulence reducer. Possible designs of turbulence reducer include a disk cut from polyethylene, nylon, ora Teflon® spheeting (4 to 6 mil), ort a glass petri dimsh attached (open face up) to a glass pipette. The ma turbulence reducer is positioned just above the sediment surface and raised as water is added. It is convednient to mark each test chamber on the seide at 950 mL and to fill with water to reach the mark. A turbulence reducer can be rinsed into sorting trays containing collection site water. Healthy, active amphipods can be removed from detritus by using a bulb pipette with clean water between replicates of a suitable size (for example, one with treatment, but a 5-mm diameter bulb).

A5.2.1 For acclimation, *L. plumulosus* can separate turbulence reducer should be used for each treatment. The test chambers

TABLE A2.3 Test Acceptability Requirements for a 28-d Sediment Toxicity Test with *Leptocheirus plumulosus* (USEPA-USACE 2001 (2))

A. It is recommended for conducting the 28-d test with *L. plumulosus* that the following performance criteria are met:

1. Neonate *L. plumulosus*, size-selected (retained between 0.25-mm and 0.6-mm screens) or age selected (<24-h old), are used to initiate the test(s).
2. Average survival of amphipods introduced at the start of the test in the negative control sediment must be greater than or equal to 80 % at the end of the test, with no single replicate having 60 % survival or less.
3. Measurable growth and reproduction should be observed in all replicates of the negative control treatment.

B. Performance-based criteria for culturing *L. plumulosus* include the following:

1. Laboratories should perform periodic 96-h water-only reference-toxicity tests (at a minimum, one test every six months) to assess the sensitivity of culture organisms (section 11.16).
2. Records should be kept on the frequency of restarting cultures.
3. Laboratories should record the pH and ammonia of the culture water at least quarterly. Dissolved oxygen and salinity should be measured weekly. Temperature should be recorded daily.
4. Laboratories should characterize and monitor background contamination and nutrient quality of food if problems are observed in culturing or testing organisms.

C. Additional requirements:

1. All organisms in a test must be from the same source.
2. Storage of sediments collected from the field should follow guidance outlined in section 10.2.
3. All test chambers (and compartments) should be identical and should contain the same amount of sediment and overlying water.
4. Negative-control sediment and appropriate solvent controls must be included in a test. The solvent control used must not adversely affect test organisms.
5. Natural physico-chemical characteristics of test sediment collected from the field should be within the tolerance limits of the test organisms.
6. Salinity, pH, and DO, in the overlying water, ammonia in pore water and test sediment grain size should be within test condition limit of the test species (Table A2.1), or else effects of the variables need to be considered during interpretation of test results.
7. The time-weighted average of daily temperature readings must be within +2°C of the desired temperature. The instantaneous temperature must always be within +3°C of desired temperature.
8. The time-weighted average of daily salinity readings should be within $\pm 2\text{‰}$ of the selected salinity and the instantaneous salinity readings should be $\pm 3\text{‰}$ of the selected salinity.

should be covered, and placed in an aquarium (for example, 40-L) containing a 1–2 cm deep layer of 0.5-mm sieved collection site temperature controlled water bath (or acceptable equivalent) in randomly assigned positions. Aeration is started when suspended sediment has settled (often overnight). A test begins when the test organisms are added to the test chambers (Day 0).

A2.3.3 Initial Measurements—On Day 0, water quality should be measured in all test chambers before adding amphipods to test chambers. If any water quality parameter is outside acceptable limits (Table A2.1), the problem should be corrected in all replicate containers of that treatment. The water quality characteristics should be re-measure and the test should be started once the water quality characteristics are acceptable. Aberrant pH values might be caused by characteristics of certain sediments and therefore may be impractical to correct.

A2.3.4 Acclimation:

A2.3.4.1 Test organisms should be cultured at a density of approximately 200 temperature near 25°C. Amphipod cultures held below 23°C need to 300 per aquarium. Aeration be acclimated to test temperature of 25°C ($\pm 3^\circ\text{C}$) before test initiation. Ideally, test organisms should be vigorous. Two cultured in the same water that will be used in testing.

A2.3.4.2 Occasionally there is a need to three days are sufficient for acclimation perform evaluations at a temperature or salinity different than those recommended in Table A2.1. Under these circumstances, it may be necessary to acclimate organisms to the desired test environment. A gradual change temperature or salinity to prevent thermal shock that could result when organisms are moved immediately from collection site water the culture temperature or salinity to the test temperature or salinity. Reproduction and growth rates in cultures may be greatly reduced at temperatures <20°C. However, reproduction and growth is desirable. This not effected at a salinity ranging from 5 ‰ and 20 ‰ (DeWitt et al., 1997a (7)). Acclimation can be accomplished achieved by gradually increasing exposing organisms to a gradual change in temperature or salinity. However, the proportion rate of test water

change should be relatively slow to prevent shock. A change in temperature or salinity not exceeding 3°C or 3 ‰ per 24-h period is strongly recommended (Section 12). Tests at temperatures other than 25°C need to 3 days:

A5.2.2 Culture techniques be preceded by studies to determine expected performance under alternate conditions.

A2.3.5 Addition of Amphipods:

A2.3.5.1 The test is initiated when amphipods are being refined. Presently, laboratory populations can added to the test chambers. See section 12.5 for procedures for obtaining neonates for testing. Amphipods should be maintained through several generations impartially selected and placed in shallow plastic tubs transfer containers (small dishes or glass aquaria eye cups) containing a 1–2 cm layer small amount of fine grained sediment from the amphipod collection site or a texturally similar sediment (62). Water exchange is static renewal, with 30 to 50 percent test water. The number of water volume amphipods in each container replaced 2 dish should be verified by recounting before organisms are added to 4 times per week. Culture containers test chambers. To facilitate recounting, amphipods may be distributed to test chambers in batches of 5 or 10 instead of the full complement of 20. Because neonates are aerated, maintained very small, caution should be taken to ensure that each test chamber receives all 20 amphipods at test initiation. The distribution of amphipods to the test chambers needs to be done in an impartial fashion. Animals need to be added to test chambers as soon as possible following their collection to minimize handling stress and exposure to temperature changes. Three impartially selected sets of approximately 20°C, 20 neonates for initial weight determination should be isolated at the start of the test.

A2.3.5.2 To facilitate the initiation process, aeration should be stopped in test chambers immediately before adding the neonates. Sediment in test chambers should not be disrupted during the initiation procedure. Neonates from a transfer container should be poured into a test chamber. Any neonates remaining in transfer containers can be washed immediately into the test chamber using a gentle stream of 20 g/kg water at appropriate temperature and salinity. Neonates trapped at the surface water can be submerged by using a photoperiod of 16h light:8h dark. Cultures receive blunt probe or by gently dribbling a mixture few drops of micro-algae (for example, *Pseudoisochrysis paradoxa*, *Phaeodactylum tricornutum*, *Tetraselmis suecica*) and approximately 0.1 g of test or culture water onto the amphipod “gorp” (a mixture from above. A disk of fish food flakes, yeast, alfalfa powder, ground cereal leaves and shrimp maturation feed) 2–3 times per week (62). Amphipods 6-mil polyethylene, nylon, or Teflon® can be separated from acclimation or culture sediments using a 0.5 mm sieve immediately prior used on the water surface to initiating minimize disruption of the toxicity test.

A5.3 Toxicity Test Specifications—The effects sediment surface, if necessary. Rinse the disk after amphipods are added to ensure that none of different physical conditions on the amphipods have stuck to the disk. The disk should be removed once the amphipods have been introduced. A separate disk should be used for each treatment to avoid cross contamination. Aeration may be discontinued for up to 1 to 2 h to allow the amphipods to burrow into the sediment. Aeration should then continue for the remainder of the test.

A2.3.5.3 After the test organisms have been added, the test chambers should be examined for individuals that did not burrow into the sediment and might have been stressed or injured during the isolation, counting, or initiation processes. Injured or stressed test organisms will not burrow into sediment and should be removed. Neonates that have not burrowed within 1 h should be replaced with test organisms from the same sieved population, unless they are repeatedly burrowing into the sediment and immediately emerging in an apparent avoidance response. In that case, the amphipods are not replaced. The number of amphipods that are replaced in each test chamber should be recorded.

A2.3.6 Test Conditions:

A2.3.6.1 Test limits for the 28-d *L. plumulosus* to toxic materials test are currently under investigation. This species is routinely tested at 20°C, but has been tested at 25°C. Salinity provided in Table A2.1. Test sediments with characteristics that exceed these limits are subject to noncontaminant effects that should to be considered during interpretation of test results.

A2.3.6.2 Aeration—The overlying water will depend on the objectives in each test chamber should be aerated continuously after an initial settling period, except during introduction of the study. Toxicity test seawater can organisms. Filtered, dry, clean air should be bubbled through a glass or plastic pipette via plastic tubing (about 3 bubbles/sec). The tip of the pipette should be suspended 2 cm to 3 cm above the same salinity as surface of the interstitial water sediment so that it does not disturb the sediment surface. The concentration of DO in the test sediment, water overlying the ambient bottom salinity at sediment in the test chambers is maintained at or near saturation by gentle aeration. Ideally, air is bubbled through the water at a rate that maintains a high percentage of saturation (for examplee, about 90 %) but does not disturb the sediment surface. If air flow to one or more test salinity chambers is interrupted (that is, for more than 1 h), DO should be measured in those test chambers to determine whether DO concentrations have fallen below 4.4 mg/L. The 28-d mean should be >4.4 mg/L DO, and daily DO measurements should be >3.6 mg/L (50 % saturation). Results may be unacceptable for test chambers in which aeration is interrupted or DO concentrations fall to below 50 % of saturation.

A2.3.6.3 Lighting—Laboratory lighting should be maintained on a 16 h light: 8 h dark photoperiod cycle throughout the range test at an intensity of 2 500 to 32 g/kg. Laboratory investigations indicate 1000 lux.

A2.3.6.4 Feeding and Water Renewal *Leptocheirus*—A Tetramarine®-only diet is recommended for the 28-d sediment toxicity test with is tolerant *L. plumulosus*. With this diet, 400 mL of overlying water is replaced three times per week (Monday-Wednesday-Friday), after which a Tetramarine® slurry is delivered to each chamber in 1-mL aliquots. Water removal and replacement should be completed using procedures that minimize disturbance of sediment types (56); however, in the test chambers. Water can be

removed by siphoning through a grain size reference tube with fine-meshed screening over the intake to prevent uptake of amphipods. A pump can also be used to remove water. Water should not be poured from test chambers because this practice can resuspend and disturb the sediment. A separate turbulence-reducer should be used for each treatment when water is replaced to avoid cross contamination (A2.3.2). Tetramarine® is fed at a rate of 20 mg per test chamber between Days 0 to 13 and 40 mg per test chamber between Days 14 to 28. To prepare the slurry, Tetramarine® is finely ground with a food mill (blender, mortar and pestle, or a similar device) and sieved through a 0.25-mm screen. Test water is added to the appropriate amount of Tetramarine®, and the slurry is mixed on a stir plate for 15 min. USEPA-USACE (2001 (2)) provides a sample calculation for preparation of food rations. The slurry is prepared fresh for each use and needs to be somewhat stressful. Fine-grained sediments mixed continuously during feeding to prevent the Tetramarine® from settling.

A2.3.6.5 Laboratory experimentation has shown that food ration can affect the response of test animals to sediment-associated contaminants. The food ration of Tetramarine® recommended in this standard was evaluated with two other food rations in an experiment in which test animals were exposed to sediments spiked with PCB29 at concentrations between 15 and 240 mg/L (USEPA-USACE 2001 (2)). Three separate feeding rates were evaluated at each PCB29 concentration included 30 mg/60 mg (Days 0 to 13/Days 14 to 18), 20 mg/40 mg and 10 mg/20 mg per test chamber. Significant reductions in survival and growth were evident only in the highest PCB29 concentration for each of the food rations. Decreased reproduction was also evident at 240 mg/L PCB29 at each food ration as well as at 120 mg/L for the 20 mg/40 mg and 10 mg/20 mg rations (USEPA-USACE 2001 (2)). Given the generally lower reproductive rates observed at the lowest food ration, the 20 mg/40 mg feeding rate is recommended for use in this standard.

A2.3.6.6 *Water Temperature*—The test temperature was selected to approximate summertime temperature experienced by *L. plumulosus* in the wild (Holland et al., 1988 (185); McGee, 1998 (172)). The exposure chamber routinely test temperature is 25°C with a daily maximum range of $\pm 3^\circ\text{C}$ and a 28-d weighted mean of $25^\circ\text{C} \pm 2^\circ\text{C}$. Water used for renewal of test chambers should be adjusted to test temperature before use in renewals.

A2.3.6.7 *Salinity*—The target test salinity for *L. plumulosus* in a 1-L glass container is 5 or 20 ‰ depending on the pore-water salinity. The recommended overlying salinity is 5 ‰ for test with a sediment pore water from 1 to 10 ‰ or 20 ‰ for test sediments with pore water $> 10 \text{ ‰}$. The 28-d mean salinity values should deviate no more than 2 ‰ from the recommended salinity (for example, 5 ‰ or 20 ‰). Pore-water salinity of 10-0-cm each test sediment should be measured before the initiation of a test. Sediment pore water can be measured in water overlying sediment in sample containers before homogenization of sediment. Pore-water salinity can also be obtained by centrifugation (Section 10). Alternatively, the salinity of the overlying water can be adjusted to a selected target salinity (for example, one representative of the salinity regime at the site of interest). If tests are conducted with procedures different from those described in Table A2.1 (for example, different salinity, lighting, temperature, feeding conditions), additional tests are required to determine comparability of results (sections 1-E.4 and 1.10).

A2.3.7 *Measurements and Observations* :

A2.3.7.1 *Temperature* should be measured at least daily in an additional replicate chamber or from the water bath or environmental chamber. The exposure temperature of the water bath or a test chamber should be covered continuously monitored with a watch glass minimum and maximum temperature recorded daily. An additional replicate container identical to test containers is recommended for continuous temperature monitoring. The time-weighted average of daily temperature readings must be $25^\circ\text{C} \pm 2^\circ\text{C}$. The instantaneous temperature must always be within $\pm 3^\circ\text{C}$ of the contents desired temperature.

A2.3.7.2 *Salinity, DO, temperature, and evaporation pH* of the overlying water and test materials. Aeration can should be provided to each measured three times per week in at least one test chamber through a 1-mL glass pipette positioned per treatment before renewal of water. Care should be taken not closer than 2 cm from to disturb the sediment surface. Each when sampling overlying water quality.

A2.3.7.3 *Total ammonia* should be measured in overlying and pore water at test chamber initiation (Day 0 or Day -1 for pore water) and at test termination (Day 28). Salinity, pH, and temperature should contain a 2-cm deep layer be measured with each ammonia measurement. Simultaneous measurements of ammonia, salinity, pH, and temperature in sediment pore water should be taken before test initiation. If test sediments are sieved, pore-water samples for ammonia should be collected before and after sieving. Pore water can be obtained by centrifugation or from overlying water in sample containers (before pretest homogenization). If ammonia levels exceed recommended limits (Table A2.1), then ammonia reduction procedures are advisable before test initiation. However, if ammonia is the chemical of concern in the test sediments, pore-water ammonia concentrations should not be deliberately manipulated.

A2.3.7.4 Each test chamber should be examined daily to create approximately ensure that airflow to the overlying water is acceptable. Daily checks for amphipods trapped at the water surface are recommended for the first three days of a 4:1 (v/v) test. Amphipods caught in the air-water interface should be gently pushed down into the water to sediment ratio. Sediment using a blunt glass probe or drops of dilution water. The number of amphipods swimming in the water column and trapped in the air-water interface should be noted and amphipods submerged before each water renewal. The number of apparently dead test organisms should be noted, but organisms should not be removed or otherwise disturbed during the test. Exuviae may be mistaken for dead amphipods; therefore, care should be taken in identifying animals as dead.

A2.3.8 *Ending a Test*:

A2.3.8.1 The contents of each test chamber are sieved to isolate the test chambers organisms. The mesh sizes for sieving the

day before contents of the amphipods are added test chambers is 0.5 to allow suspended sediment particles 0.6 mm to settle; isolate adults and 0.25 mm to allow time for equilibration of temperature and isolate offspring. The 0.6-mm sieve should not be stacked atop the 0.25-mm sieve for thims procenss. Test- water should be used for sieving. Materfial retained on each sieve should be washed into a sorting tray with clean test water. *L. plumulosus*

A5.3.1 After overnight equilibration are easily removed from the sediment by the sieving process.

A2.3.8.2 Material that has been washed from the sieve into a sorting tray should be carefully examined for the presence of amphipods. A small portion of the test chambers, material should be sorted through at a time, and amphipods can should be randomly distributed to each of removed as they are found. Amphipods and residual sediment retained on the containers. It is desirable 0.25-mm sieve should be rinsed briefly with freshwater to saerifice remove salts and washed into a random labeled sample jar using 70 % alcohol (either ethyl or isopropyl). Use of at wide funnel supported by a ring st-20and facilitates this process. Because offspring are very small, care is needed to transfer all organisms from those being sorted on day 0 the screen to provide an initial size range estimate the sample jar. Add sufficient 70 % alcohol to preserve the amphipods, and add about 3 mL of rose bengal solution (about 1 g/L) to stain the organisms. Offspring can be counted on test animals. Twenty termination day, but waiting 2 to 3 d allows the amphipods to be more darkly stained.

A2.3.8.3 *Survival*—Numbers of live and dead adult amphipods should be tested per replicate. Animals caught on determined and recorded for each test chamber. Missing adult organisms are assumed to have died, decomposed, and disintegrated during the water's surface can test; they should be g included in the number dead in caly pculations of the percentage survival for each replicate treatment. Amphipods that are inactive but not obviously dead are observed using a g laow-power dissecting microscope or a hand-held magnifying glass. Any organismp that fails to exhibit movement (that is, neuromuscular twitch of pleopods or antennae) upon gentle prodding with a probe should be considered dead. An independent count of the number of isolated amphipods that are dead, alive, ovr moribund should be mad-5e in 10 % of replicates by a second observer. Based one the experience of one laboratory, the intralaboratory median CV for survival (sample size of 88 treatments) can be expected to be 11 % (DeWitt et al. 1997b (25); section 15.6.1). Based on one study involving 10-min to burrow into laboratories, the test sediments. Amphipods interlaboratory CV for survival ranged from 4 to 19 % (DeWitt el a. 1997b (25); section 15.6.2). It should be expected that have not burrowed within that intralaboratory CV for survival will decrease over time as a laboratory gains experience using this method. Similarly, the interlaboratory CV for survival should decrease from reported values here as more laboratories gain experience using this method.

A2.3.8.4 *Growth Rate*—Growth rate of amphipods can be replorted as daily change of average ind-wividual length or weight. However, measuring lengthy is more laborious and therefore more expensive than measurring weight to determine growth rate, and does not result in an increase in sensitivity in *L. plumulosus* 28-d test (DeWitt et al., 1997a (7)). Dry weight of amphipods can be determined as follows: (1) transferring the archived amphipods are repeatedly burrowing from a replicate out of the preservative into a crystallizing dish; (2) rinsing amphipods with deionized water; (3) transferring these rinsed amphipods to a preweighed aluminum pan; (4) drying these samples to constant weight at 60°C; and (5) weighing the sediment pan and immediately emerging in an apparent avoidance response. In that case, dried amphipods on a balance to the nearest 0.01 mg. Average dry weight of individual amphipods are not replaced. Amphipods are not removed in each replicate is calculated from these data. Due to the surface small size of test sediments the amphipods, caution should be taken during weighing 20 dried amphipods after 28-d sediment exposure may weigh less than 25 mg). The average per-capita dry weight of adult amphipods for each replicate is the course difference between the tarred weight of the toxicity test even if they appear dead, since some boat and the total weight of the boat plus dried amphipods, divided by the number of amphipods in the weigh boat. The growth rate endpoint (mg/d) is the difference between per capita adult and neonate dry weights, divided by 28 d. In other words, for each replicate, calculate: Growth Rate (mg/individual/day) = (mean adult dry weight - mean neonate dry weight)/28 (note that seem dead might actually this pooled weight of neonates is typically very small). Weigh pans need to be carefully handled using powder-less gloves and might later rebury into test substrate.

A5.3.2 The toxicity test the balance should be calibrated with standard weights with each use. Forceps can also be terminated after 40 days by sieving amphipods used to handle the weigh pans. Use of small aluminum pans will help reduce variability in measurements of dry weight. Weigh boats can also be constructed from test sediments using a 0.5-mm mesh screen. Mortality is sheets of aluminum foil. Amphipod body length (± 0.1 mm) can be measured from the base of the first antennae to the tip of the third uropod along the curve of the dorsal surface. The use of a digitizing system and microscope to measure length has been described in Kemble et al. (1994 (51)) for this short-term test. Burrows generally disintegrate during sieving *Hyaella azteca* and DeWitt et al. (1992a (6) and 1997a (7)) for *Leptocheirus plumulosus*. Based on the experience of one laboratory, the intralaboratory median CV for growth (sample size of 87 treatments) can be transferred expected to be 3 % (DeWitt et al. 1997b (25); section 15.6.1). Based on one study involving 10 laboratories, the interlaboratory CV for growth ranged from 14 to 38 % (DeWitt el a. 1997b (25); section 15.6.2). It should be expected that intralaboratory CV for growth rate will decrease over time as a s laboratory gains experience using this method. Similarly, the interlaboratory CV for growth raten should decrease from reported values here as more laboratories gain experience using this method.

A2.3.8.5 *Reproduction*—The ability offspring should be counted within 2 weeks of surviving amphipods terminating the test. It may be possible to count the offspring the day the experiment is broken down. If not, preserve offspring in 70 % alcohol (either ethyl or isopropyl). Transfer preserved, stained offspring to a fine screen (<0.25-mm mesh) and rinse with freshwater to remove

alcohol and excess stain. Rinse the live or preserved neonates into a shallow dish and count them under magnification, such as a dissecting microscope. Record the number of offspring. For QA, 10 % of the samples should be recounted by a second analyst. The reproduction endpoint is calculated as the number of offspring per living adult. Based on the experience of one laboratory, the intralaboratory median CV for reproduction (sample size of 88 treatments) can be expected to be 18 % (DeWitt et al. 1997b (25; section 15.6.1). Based on one study involving 10 laboratories, the interlaboratory CV for survival ranged from 35 to 102 % (DeWitt et al. 1997b (25; section 15.6.2). It should be expected that intralaboratory CV for reproduction will decrease over time as a laboratory gains experience using this method. Similarly, the interlaboratory CV for reproduction should decrease from reported values here as more laboratories gain experience using this method.

A2.3.9 Control Performance Issues and Revisions to the Method—The *Leptocheirus plumulosus* 28-d sediment toxicity test, like all experimental systems, is subject to occasional failures. Because the *L. plumulosus* 28-d sediment toxicity test endpoint.

A5.3.3 Other Testing—Partial life cycle is more complex and of longer duration than the 10-d sediment toxicity tests (28 described in Annex A1, there are more opportunities for problems to 30 days) initiated occur in this long-term test than in the short-term tests. Problems with juveniles the test are most readily detected by failure to meet test acceptability criteria in the control treatment (Table A2.1 and Table A2.3), such as mortality <20 % or failure of amphipods to grow or reproduce. Test failures usually can be attributed to a failure to maintain one or more test requirements described in Table A2.1 and Table A2.3; however, tests sometimes fail inexplicably. Possible causes for unaccountable test failures have included overfeeding (for example, leading to anoxia or increased production of hydrogen sulfide), poor health of test animals (that is, culture failure), or accidental introduction of toxic materials into test chambers. Scientists from the USEPA and the USACE observe that the frequency of failure decreases as the laboratory and staff using the test gain more experience through conducting the test; however, neither agency has explicit data on the frequency of failure. Users of this species, with amphipod length test should be aware of this possibility and survivorship prepare for the possibility to rerun the test on occasion. The method for the *L. plumulosus* 28-d sediment toxicity test will be revised as new experimentabl data reveal test conditions that reduce the probability of possible test failure.

A2.4 Interpretation of Results

A2.4.1 This section describes information that is currently underway useful in helping to determine interpret the optimum conditions for these tests:

A5.4 results of sediment toxicity tests with *Life Cycle and Age Classes*—*Leptocheirus**L. plumulosus*. Section 13 provides additional information on analyses and reporting of toxicity test data.

A2.4.2 Influence of Indigenous Organisms—Indigenous organisms may be present in field-collected sediment. The presence of organisms taxonomically similar to the test organism or the presence of the test organisms in the sample (McGee et al. 1999 (41)) can make interpretation of treatment effects difficult. Predatory organisms can adversely affect test organism survival. For example, Redmond and Scott (1989 (104)) showed that the polychaete *Nephtys incisa* † can consume amphipods under test conditions. All control, reference, and test sediment should be press-sieved through 0.25-mm mesh to avoid these complications. If test sediment is not sieved, the number and species-capable of p indigenous organisms should be determined to better interpret results.

A2.4.3 Effects of Sediment Grain Size—*L. plumulosus* tolerates a least two broods, with peak periods wide range of sediment types. There is generally little effect on survival, growth rate, or reproduction when coarse-grained (sand) or fine-grained (predominantly silt and clay) sediment is used. See section 12.1.3 for additional detail. In some studies, *L. plumulosus* has exhibited >90 % survival in clean sediment ranging from nearly 100 % sand to m nearly 100 % silt + clay (Schlekat et al., 1992 (13), USEPA 1994a (1)). However, adverse effects can occur in sediment with very high levels of clay or sand. Laboratory studies have shown significant reduction in survival when clay content exceeded 84 %, and survival, growth and reproduction were significantly reduced in the fall 100 % sand (Emery et al., 1997 (56, 67). Gravid females(8)). Results have been equivocal from controlled tests with mixed grained sediments (between 10 % and 90 % silt/clay). Emery et al. (1997 (8)) found an increase in Chesapeake Bay growth as sediment coarseness increased up to 75 % sand. DeWitt et al. (1997a (7)) reported enhanced growth in finer-grained sediment as D compared with more coarse-grained material, but the difference in growth was not considered to be biologically significant (DeWitt et al., 1997a (7)). Therefore, *L. plumulosus* should be tested with sediment with silt/clay content between 5 % and as early as February, indicating that timing of reproduction varies yearly depending on climatic conditions. In cultured populations, females produce multiple broods 85 % (Table A2.1). If sediment characteristics exceed these bounds, an appropriate clean control and gravid females are available year-round reference sediment should be incorporated into the test to separate effects of sediment-associated contaminants from effects of particle size.

A2.4.4 Effects of Pore-water Salinity (62). Size:

A2.4.4.1 The range of field-collected test organisms might depend on salinity in which a given species can survive when the size structure overlying water salinity is matched to that of the field population, as pore-water salinity is the m salinity tolerance range. The potential for a toxic response caused by salinity alone exists if a species is exposed to conditions outside of amphipods collected in early spring its range of tolerance. For estuarine sediment, it is generally greater than those collected in important to know the pore-water salinity of each sediment before testing is started and to use overlying water of an appropriate salinity. *L. plumulosus* is not recommended for testing with freshwater sediments (<1 ‰ pore-water salinity or with sediments having pore-water salinity >35 ‰ until further testing is completed to confirm acceptable response in organisms (DeWitt et al. S, 1997a

(7)). This standard recommends use of standard salinity of overlying water for testing (that is, 5 ‰ or 20 ‰; Table A2.1).

A2.4.4.2 *L. plumulosus*, a euryhaline species, can survive and thrive in a wide range of cultured amphipods is less variable seasonally. Immature salinity conditions. The salinity tolerance and adult amphipods, approximately 3 application range for this amphipod is 1 to 35 ‰ (DeWitt et al., 1989 (11); DeWitt et al., 1992a (6); Schlekot et al., 1992 (13); DeWitt et al., 1997a (7)). Although there is some evidence of salinity-related stress for *L. plumulosus* at salinity extremes, the base breadth of salinity tolerance exhibited by this species is most likely sufficient for application to the majority of sediments that might be encountered in an estuarine system (that is, pore-water salinity from 1 to >30 ‰).

A2.4.4.3 This method recommends testing with an overlying water salinity of either 5 or 20 ‰; the end choice of overlying water salinity is dependant on the third pleon segment along pore-water salinity of test sediment. Alternatively, the dorsal surface, should salinity of the overlying water can be adjusted to a selected target salinity (for example, one representative of the salinity regime at the site of interest). If tests are conducted with procedures different from those described in Table A2.1 (for example, different salinity, lighting, temperature, feeding conditions), additional tests because they are easy required to handle determine comparability of results (sections 1.4 and 1.10).

A2.4.4.4 Although matching overlying and pore-water salinity values in test containers might be appropriate for some study designs, this practice is logistically complicated and normally impractical to accomplish. Acclimation of age, size, sex, and seasonal variation amphipod cultures to the appropriate salinity is required. Moreover, if sediment samples to be tested have different pore-water salinity values, care needs to be exercised to ensure that renewals are completed with water of field-collected organisms on the sensitivity appropriate salinity.

A2.4.5 *Effects of Sediment-associated Ammonia:*

A2.4.5.1 Field-collected sediment may contain concentrations of pore-water ammonia that are toxic to amphipods. The water-only NOEC for *L. plumulosus* to contaminants is currently being examined. Evidence 60 mg/L (USEPA, 1994a (1)). If ammonia concentrations are above this value at test initiation, mortality may be due in part to date indicates mixed-sex populations within effects of ammonia. Depending on test application, it might be desirable to lower the ammonia concentration by manipulating the test system before introduction of test organisms if measured ammonia in the pore water or overlying water is greater than the NOEC. However, if ammonia is the chemical of concern in the test sediments, pore-water ammonia concentrations should not be deliberately manipulated. If sediment toxicity tests are conducted to evaluate the acceptability of dredged material for disposal, the manipulations could be performed. Section 13.3.6 references methods for conducting TIEs to determine whether ammonia is contributing to the toxicity of sediment samples. Manipulations involve flushing the test system by renewing a specified amount of overlying water until ammonia concentrations are reduced (section A1.4.5). The effects of dilution of ammonia on pore-water concentration is not known. Due to this uncertainty, one option could be to monitor pore-water concentrations.

A2.4.5.2 If ammonia is of concern to the regulatory application associated with the sediment toxicity test, overlying water should be sampled about 1 cm above the sediment surface before introduction of test organisms on Day 0. Pore-water ammonia should be measured when sediment samples are prepared for testing. If both the pore water and overlying water ammonia concentrations are <60 mg/L, then the test may proceed normally. If the ammonia concentration is >60 mg/L in a given sample, then ammonia level can be reduced by aerating the sample to saturation and replacing 2 volumes of overlying water per day. Purging pore-water ammonia (up to 60 mg/L) from test sediments before starting the toxicity test, and 96 h employing the routine replacement of overlying water in each test chamber every other day (M-W-F) did result in a consistently reduced pore-water ammonia concentration throughout the 28 days from about 60 mg/L to a minimum about 1 mg/L (DeWitt et al., 1997a (56, 59-61)).

A5.5 (7) *Control Survival—Mean control survival using*. **Similar results were obtained by other researchers (Moore et al. 1997 *Leptocheirus plumulosus* (26); Moore et al. 1995 must be at least 90% (214)).** The analyst should measure the pore-water ammonia concentration each day until it is <60 mg/L. The pore-water ammonia threshold for the chronic sediment toxicity test was based on 28-d exposures of the amphipods to be considered valid.

A5.6—sediments with experimentally-elevated pore-water ammonia (up to 60 mg/L), employing the specified purging technique before starting the toxicity test exposure, and employing the routine replacement of overlying water (M-W-F; DeWitt et al., 1997a *Sensitivity—Leptocheirus plumulosus* (7) is tolerant). **No lethal or sublethal toxicity was observed in this experiment at any one of the tested pore-water ammonia concentrations, which is most likely caused by loss of ammonia from the test system due to diffusion of pore-water ammonia from the sediments to the overlying water and the replacement of the overlying water three times per week. Because additional replicate containers are required for pore-water measurements, a range minimum of sediment types two additional replicate containers are required (one for Day 0 and salinities. The sensitivity of this species is comparable one for Day 28). Additional replicate containers should be prepared if pore-water ammonia levels are high enough to require several successive days for pore-water ammonia reduction. When ammonia concentrations are reduced to <60 mg/L, testing should be initiated by adding test organisms.**

A2.4.6 *Hydrogen Sulfide*—Hydrogen sulfide occurs naturally in anoxic marine sediments. Sims and Moore (1995 *Hyatetta azteca* (213)) conducted an extensive review of the literature that focused on the effects of hydrogen sulfide on benthic organisms. Sims and Moore (1995 in 96 h (213)) reported that tube-building amphipods circulate oxygenated water through their burrows, thus reducing or eliminating exposures to pore-water hydrogen sulfide. In acute experiments, however, dissolved sulfides have been

shown to be toxic to marine amphipods *R. abronius* (56, 61). A review and *E. estuarius* (48-h LOECs of benthic surveys 1.47 and sediment contamination in Chesapeake Bay indicates a negative correlation between 1.92 mg/L total sulfide respectively; Knezovich et al. 1996 (215)). Currently, no data exist regarding the presence sensitivity of *L. plumulosus* and to hydrogen sulfide in 28-d exposures. Additional information on the degree tolerance of aquatic organisms to sulfides can be found in Batgariõnao (1992 (66, 68)).

A5.7 (216).

A2.4.7 Total Organic Carbon (TOC) Interpretation—When interpreting—Test sediment TOC content can vary greatly, ranging from near 0 to >10 %. The amount of TOC can affect test organism survival, growth, and reproduction. Limited evidence suggests that the *L. plumulosus* chronic test is tolerant to most TOC concentrations; however, Scott et al. (1996 (217)) reported that growth and reproduction may be lower in uncontaminated field sediments having <2 % TOC concentrations. An analysis of organism response over a wide range of sediment TOC was completed by DeWitt et al. (1997b (25)) using reference sediment data from two studies. No effect on survival, growth, or reproduction was detected for sediments with TOC concentrations ranging from 1 to 7 % TOC. There was some evidence of significantly decreased survival, growth, and reproduction in <1 % TOC sediments. No data were available for test sediments with TOC >7 %. Therefore until additional data are generated, if test sediment TOC concentrations are <1 % or >7 %, a TOC control or reference sediment with similar TOC should be kept in mind tested concurrently.

A2.4.8 Future Research— Research to find methods that reduce the early life stage, variability of the reproductive ability, or growth rate and reproduction endpoints could lead to improvements in the long-term survival statistical power of the *L. plumulosus* might be affected by contaminants at concentrations lower than those that produce a chronic toxicity test. A second “round-robin” study using the revised feeding regime and using only laboratories with considerable experience running this toxicity test, could provide improved estimates of the interlaboratory accuracy and precision of each endpoint. Additional research is needed to evaluate the relative toxicological sensitivity of the lethal response. Partial life cycle and sublethal endpoints to a wide variety of chemicals spiked in sediment and to mixtures of chemicals in sediments from contaminant gradients in the field. Additional research is needed to evaluate the ability of the test’s lethal and sublethal endpoints to estimate the responses of populations and communities of benthic invertebrates to contaminated sediments. Additional research is also needed to link the toxicity test procedures are under development for endpoints to a field validated population model of *L. plumulosus* (McGee and Spencer 2001 and should resolve these questions:

(218), Spencer and McGee 2001 (219)) which would provide additional ecological relevant interpretative guidance for the toxicity test.

REFERENCES

- (1) USEPA. 1994a . Methods for measuring assessing the toxicity of sediment-associated contaminants with estuarine and marine amphipods. EPA-600/R-94/025, Narragansett, RI: EPA/600/R-94/025, Office of Research and Development, Washington, D.C.
- (2) USEPA-USACE. 2001999. Methods for assessing the chronic toxicity of marine and estuarine sediment-associated contaminants with the amphipod *Leptocheirus plumulosus*, 1st edition. EPA-series number pending, Washington, DC: 600/R-01/020, Washington DC and Vicksburg, MS.
- (3) Swartz, R. C., DeBen, W. A., Jones, J.Ho, K. P., Lamberson, J. O., and Cole, T. A. Kuhn, M. Pelletier, F. A., “Phoxocephalid Amphipod Bioassay for Marine Sediment Toxicity.” *Aquatic Toxicology and Hazard Assessment: Seventh Symposium, ASTM-STP 854*, Cardwell, Mc Gee, R. D., Purdy, R., M. Burgess, and Bahner, R. C., Eds., ASTM, Philadelphia, 1985, pp. 284–307. J. Serbst. 2000. Sediment toxicity assessment: comparison of standard and new testing designs. *Arch. Environ. Contam. Toxicol.* 39:462-468.
- (4) Swartz, R. C., DeBen, W.Ferretti, J. A., and Cole, D. F.A., “A Bioassay for the Toxicity Calessio, J. M. Lazorchak, and C. O. Durham. 2002. Evaluation of Sediment to Marine Macrobenthos,” reduced sediment volume toxicity test procedures using the marine amphipod *Ampelisca abdita*. *Environ. Toxicol. Chem. Journal of the Water Pollution Control Federation*, Vol 51, 1979, pp. 944–950: 21: 2372-2377.
- (5) Bolton, H. S., Breteler, R. J., Vigon, B. W., Seanton, J. A., Environment Canada. 1992. Biological test method: Acute test for sediment toxicity using marine or estuarine amphipods. Environment Canada, Conservation and Clark, S. L., “National Perspective on Sediment Quality,” *U.S. Environmental Protection Agency Contract No. 68-01-6986*, Battelle, Washington, DC, 1985. Protection, Ottawa, Ontario. 83 p., EPS 1/RM/26.
- (6) International Technical Information Institute, *Toxic DeWitt, T.H., M.S. Redmond, J.E. Sewall, and Hazardous Industrial Chemicals Safety Manual*, Tokyo, Japan, 1977; Sax, N. I., *Dangerous Properties R.C. Swartz, 1992a. Development of Industrial Materials*, 6th ed., Van Nostrand Reinhold Co., New York, NY, 1984; Patty, F. A., ed., *Industrial Hygiene and Toxicology*, Vol II, 2nd ed., Interscience, New York, NY, 1963; Hamilton, A., and Hardy, H. L., *Industrial Toxicology* 3rd ed., Publishing Sciences Group, Inc., Acton, MA, 1974; Goselin, R. E., Hodge, H. C., Smith, R. P., and Gleason, M. N., *Clinical Toxicology of Commercial Products*, 4th ed., Williams and Wilkins Co., Baltimore, MD, 1976. a chronic sediment toxicity test for marine benthic amphipods. U.S. Environmental Protection Agency. CBP/TRS/89/93.



- (7) Green, N. E., DeWitt, T. H., L. A. Niewolny, S. L. Niekirk, B. D. Gruendell, W. W. Gardiner, and Turk, A., *Safety in Working A.B. Borde. 1997a. Support for development of a standard chronic sediment toxicity protocol with Chemicals*, MacMillan, New York, NY, 1978; National Research Council, the estuarine amphipod, *Prudent Practices Leptocheirus plumulosus. Prepared for Handling Hazardous Chemicals in Laboratories*, National Academy Press, Washington, DC, 1981; Walters, D. B., ed., *Safe Handling the U.S. Environmental Protection Agency, Office of Chemical Carcinogens, Mutagens, Teratogens and Highly Toxic Substances*, Ann Arbor Science, Ann Arbor, MI, 1980; Fawcett, H. H., and Wood, W. S., eds., *Safety and Accident Prevention in Chemical Operations*, 2nd ed., Wiley-Interscience, New York, NY, 1982. *Water. PNWD-2369. Battelle Pacific Northwest Division, Richland, WA.*
- (8) National Council on Radiation Protection Emery, V.L., Jr., D.W. Moore, B.R. Gray, B.M. Duke, A.B. Gibson, R.B. Wright, and Measurement, "Basic Radiation Protection Criteria," J.D. Farrar. 1997. Development of a chronic sublethal sediment bioassay using the estuarine amphipod *Leptocheirus plumulosus* NCRP Report No. 39, Washington, DC, 1971; Shapiro, J., *Radiation Protection*, 2nd ed., Harvard University Press, Cambridge, MA, 1981. (Shoemaker). *Environ. Toxicol. Chem.* 16:1912-1920.
- (9) National Institutes of Health, "NIH Guidelines Emery, V.L., Jr., and D.W. Moore. 1996. Preliminary protocol for conducting 28-day chronic sublethal bioassays using the Laboratory Use of Chemical Carcinogens," estuarine amphipod *Leptocheirus plumulosus* NIH Publication No. 81-2385, Bethesda, MD, May 1981. (Shoemaker). Technical Note EEDP-01-36. U.S. Army Corps of Engineers, Vicksburg, MS.
- (10) Drummond, R. A., Swartz, R.C., W.A. DeBen, J.K.P. Jones, J.O. Lamberson, and Dawson, W. F., "An Inexpensive Method F.A. Code. 1985. Phoxocephalid amphipod bioassay for Simulating Diel Pattern of Lighting in the Laboratory," marine sediment toxicity. In *Transactions of the Aquatic Toxicology and Hazard Assessment: 7th Volume, ASTM STP 854, R.D. Cardwell, R. Purdy, and R.C. Bahner, eds., American Fisheries Society, Vol 99, 1970, pp. 434-435; Everest, F. H., and Rogers, J., "Two Economical Photoperiod Controls for Laboratory Studies," Progressive Fish-Culturist, Vol 44, 1982, Testing and Materials, Philadelphia, PA. pp. 113-114; 284-307.*
- (11) Carmignani, G. M., DeWitt, T.H., R.C. Swartz, and Bennett, J. P., "Leaching J.O. Lamberson. 1989. Measuring the acute toxicity of Plastics Used in Closed Aquaculture Systems," estuarine sediments. *Environ. Toxicol. Chem. Aquaculture*, Vol 7, 1976, pp. 89-91. 8:1035-1048.
- (12) U.S. Environmental Protection Agency, "Ambient Aquatic Life Water Quality Criteria for Chlorine 1984," Scott, K.J. and M.S. Redmond. 1989. The effects of a contaminated dredged material on laboratory populations of the tubicolous amphipod, EPA 440/584-030, National Technical Information Services, Springfield, VA, 1985. *Ampelisca abdita. In Aquatic Toxicology and Hazard Assessment: 12th Volume, ASTM STP 1027, U.M. Cowgill and L.R. Williams, eds. American Society for Testing and Materials, Philadelphia, PA. pp. 289-303.*
- (13) "Reagent Chemicals, American Chemical Society Specifications," Am. Chemical Soc., Washington, DC. For suggestions on the testing of reagents not listed by the American Chemical Society, see "Reagent Chemicals Schlegel, C.E., B.L. McGee, and Standards," by Joseph Rosin, D. Van Nostrand Co., Inc., New York, NY, and E. Reinharz. 1992. Testing sediment toxicity in Chesapeake Bay with the "United States Pharmacopeia." amphipod *Leptocheirus plumulosus*: An evaluation. *Environ Toxicol Chem.* 11:225-236.
- (14) Kester, D. R., Duedall, I. W., Connors, D. N., and Pytkowicz, R. M., "Preparation of Artificial Seawater," *Limnology and Oceanography*, Vol 12, 1967, pp. 176-179; Zarogian, G. E., Pesch, G., and Morrison, G., "Formulation of an Artificial Sea Water Media Suitable USEPA. 1992. Proceedings from workshop on tiered testing issues for Oyster Larvae Development," *American Zoologist*, Vol 9, 1969, p. 1144; Zillioux, E. J., Foulk, H. R., Prager, J. C., freshwater and Cardin, J. A., "Using Artemia to Assay Oil Dispersant Toxicities," *Journal of the Water Pollution Control Federation*, Vol 45, 1973, pp. 2389-2396. marine sediments. Washington, D.C., September 16-18, 1992.
- (15) Davey, E. W., Gentile, J. H., Erickson, S. J., Williams, L.G., P.M. Chapman, and Betzer, P., "Removal T.C. Ginn. 1986. A comparative evaluation of Trace Metals from Marine Culture Media," *Limnology marine sediment toxicity using bacterial luminescence, oyster embryo, and Oceanography amphipod bioassays. Marine Environmental Research*, Vol 15, 1970, pp. 486-488-19: 225-249.
- (16) Bullock, G. L., Word, J.Q., J.A. Ward, B. Brown, B.D. Walls, and Stuckey, H. M., "Ultraviolet Treatment S. Lemlich. 1989. Relative sensitivity and cost of Water for Destruction of Five Gram-Negative Bacteria Pathogenic to Fishes," amphipod bioassays. In: *Journal of the Fisheries Research Board of Canada*, Vol 34, 1977; *Oceans '89 Volume 2: Ocean Pollution*. IEEE Publication No. 89CH2780-5. pp. 1244-1249-467-473.
- (17) *Water Quality Criteria*, National Academy of Sciences, Long, E.R. and National Academy M.F. Buchman. 1989. An evaluation of Engineering, 1972; EPA-R3-73-033; candidate measures of biological effects for the National Status and Trends Program. NOAA Technical Information Service, Springfield, VA, 1973, pp. 172-193; *Federal Register*, Vol Memorandum NOS OMA 45, Nov. 28, 1980, pp. 79318-79379; *Federal Register*, Vol 50, July 29, 1985, pp. 30784-30796. Seattle, WA. 105 p.
- (18) Malueg, K. W., Schuytema, G. S., Pastorak, R.A., and Krawczyk, D. F., "Effects D.S. Becker. 1990. Comparative sensitivity of Sample Storage on a Copper-Spiked Freshwater Sediment," *Environmental Toxicology sediment bioassays at three Superfund sites in Puget Sound. In: W.G. Landis and Chemistry*, Vol 5, 1986, W.H. van der Schalie, Eds., *Aquatic toxicology and risk assessment: Thirteenth volume, STP 1096*, Philadelphia, PA: American Society for Testing and Materials, pp. 245-253-123-139.
- (19) Robinson, A. M., Lamberson, J. O., Cole, F. A., McGee, B.L., C.E. Schlegel, and Swartz, R. C., "Effects E. Reinharz. 1993. Assessing sublethal levels of Culture Conditions on sediment contamination with the Sensitivity of the Phoxocephalid Amphipod; estuarine amphipod, *Rhepoxynius abronius*, to Cadmium in Sediment," *Leptocheirus plumulosus. Environmental Toxicology and Chemistry*, Vol 7, 1988, pp. 953-959. *Environ Toxicol. Chem.* 12:577-588.
- (20) DeWitt, T. H., Ditsworth, G. R., and Swartz, DeWitt, T.H., B.L. McGee, R. C., "The Effects Alden, H. Phelps, and A. Pinkney. 1992b. A multi-species, multi-lab comparison of Natural Sediment Features on the Phoxocephalid Amphipod, *Rhepoxynius*

- abronius*,” *Marine toxicity of a Chesapeake Bay sediment. Presented at the Thirteenth Annual Meeting of the Society for Environmental Research*, Vol 25, 1988, pp. 99–124. *Toxicology and Chemistry*, Cincinnati, OH, November 8-12.
- (21) Swartz, R. C., Schults, D. W., Ditsworth, G. R., and DeBen, W. A., “Toxicity of *Schlekat*, C.E., K.J. Scott, R.C. Swartz, B. Albrecht, L. Antrim, K. Doe, S. Douglas, J. A. Ferretti, D.J. Hansen, D.W. Moore, C. Mueller and A. Tang. 1995. Interlaboratory comparison of Sewage Sludge to *Rhepoxynius abronius*, a Marine Benthic Amphipod,” 10-day sediment toxicity test method using *Ampelisca abdita*, *Eohaustorius estuarius* *Archives of Environmental Contamination and Toxicology*, Vol 13, 1984, pp. 207–216. *Leptocheirus plumulosus*. *Environ. Toxicol. Chem.* 14:2163-2174.
- (22) Vieth, G. D., and Comstock, V. M., “Apparatus for Continuously Saturating Water with Hydrophobic Organic Chemicals,” *Journal of the Fisheries Research Board of Canada*, Vol 32, 1975, pp. 1849–1851; Geringerich, W. H., Seim, W. K., ambient toxicity and Schonbrod, R. D., “An Apparatus for the Continuous Generation of Stock Solutions of Hydrophobic Chemicals,” *Bulletin of Environmental Contamination and Toxicology*, Vol 23, 1979, pp. 685–689; Phipps, H. L., Holecombe, G. W., and Fiandt, J. T., “Saturator System for Generating Toxic Water Solutions for Aquatic Bioassays,” *geological assessment. Progressive Fish-Culturist Environ Toxicol Chem.*, Vol 44, 1982, pp. 115–116. 19:1753-1763.
- (23) Barnard, J. L., Word, J., W. Gardiner, and Barnard, C. M., “The Genus *Rhepoxynius* (Crustacea: Amphipoda: Phoxocephalidae) in American Seas,” *Smithsonian Contributions for Zoology*, No. 357, Smithsonian Institution Press, Washington, DC, 1982; M. Pinza. 1997. Confounding factors related to marine sediment ecotoxicological testing. PNWD-SA-4435. 1997 Short Course, Society of Environmental Toxicology and Chemistry.
- (24) DeWitt, Kohn, N.P., J.Q. Word, D.K. Niyogi, L.T. Ross, T.-H., Swartz, R. C., Dillon, and Lamberson, J.-O., “Measuring the Toxicity of Estuarine Sediments,” *Mar. Environ. Res. Environmental Toxicology and Chemistry*, Vol 8, 1989, pp. 1035–1048. 38: 1-15.
- (25) Oakden, J. M., Oliver, J. S., DeWitt, T.H., M.R. Pinza, L.A. Niewolny, V.I. Cullinan, and Flegel, A. R., “EDTA-Chelation of Zinc in Sediment: Effects on the Behavior and Mortality of Two Infaunal Amphipods,” *standard marine/estuarine chronic sediment toxicity test method using Marine Biology*, Vol 84, 1984, pp. 125–130; Reichert, W. L., Eberhart, B.T. L., and Varanasi, U., “Exposure of *Leptocheirus plumulosus*. Prepared for the U.S. Environmental Protection Agency, Office of Two Species of Deposit-Feeding Amphipods to Sediment-Associated [³H]benzo[a]pyrene: Uptake, Metabolism and Covalent Binding to Tissue Macromolecules,” *Aquatic Toxicology*, Vol 6, 1985, pp. 45–56; Varanasi, U., Reichert, W. L., Stein, J. E., Brown, D. W., and Sanborn, H. R., “Bioavailability and Biotransformation of Aromatic Hydrocarbons in Benthic Organisms Exposed to Sediment from an Urban Estuary,” *Environmental Science and Technology*, Vol 19, 1985, pp. 836–841. *Technology*. PNNL-11768. Pacific Northwest National Laboratory, Richland, WA.
- (26) Bousfield, E. L., “Haustoriidae of the United States National Museum,” *Proceedings of the American Academy of Natural Sciences*, Vol 117, No. 3512, 1965, pp. 159–240; Bousfield, E. L., “Adaptive Radiation in Sand-Burrowing Amphipod Crustaceans,” *estuarine amphipod Chesapeake Science Leptocheirus plumulosus. Environ. Toxicol. Chem.*, Vol 11, pp. 143–154. 16:1020-1027.
- (27) Bosworth, W. S., Jr., “Biology of the Genus *Eohaustorius* (Amphipoda: Haustoriidae) on the Oregon Coast,” PhD Thesis, Oregon State University, Corvallis, OR, 1976. *sediments: Review of previous methods and recommendations for future test protocols*. IEC Beak consultants, Ltd. Mississauga, Ontario.
- (28) Davis, Long, E.R., M.F. Buchman, S.M. Bay, R.J. Breteler, R.S. Carr, P.M. Chapman, J.E. Hose, A.L. Lissner, J.-C., Scott, and Hoos, R. A. W., “Use of Dehydroabietic Acid as Reference Toxicants for Salmonid Bioassays,” *Tomales Bay, CA. Journal of the Fisheries Research Board of Canada*, Vol 32, 1975, pp. 411–416; LaRoche, G., Eisler, R., and Tarzwell, C. M., “Bioassay Procedures for Oil and Oil-Dispersant Toxicity Evaluation,” *Journal of the Water Pollution Control Federation Environ Toxicol. Chem.*, Vol 42, 1970, pp. 1982–1989; Adelman, I. R., and Smith, L. L., Jr., “Standard Test Fish Development, Part I,” *EPA-600/3-76061a*, National Technical Information Service, Springfield, VA, 1976; Fogels, A., and Sprague, J. B., “Comparative Short-Term Tolerance of Zebrafish, Flagfish, and Rainbow Trout to Five Poisons Including Potential Reference Toxicants,” *Water Research*, Vol 11, 1977, pp. 811–817; Cardwell, R. D., Woelke, C. E., Carr, M. I., and Sanborn, E., “Appraisal of a Reference Toxicant for Evaluating the Quality of Oyster Larvae,” *Bulletin of Environmental Contamination and Toxicology*, Vol 18, 1977, pp. 719–725; Roberts, M. H., Jr., Warriner, J. E., Tsai, C-F., Wright, D., and Cronin, L. E., “Comparison of Estuarine Species Sensitivities to Three Toxicants,” *Archives of Environmental Contamination and Toxicology*, Vol 11, 1982, pp. 681–692. 9:1193-1214.
- (29) Hurlbert, S. H., “Pseudoreplication in Ecological Field Experiments,” *inorganic and organic selenium to Daphnia magna Ecological Monographs (Cladocera) and Chironomus riparius*, Vol 54, 1984, pp. 187–211. (Diptera). *Environ. Toxicol. Chem.* 9:1171-1181.
- (30) Green, R. H., Ingersoll, C.G. and M.K. Nelson. 1990. Testing sediment toxicity with *Hyalella azteca* *Sampling Design (Amphipoda) and Statistical Methods for Environmental Biologists Chironomus riparius*, Wiley, New York, 1979. (Diptera). In *Aquatic Toxicology and Risk Assessment, 13th volume. ASTM STP 1096 W.G. Landis and W.H. van der Schalie, eds.* Philadelphia, PA. pp. 93-109, 1990.
- (31) Snedecor, G. W., Burton, G.A., Jr. and C.G. Ingersoll. 1994. Evaluating the toxicity of sediments. In *Statistical Methods*, 5th ed., Iowa State University Press, Ames, IA, 1956, pp. 60–62. *The Assessment of Contaminated Great Lakes sediment*. US Environmental Protection Agency Report, Region V, Chicago, IL. EPA-905-B94-002.
- (32) Cohen, J., Reish, D.J. 1988. The use of toxicity testing in marine environmental research, Chapter 10. In *Statistical Power Analysis for the Behavioral Sciences*, Academic Press, *Marine Organisms as Indicators*. D.F. Soule and G.S. Kleppel eds. Springer-Verlag, New York, NY, 1977; Natrella, M. G., “The Relationship Between Confidence Intervals and Tests of Significance,” *American Statistician*, Vol 14, 1960, NY. pp. 20–22. 213-245.
- (33) Berg, E. L., ed., “Handbook for Sampling” *Harkey, G.A., P.F. Landrum, and Sample Preservation S.J. Klaine*. 1994. Preliminary studies on the effect

- of Water and Wastewater," feeding during whole sediment bioassays using *EPA-600/4-82-029*, National Technical Information Service, Springfield, VA, 1982. *Chironomus riparius* larvae. *Chemosphere* 28:597-606.
- (34) U.S. Environmental Protection Agency, "Methods for Chemical Analysis of Water Landrum, P.F. 1989. Bioavailability and Wastes," *EPA-600/4-79-020 (Revised March 1983)*, National Technical Information Service, Springfield, VA, 1983; Strickland, J. D. H., and Parsons, T. R., *A Practical Handbook toxicokinetics of Seawater Analysis, Bulletin 167*, Fisheries Research Board of Canada, Ottawa, 1968; U.S. Geological Survey, *National Handbook of Recommended Methods polycyclic aromatic hydrocarbons sorbed to sediments for Water-Data Acquisition*, U.S. Department of the Interior, Reston, VA, 1977; American Public Health Association, American Water Works Association, and Water Pollution Control Federation, amphipod, *Standard Methods for the Examination of Water and Wastewater*, 16th ed., Washington, DC, 1985. For analysis of sediments, see Plumb, R. H., Jr., "Procedures for Handling and Chemical Analysis of Sediment and Water Samples," *U.S. Environmental Protection Agency/Corps of Engineers Technical Report EPA/CE-81-1*, U.S. Army Engineer Waterways Experiment Station, Vicksburg, MS, 1981; Buchanan, J. B., "Sediment Analysis," *Methods for the Study of Marine Benthos*, 2nd ed., N. A. Holme, and A. D. McIntyre, eds., Blackwell Scientific Publications, Oxford, 1984, Chapter 3, pp. 41-65. *Pontoporeia hoyi*. *Environ. Sci. Tech.* 23:588-595.
- (35) Litchfield, J. T., Jr., Boese B.L., Lee II, H., Specht, D.T., Randall, R.C., and Wileoxon, F., "A Simplified Method Winsor, M.H. 1990. Comparison of Evaluating Dose-Effect Experiments," *Journal of Pharmacology aqueous and Experimental Therapeutics*, Vol 96, 1949, pp. 99-113; Finney, D. J., *Statistical Method solid-phase uptake for hexachlorobenzene in Biological Assay*, 2nd ed., Hafner Publishing Co., New York, NY, 1964; Finney, D. J., the tellinid clam *Probit Analysis*, 3rd ed., Cambridge University Press, London, 1971; Stephan, C. E., "Methods for Calculating an LC50," *Aquatic Toxicology and Hazard Evaluation Macoma nasuta*, F. L. Mayer, and J. L. Hamelink, eds., A mass balance approach. *Environ. Toxicol. Chem. ASTM-STP-634*, ASTM, Philadelphia, PA, 1977, pp. 65-84; Hamilton, M. A., Russo, R. C., and Thurston, R. V., "Trimmed Spearman-Kärber Method for Estimating Median Lethal Concentrations in Toxicity Bioassays," *Environmental Science and Technology*, Vol 11, 1977, pp. 714-719. Correction in Vol 12, 1978, p. 417. 9:221-231.
- (36) Sokal, R. R., Knezovich, J.P., F.L. Harrison, and Rohlf, F. J., R.G. Wilhelm. 1987. The bioavailability of sediment-sorbed organic chemicals: A review. *Water Air Soil Pollut. Biometry*, 2nd ed., W. H. Freeman and Co., San Francisco, CA, 1981, pp. 229-231. 32:233-245.
- (37) Chew, V., *Comparisons Among Treatment Means Di Toro, D.M., J.D. Mahony, D.J. Hansen, K.J. Scott, M.B. Hicks, S.M. Mays, and M.S. Redmond. 1990. Toxicity of cadmium in an Analysis sediments: The role of Variance, ARS/H/6*, Agricultural Research Service, U.S. Department of Agriculture, 1977. acid volatile sulfide. *Environ. Toxicol. Chem.* 9:1489-1504.
- (38) Gelber, R. D., Lavin, P. T., Mehta, C. R., Di Toro, D.M., C.S Zarba, D.J. Hansen, W.J. Berry, R.C. Swartz, C.E. Cowan, S.P. Pavlou, H.E. Allen, N. A. Thomas, and Schoenfeld, D. A., P.R. Paquin. 1991. Technical basis for establishing sediment quality criteria for nonionic organic chemicals using equilibrium partitioning. *Fundamentals of Aquatic Toxicology: Methods and Applications*, McGraw-Hill, 1985. *Environ. Toxicol. Chem.* 10:1541-1583.
- (39) Swartz, R.C., Code, F.A., Lamberson, J.O, Ferraro, S.P., Schults, D.W., DeBen, W. A., Sereu, K. A., W.A., Lee II, H., and Lamberson, J. O., "Sediment Toxicity Ozretich, R.J. 1994. Sediment toxicity, contamination, and Distribution of Amphipods amphipod abundance at a DDT and dieldrin-contaminated site in Commencement Bay, Washington, USA," San Francisco Bay. *Environ. Toxicol. Chem. Marine Pollution Bulletin*, Vol 13, 1982, pp. 359-364. 13:949-962.
- (40) Kemp, P. F., Cole, F. A., Anderson, B.S., Hunt, J.W., Phillips, B.M., Fairey, R., J., Oakden, J.M., Puckett, H.M., Stephenson, M., Tjeerdema, R.S., Long, E.R., Wilson, C.J., and Swartz, R. C., "Life History and Production of the Phoxocephalid Amphipod Lyons, M. 2001. Sediment quality in Los Angeles Harbor: A triad assessment. *Rhepoxymius abronius* (Barnard)," *Journal of Crustacean Biology*, Vol 5, 1985, pp. 449-464. *Environ Toxicol. Chem* 20:359-370.
- (41) Oliver, J. S., Oakden, J. M., McGee, B.L., D.J. Fisher, L.T. Yonkos, G.P. Ziegler, and Slattery, P. S., "Phoxocephalid Amphipods as Predators on Larvae S. Turley. 1999. Assessment of sediment contamination, acute toxicity and J population viability of the estuarine amphipod *Leptocheirus plumulosus* in Marine Soft-bottom Communities," Baltimore Harbor. *Marine Ecology Progress Service*, Vol 7, 1982, pp. 179-184. *Environ. Toxicol. Chem.* 18:2151-2160.
- (42) Swartz, R. C., Ditsworth, G. R., Schults, D. W., McGee, B.L. and Lamberson, J. O., "Sediment Toxicity to a Marine Infaunal Amphipod: Cadmium and Its Interaction D.J. Fisher. 1999. Field validation of the chronic sediment bioassay with Sewage Sludge," the estuarine amphipod *Marine Leptocheirus plumulosus*. *Final Report. Prepared for the U.S. Environmental Protection Agency, Office of Science and Technology by the University of Maryland, Wye Research, Vol 18, 1985, pp. 133-153. and Education Center, Queenstown, MD.*
- (43) Bosworth, W. S., Jr., "Three New Species of *Eohaustorius* (Amphipoda. Haustoriidae) from Swartz, R.C., W.A. DeBen, K.A. Secru, and J.O. Lamberson. 1982. Sediment toxicity and the Oregon Coast," *Crustaceana*, Vol 25, 1973, pp. 253-260. distribution of amphipods in Commencement Bay, Washington, USA. *Mar. Poll. Bull.* 13:359-364.
- (44) Bousfield, E. L., *Shallow-water Gammaridean Amphipoda Green, A.S. and G.T. Chandler. 1996. Life-table evaluation of New England*, Cornell University Press, Ithaca, New York, 1973. sediment-associated chlorpyrifos chronic toxicity to the benthic copepod, *Amphiascus tenuiremis*. *Arch. Environ. Contam. Toxicol.* 31:77-83.
- (45) Nichols, F. H., Levin, L., H. Caswell, T. Bridges, C. Dibacco, D. Cabrera, and Thompson, J. K., "Persistenece G. Plaia. 1996. Demographic responses

- of an Introduced Mudflat Community in South San Francisco Bay, California.” estuarine polychaetes to pollutants: life table response experiments. *Marine Ecology Progress Series*, Vol 24, 1985, pp. 83–97. *Ecological Applications*, 6:1295-1313.
- (46) Mills, E. L., “The Biology of Ciarelli, S., W.A., P.M.A. Vonck, N.M. van Straalen, and J. Stronkhorst. 1998. Ecotoxicology assessment of an Amphipod Crustacean-Sibling Species Pair,” contaminated dredged material with the marine amphipod *Corophium volutator*. *Arch Environ. Contam. Journal of the Fisheries Research Board of Canada*, Vol 24, 1967, pp. 305–355; *Toxicol.* 34:350-356.
- (47) Stickney, A. P., Meador, J.P. and Stringer, L. D., “A Study of C.A. Rice. 2001. Impaired growth in the Invertebrate Bottom Fauna of Greenwich Bay, Rhode Island,” polychaete *Armandia brevis* exposed to tributyltin in sediment. *Ecology*, Vol 38, 1957, pp. 111–122. *Marine Environmental Research*, 51:113-129.
- (48) Scott, K. J., and Redmond, M. S., “The Effects of a Contaminated Dredged Material on Laboratory Populations of the Tubicolous Amphipod, *Ampelisca abdita*,” *Aquatic Toxicology and Hazard Assessment: Twelfth Volume, ASTM STP 1027*, Cowgill, U. M., and Williams, L. R., eds., ASTM, Philadelphia, PA, 1989, pp. 289–303. Estuarine or marine Protection, Washington, D.C.
- (49) Long, E. R., and Buchman, M. F., “An Evaluation of Candidate Measures of Biological Effects of Dredged Material Proposed for Discharge in Waters of the National Status and Trends Program,” *Technical Memorandum NOS-OMA 45*, National Oceanic and Atmospheric Administration, 1989. U.S. testing manual. EPA-823-B-98-004 Washington, D.C.
- (50) Chapman, J. W., Ingersoll, C.G., E.L. Brunson, F.J. Dwyer, D.K. Hardesty, and Dorman, J. A., “Diagnosis, Systematics and Notes on *Grandidierella japonica* (Amphipoda, Gammaridea) and its Introduction to the Pacific Coast N.E. Kemble. 1998. Use of sublethal endpoints in sediment toxicity tests with the United States,” amphipod *Bulletin of the Southern California Academy of Science* *Hyalella azteca*. *Environ. Toxicol. Chem.*, Vol 74, 1975, pp. 104–108. 17:1508-1523.
- (51) Nipper, M. G., Greenstein, D. J., Kemble, N.E., W.G. Brumbaugh, E.L. Brunson, F.J. Dwyer, C.G. Ingersoll, D.P. Monda, and Bay, S. M., “Short- and Long-Term Sediment Toxicity Test Methods with of metal-contaminated sediments from the Amphipod upper Clark Fork River, MT, to aquatic invertebrates in laboratory exposures. *Environ. Toxicol. Chem. Grandidierella japonica*,” *Environmental Toxicology and Chemistry*, Vol 8, 1989, pp. 1191–1200. 13:1985-1997.
- (52) Reish, D. J., McGee, B.L., C.E. Schlekot, D.M. Boward, and LeMay, J. A., *Bioassay Manual for Dredged Sediments*, U.S. Army Corps of Engineers, Los Angeles District, Marine Biological Consultants T.L. Wade. 1995. Sediment contamination and Southern California Coastal Water Research Project, 1980; biological effects in a Chesapeake Bay marina. *Ecotoxicology*, 4:39-59.
- (53) Hong, J.-S., and Reish, D. J., “Acute Toxicity of Cadmium to Eight Species of Marine Amphipod Contaminated Sediments on marine benthic biota and Isopod Crustaceans from Southern California,” communities. In *Bulletin of Environmental Contamination Contaminated Marine Sediments Assessment and Toxicology*, Vol 39, 1987, Remediation. *National Research Council. National Academy Press, Washington, D.C.* pp. 884–888. 132-154.
- (54) Bousfield, E. L., Pfitezenmeyer, H. 1975. Benthos. In *Shallow-water Gammaridean Amphipoda A Biological Study of New England Baltimore Harbor*, Cornell University Press, Ithaca T. Koo, ed., Contribution No. 621. Center for Environmental and London, 1973. *Estuarine Studies*, University of Maryland, Solomons, MD.
- (55) Feeley, J. B., and Wass, M. L., “The Distribution and Ecology of Reinharz, E. 1981. Animal sediment relationships: A case study of the Gammaridea (Crustacea: Amphipoda) of the Lower Chesapeake Estuaries,” Virginia Institute of Marine Science, Gloucester Point, VA, 1971. Patapsco River. Open File No.6. Maryland Geological Survey, Baltimore, MD.
- (56) Schlekot, C. E., McGee, B. L., Gray, B.R., V.L. Emery, D.L. Brandon, R.B. Wright, B.M. Duke, J.D. Farrar, and Reinharz, E., “Testing Sediment Toxicity in Chesapeake Bay using D.W. Moore. 1998. Selection and optimal measures of growth and reproduction for the Amphipod sublethal *Leptocheirus plumulosus*: An Evaluation,” *Environmental Toxicology sediment bioassay*. *Environ. Toxicol. and Chemistry*, 1991, 11:225–236. 17: 2288-2297.
- (57) Holland, A. F., Shaughnessy, A. T., Scott, L. C., Dickens, V. A., Ranasinghe, J. A., and Summers, J. K., Progress Report: Ingersoll, C.G. 1995. Sediment toxicity tests. In *Long-term Benthic Monitoring and Assessment Program for the Maryland Portion Fundamentals of Chesapeake Bay Aquatic Toxicology* (July 1986–October 1987); PPRP-LTB/EST-88-1, VERSAR, Inc., Columbia, MD, 1988. 2nd edition, G. Rand, ed. Taylor and Francis, Washington, D.C. pp. 231-255.
- (58) Jordan, R. A., Ankley, G., and Sutton, C. E., “Oligohaline Benthic Invertebrate Communities at Two Chesapeake Bay Power Plants,” Thomas, N., *Interstitial Water Toxicity Identification Evaluation Approach*, *Estuaries*, 1984, 7:192–212. *Sediment classification methods compendium*, EPA 823-R-92-006, Washington, DC, 1992, pp. 5-1 to 5-14.
- (59) Maryland Department of the Environment, DeWitt, T. H., Ditsworth, G. R., and Swartz, R. C., “Effects of Natural Sediment Features on the Environment, Ecological Assessment Division,” Biological Assessment of Marina-Related Impact in Chesapeake Bay: A Pilot Study,” Final Report to U.S. Fish and Wildlife Service, Chesapeake Bay Estuary Program, Annapolis, MD, 1991. Phoxocephalid Amphipod, *Rhepoxynius abronius*: Implications for Sediment

- Toxicity Bioassays. *Marine Environmental Research*, Vol 25, 1988, pp. 99-124.
- (60) Maryland Department of the Environment, Anklely, G. T., Benoit, D. A., Balough, J. C., Reynoldson, T. B., Day, K. E., and Hoke, R. A., "Evaluation of the Environment, Ecological Assessment Division, "Assessment of Ambient Potential Confounding Factors in Sediment Toxicity of Chesapeake Bay Sediments," Final Report to Maryland Department of Natural Resources, Annapolis, MD, 1991. Tests with Three Freshwater Benthic Invertebrates," *Environmental Toxicology and Chemistry*, Vol 13, 1994, pp. 627-635.
- (61) Pinkney, A. Walsh, G. E., Gowda, S., and Rzemien, Weber, D. E., "Sediment Toxicity Testing Simon, T. L., and Brashers, L. K., "Toxicity Tests of the Baltimore Harbor Effluents With Marsh Plants in Water and C & D Canal Approach Channels with the Amphipod, Sediment," *Leptocheirus plumulosus*," Final report to the Maryland Department of Natural Resources, VERSAR, Inc., Columbia, MD, 1991. *Environmental Toxicology and Chemistry*, Vol 10, 1991, pp. 517-525.
- (62) Sewall, Suedel, B. C., and Rodgers Jr., J. E., DeWitt, T. H., and Swartz, R. C., "Laboratory Culture "Development of Chesapeake Bay Amphipods Formulated Reference Sediments for Chronic Freshwater and Acute Estuarine Sediment Quality Toxicity Testing," Abstract *Environmental Toxicology and Chemistry*, Vol 13, pp. 1163-1176; Suedel, B. C., and Rodgers, J. H. Responses of poster presented at SETAC, November 1991. *Hyalella azteca* and *Chironomus tentans* to particle size distribution and organic matter content of formulated and natural sediment. *Environmental Toxicology and Chemistry*, Vol 13, 1994, pp. 1639-1648.
- (63) Mills, E. L., "Ampelisca abdita, a New Amphipod Crustacean from Eastern North America," *Canadian Journal of Zoology*, Vol 42, 1964, pp. 559-575. freshwater test organisms to formulated sediments for use in whole-sediment toxicity tests. *Environ. Toxicol. Chem.* 18:222-230.
- (64) Santos, S. L., USEPA. Procedures for collection, handling, and Simon, J. L., "Response manipulation of Soft-Bottom Benthos to Catastrophic Disturbance in a South Florida Estuary," *Marine Ecology Progress Series*, Vol 3, 1989, pp. 347-355. aquatic sediments for toxicological testing and chemical analysis October 2000.
- (65) Rogerson, P. F., Schimmel, S. C., Green, R. H., "Sampling Design and Hoffman, G., "Chemical and Biological Characterization of Black Rock Harbor Dredged Material," *Technical Report D-85-9*, U.S. Army Engineer Waterways Experiment Station, 1985. Statistical Methods for Environmental Biologists," Wiley-Interscience, New York. 257 p., 1979.
- (66) Reinharz, E., "Animal Sediment Relationships: A Case Study Stemmer, B. L., Burton Jr., G. A., and Sasson-Brickson, G., "Effect of the Patapsee River," Open file No. 6, Maryland Geological Survey, Baltimore, MD, 1981. Sediment Spatial Variance and Collection Method on Cladoceran Toxicity and Indigenous Microbial Activity Determinations," *Environmental Toxicology and Chemistry*, Vol 9, 1990a, pp. 1035-1044.
- (67) Ray, Stephan, C. E., Mount, D. I., Hansen, D. J., Gentile, J. H., Chapman, G. L., "The Ecology A., and Brungs, W. A., "Guidelines for Deriving Numerical National Water Quality Criteria for the Protection of Benthic Macro-invertebrates in Two New Jersey Salt Marsh Waterways," Rutgers University, Ph.D. Thesis, 1982. Aquatic Organisms and Their Uses," PB85-227049, National Technical Information Service, Springfield, VA, 1985.
- (68) Chapman, P. M., "Current Approaches to Developing Sediment Quality Criteria," *Environmental Toxicology and Chemistry*, Vol 8, 1989, pp. 589-599.
- (69) Burton, G. A., "Assessment of Freshwater Sediment Toxicity," *Environmental Toxicology and Chemistry*, Vol 10, 1991, pp. 1585-1627.
- (70) Di Toro, D. M., Zarba, C. S., Hansen, D. J., Berry, W. J., Swartz, R. C., Cowan, C. E., Pavlou, S. P., Allen, H. E., Thomas, N. A., and Paquin, P. R., "Technical Basis for Establishing Sediment Quality Criteria for Nonionic Organic Chemicals Using Equilibrium Partitioning," *Environmental Toxicology and Chemistry*, Vol 10, 1991, pp. 1541-1583.
- (71) Giesy, J. P., Graney, R. L., Newsted, J. L., Rosiu, C. J., Benda, A., Kreis Jr., R. G., and Horvath, F. J., "Comparison of Three Sediment Bioassay Methods Using Detroit River Sediments," *Environmental Toxicology and Chemistry*, Vol 7, 1988, pp. 483-498.
- (72) Kemp, P. F., and Swartz, R. C., "Acute Toxicity of Interstitial and Particle-Bound Cadmium to A Marine Infaunal Amphipod," *Marine Environmental Research*, Vol 26, 1988, pp. 135-153.
- (73) *Methods for Measuring the Toxicity and Bioaccumulation of Sediment-Associated Contaminants With Freshwater Invertebrates*, Second Edition EPA 600/R-99/064, U.S. Environmental Protection Agency, Duluth, MN, 2000.
- (74) *Procedures for Assessing the Toxicity and Bioaccumulation of Sediment-Associated Contaminants With Marine Invertebrates*, EPA 600/R-94/025, U.S. Environmental Protection Agency, Duluth, MN, 1994b.
- (75) Environment Canada, Biological Test Method: Test for growth and survival in sediment using the freshwater amphipod *Hyalella azteca* Environment Canada Ottawa, Ontario, EPSRN33, 1997a.
- (76) Environment Canada, Biological Test Method: Test for Growth and survival in sediment using larvae of freshwater midges (*Chironomus tentans* or *Chironomus riparius*). Environment Canada, Ottawa, Ontario, EPSRN32, 1997b.
- (77) Sibley, P.K., Anklely, G.T., Cotter, A.M., and Leonard, E.N., Predicting chronic toxicity of sediments spiked with zinc: An evaluation of the acid volatile sulfide (AVS) model using a life-cycle test with the midge *Chironomus tentans*. *Environ. Toxicol. Chem.* 15:2102-2112, 1996.
- (78) Sibley, P.K., Benoit, D.A., and Anklely, G.T., the significance of growth in *Chironomus tentans* sediment toxicity tests: Relationship to reproduction and demographic endpoints. *Environ. Toxicol. Chem.* 16:336-345, 1997a
- (79) Sibley, P.K., Monson, P.D., and Anklely, G.T., The effect of gut contents on dry weight estimates of *Chironomus tentans* larvae: Implications for interpreting toxicity on freshwater sediment toxicity tests. *Environ. Toxicol. Chem.* 16: 1721-1726, 1997b.
- (80) Benoit, D.A., Sibley, P.K., Juenemann, J.L., and Anklely, G.T. *Chironomus tentans* life-cycle test: Design and evaluation for use in assessing toxicity of contaminated sediments. *Environ. Toxicol. Chem.* 16:1165-1176, 1997.
- (81) Ingersoll, C.G., Brunson, E.L., F.J., Hardesty, D.K., and Kemble, N.E. Use of sublethal endpoints in sediment toxicity test with the amphipod *Hyalella azteca* *Environ. Toxicol. Chem.* 17:1508-1523, 1998.
- (82) Kemble, N. E., Brumbaugh, W. G., Brunson, E. L., Dwyer, F. J., Ingersoll, C. G., Monda, D. P., and Woodward, D. F., "Toxicity of Metal-contaminated Sediments from the Upper Clark Fork River, MT to Aquatic Invertebrates in Laboratory Exposures," *Environmental Toxicology and Chemistry*, Vol 13, pp. 1895-1997, 1994
- (83) Pittinger, C.A., Adams W.J. Assessing ecological risks to benthic species in product and technology development. In *Ecological Risk Assessment of Contaminated Sediment*. Ingersoll C.G., Dillon T.-K., Biddinger G.R. (eds.) SETAC Press, Pensacola, FL, pp 11-21, 1997.
- (84) Stemmer, B. L., Burton Jr., G. A., and Leibfritz-Frederick, S., "Effect of Sediment Test Variables on Selenium Toxicity to *Daphnia magna*," *Environmental Toxicology and Chemistry*, Vol 9, 1990b, pp. 381-389.
- (85) Landrum, P. F., "Bioavailability and Toxicokinetics of Polycyclic Aromatic Hydrocarbons Sorbed to Sediments for the Amphipod *Pontoporeia hoyi*," *Environmental Science and Technology*, Vol 23, 1989, pp. 588-595.
- (86) Word, J. Q., Ward, J. A., Franklin, L. M., Cullinan, V. I., and Kiesser, S. L., "Evaluation of Equilibrium Partitioning Theory for Estimating the

- Toxicity of the Nonpolar Organic Compound DDT to the Sediment Dwelling Amphipod *Rhepoxynius abronius*,” USEPA Criteria and Standards Division. SCD No. 11, Washington, DC, 1987.
- (87) Landrum, P. F., and Faust, W. R., “Variation in the Bioavailability of Polycyclic Aromatic Hydrocarbons Sorbed to Sediments for the Amphipod *Pontoporeia hoyi*,” *Environmental Toxicology and Chemistry*, Vol 11, 1992, pp. 1197-1208.
- (88) Di Toro, D. M., Mahony, J. H., Hansen, D. J., Scott, K. J., Hicks, M. B., Mayr, S. M., and Redmond, M., “Toxicity of Cadmium in Sediments: The Role of Acid Volatile Sulfides,” *Environmental Toxicology and Chemistry*, Vol 9, 1990, pp. 1487-1502.
- (89) USEPA. Proceedings From Workshop on Tiered Testing Issues for Freshwater and Marine Sediments. Washington, DC, September 16-18, 1992a.
- (90) USEPA. Evaluation of the Apparent Effects Threshold (AET) Approach For Assessing Sediment Quality. Report of the Sediment Criteria Subcommittee, SAB-EETFC-89-027, Washington, DC, 1989a.
- (91) USEPA. Evaluation of the Equilibrium Partitioning (EqP) Approach For Assessing Sediment Quality. Report of the Sediment Criteria Subcommittee of the Ecological Processes and Effects Committee. EPA-SAB-EPEC-90-006, Washington, DC, 1990a.
- (92) USEPA. Evaluation of the Sediment Classification Methods Compendium. Report of the Sediment Criteria Subcommittee of the Ecological Processes and Effects Committee, EPA-SAB-EPEC-90-018, Washington, DC, 1990b.
- (93) USEPA. An SAB report: Review of Sediment Criteria Development Methodology for Non-Ionic Organic Contaminants. Report of the Sediment Criteria Subcommittee of the Ecological Processes and Effects Committee. EPA-SAB-EPEC-93-002, Washington, DC, 1992b.
- (94) Wenning RJ, Ingersoll, CG, editors. 2002. Use of sediment quality guidelines (SQGs) and related tools for the assessment of contaminated sediments: Summary from a SETAC Pellston workshop. Pensacola FL: SETAC Press.
- (95) USEPA. Sediment Classification Methods Compendium. EPA 823-R-92-006, Washington, DC, 1992c.
- (96) Long, E. R., and Morgan, L. G., The Potential for Biological-Study Effects of Sediment-Sorbed Contaminants Tested In the National Status and Trends Program. NOAA Technical Memorandum NOS OMA 52, Seattle, WA, 1991.
- (97) MacDonald, D.D., Carr, R.S., Calder, F.D., Long E.R., and Ingersoll, C.G. Development and evaluation of sediment quality guidelines for Florida coastal waters, *Ecotoxicology* 5: 253–278, 1996.
- (98) Ingersoll, C.G., Haverland, P.S., Brunson, E.L., Canfield, T.J., Dwyer, F.J., Henke, C.E., and Kemble, N.E., Calculation and evaluation of sediment effect concentrations for the amphipod *Hyaella azteca* and the midge *Chironomus riparius*, *J. Great Lakes Res.* 22:602–623, 1996.
- (99) Ingersoll, C.G., Dillon, T., Biddinger, R.G. (eds.). Methodological uncertainty in sediment ecological risk assessment. In *Ecological Risk Assessments of Contaminated Sediment*, SETAC press, Pensacola, FL, 389p. 1997.
- (100) Chapman, P. M., Power, E. A., and Burton, G. A., Jr., *Integrated Assessments in Aquatic Ecosystems*. In: G. A. Burton (ed.), *Sediment Toxicity Assessment*. Lewis Publishers, Boca Raton, FL, 1992.
- (101) Chapman, P.M., Anderson, B., Carr, S., Engle, V., Green, R., Hameedi, J., Harmon, M., Haverland, P., Hyland, J., Ingersoll, C.G., Long, E., Rodgers, J., Salzar, M., Sibley, P.J., Smith, P.J., Swartz, R.C., Thompson, B., and Windom, H. General guidelines for using the sediment quality triad. *Mar. Pollut. Bull.* 34:368–372, 1997.
- (102) Williams, L. R., “Methods for the EPA’s Regulatory Program,” *Environmental Testing and Analysis*, Vol 2, 1993, p. 36.
- (103) USEPA, EPA requirements for quality assurance project plans for environmental data operations, Draft interim final, EPA QA/R-5, Office of Research and Development, Washington, DC, 1994a.
- (104) Redmond, M.S. and K.J. Scott. 1989. Amphipod predation by the infaunal polychaete, *Nephtys incisa*. *Estuaries*. 12(3):205-207.
- (105) Karickhoff, S. W., and Morris, K. R., “Sorption Dynamics of Hydrophobic Pollutants in Sediment Suspensions,” *Environmental Toxicology and Chemistry*, Vol 4, 1985, pp. 469-479.
- (106) Davenport, R., and Spacie, A., “Acute Phototoxicity of Harbor and Tributary Sediments From Lower Lake Michigan,” *Journal of Great Lakes Research*, Vol 17, 1991, pp. 51-56.
- (107) Oris, J. T., and Giesy, J. P., “The Photoenhanced Toxicity of Anthracene to Juvenile Sunfish (*Lepomis spp.*),” *Aquatic Toxicology*, Vol 6, 1985, pp. 133-146.
- (108) Ankley, G.T., Collyard, S.A., Monson, P.D. and Kosian P.A. Influence of ultraviolet light on the toxicity of sediments contaminated with polycyclic aromatic hydrocarbons. *Environ. Toxicol. Chem.* 11: 1791–1796, 1994b.
- (109) Lamberson, J. O., and Swartz, R. C., Use of Bioassays in Determining the Toxicity of Sediment to Benthic Organisms. In: M. S. Evans (ed.), *Toxic Contaminants and Ecosystem Health: A Great Lakes Focus*, John Wiley and Sons, New York, 1988, pp. 257-279.
- (110) McNulty, E.W., Dwyer, F.J., Ellersieck, M.R., Greer, I.E., Ingersoll, C.G., and Rabeni, C.F., Evaluation of the ability of reference-toxicity tests to identify stress in laboratory populations of the amphipod *Hyaella azteca*. *Environ. Toxicol. Chem.* 18:544–548, 1999.
- (111) Harkey, G. A., Lydy, M. J., Kukkonen, J., and Landrum, P. F., “Feeding Selectivity and Assimilation of PAH and PCB in *Diporeia spp.*” *Environmental Toxicology and Chemistry*, Vol 13, 1995, pp. 1445-1455. Harkey, G. A., Landrum, P. F., and Klaine, S. J., “Preliminary Studies on the Effect of Feeding During Whole-sediment Bioassays Using *Chironomus riparius* Larvae,” *Chemosphere*, Vol 28, 1994, pp. 597-606.
- (112) Wiederholm, T., Wiederholm, A. M., and Goran, M., “Freshwater Oligochaetes,” *Water, Air, and Soil Pollution*, Vol 36, 1987, pp. 131-154.
- (113) Swartz, R. C., Kemp, P. F., Schultz, D. W., and Lamberson, J. O., “Effects of Mixtures of Sediment Contaminants on the Marine Infaunal Amphipod, *Rhepoxynius abronius*,” *Environmental Toxicology and Chemistry*, Vol 7, 1988, pp. 1013-1020.
- (114) USEPA-USACE (U.S. Army Corps of Engineers). Ecological Evaluation of Proposed Discharge of Dredged Material Into Ocean Waters, Technical Committee on Criteria for Dredged and Fill Material, *Environmental Effects Laboratory*, U.S. Army Engineer Waterways Experiment Station, Vicksburg, MS, 1977.
- (115) USEPA-USACE (U.S. Army Corps of Engineers). Evaluation of Dredged Material Proposed for Ocean Disposal. EPA-503/8-91/001, Washington, DC, 1991.
- (116) USEPA-USACE (U.S. Army Corps of Engineers). Evaluation of Dredged Material Proposed for Discharge in Waters of the U.S. - testing material. EPA 823-B-98-004, Washington, DC, 1998.
- (117) Dillon T.M. Moore, D.W. and Jarvis, A.S., “The Effects of Storage Temperature and Time on Sediment Toxicity,” *Archives of Environmental Contamination Toxicology*, Vol 27, 1994, pp. 51–53
- (118) Canfield T.J., Dwyer, F.J., Fairchild, J.F., Haverland P.S., Ingersoll, C.G., Kemble N.E., Mount D.R., La Point T.W., Burton, G.A., Swift, M.C. Assessing contamination in Great Lakes sediments using benthic invertebrate communities and the sediment quality triad approach, *J. Great Lakes Res.* 22:565–583, 1996.
- (119) USEPA. Techniques for Study of Effects and Uptake of Sediment Associated Chemicals, Office of Toxic Substances. USEPA, Gulf Breeze, FL, 1986a.

- (120) Swartz, R. C., Marine Sediment Toxicity Tests, In: Contaminated Marine Sediments-Assessment and Remediation. National Research Council, National Academy Press, Washington, DC. 1989, pp. 115-129.
- (121) USEPA. 1993. Methods for Measuring the Acute Toxicity of Effluents and Receiving Waters to Freshwater and Marine Organisms. Fourth edition. Office of Research and Development. Washington, D.C. EPA-600/4-90/027F.
- (122) Sanders, H.L., P.C. Mangelsdorf, Jr., and G.R. Hampson. 1965. Salinity and faunal distribution in the Pocasset River, Massachusetts. *Limnology and Oceanography Supplement to Vol 10*:R216-R229.
- (123) Woelke, C.E. 1972. Development of a Receiving water quality bioassay criterion based on the 48-h Pacific Oyster (*Crassostrea giga*) embryo. Washington Department of Fisheries Technical Report No.9 pp 1-93.
- (124) Woelke, C.E. 1968. Development and validation of a field bioassay method with the Pacific oyster, *Crassostrea gigas*, embryo. Ph.D. Dissertation, University of Washington, Seattle, Washington. 141 p.
- (125) Cardwell, R.D., Woelke, C.E., Carr, M.I., and Sanborn, E.W. 1977. Evaluation of water quality of Puget Sound and Hood Canal in 1976. Technical Memo No. 21, National Oceanic and Atmospheric Administration, pp. 1-36.
- (126) Cardwell, R.D., S. Olsen, M.I. Carr, and E.W. Sanborn. 1979. Causes of oyster larvae mortality in south Puget Sound. NOAA Technical Memorandum ERA MESA-39. 73 p.
- (127) USEPA. 1991. Methods for measuring the acute toxicity of effluents to freshwater and marine organisms. Fourth edition. EPA 600/4-90/027F, Cincinnati, OH.
- (128) USEPA. 1994b. Short-term methods for estimating the chronic toxicity of effluents and surface waters to estuarine or marine organisms. Second edition. EPA-600/4-91/003, Cincinnati, OH.
- (129) USEPA. Quality Criteria for Water, EPA-440/5-86-001, Washington, DC, 1986b.
- (130) Walters, D. B., and Jameson, C. W., Health and Safety for Toxicity Testing, Butterworth Publications, Woburn, MA, 1984.
- (131) Bureau of National Affairs, Inc. U.S. Environmental Protection Agency General Regulation for Hazardous Waste Management, Washington, D.C., 1986.
- (132) *Methods for Measuring the Acute Toxicity of Effluents and Receiving Waters to Freshwater and Marine Organisms*, Fourth edition, EPA 600/4-90/027F, U.S. Environmental Protection Agency, Cincinnati, OH, 1993a.
- (133) USEPA. Methods for Chemical Analysis of Water and Wastes, EPA-600/4-79-020, Cincinnati, OH, 1979b.
- (134) USEPA. Standard Guide for Collection, Storage, Characterization, and Manipulation of Sediments for Toxicological Testing, USEPA Series Number Pending, Washington, DC, October 2000.
- (135) USEPA. Contaminated sediment management strategy, EPA 823-R-98-001. Office of Water, Washington, DC, 1998.
- (136) Dillon, T. M., Moore, D. W., and Jarvis, A. S., "The Effects of Storage Temperature and Time on Sediment Toxicity," *Archives of Environmental Contamination Toxicology*, Vol 27, 1994, pp. 51-53.
- (137) Becker, D.S., Rose, C.D., and Bigham, G.N. Comparison of the 10-day freshwater sediment toxicity tests using *Hyalella azteca* and *Chironomus tentans* Environ. Toxicol. Chem. 14:2089-2094, 1995.
- (138) Carr, R.S. and Chapman, D.C. Comparison of methods for conducting marine and estuarine sediment porewater toxicity tests, extraction, storage, and handling techniques. Arch. Environ. Contam. Toxicol. 28: 69-77, 1995.
- (139) Moore, D.W. and Farrar, J.D. Effect of growth on reproduction in the freshwater amphipod, *Hyalella azteca*. *Hydrobiologia* 328:127-134, 1996.
- (140) Sarda, N. and G.A. Burton, Jr. Ammonia variation in sediments: spatial, temporal and method-related effects, Environ. Toxicol. Chem. 14:1499-1506, 1995.
- (141) Sijm, R.T.H., Haller, M. and Schrap, S.M. Influence of storage on sediment characteristics and of drying sediment on sorption coefficients of organic contaminants. Bull. Environ. Contam. Toxicol. 58:961-968, 1997.
- (142) DeFoe, D.L. and Ankley, G.T. Influence of storage time on toxicity of freshwater sediments to benthic macroinvertebrates, Environ. Pollut. 99:121-131, 1998.
- (143) Ingersoll, C. G., Buckler, D. R., Crecelius, E. A., and La Point, T. W., U.S. Fish and Wildlife Service and Battelle final report for the USEPA GLNPO Assessment and Remediation of Contaminated Sediment (ARCS) Project: Biological Assessment of Contaminated Great Lakes Sediment, EPA 905-R93-006, Chicago, IL, 1993.
- (144) Schuytema, G. S., Nebeker, A. V., Griffis, W. L., and Miller, C. E., "Effects of Freezing on Toxicity of Sediments Contaminated With DDT and Endrin," *Environmental Toxicology and Chemistry*, Vol 8, 1989, pp. 883-891.
- (145) Shuba, P. J., Tatem, H. E., and Carroll, J. H., Biological Assessment Methods to Predict the Impact of Open-Water Disposal of Dredged Material, U.S. Army Corps of Engineers Technical Report D-78-5Q, Washington, DC, 1978.
- (146) Reynoldson, T. B., Day, K. E., Clarke, C., and Milani, D., "Effect of Indigenous Animals on Chronic Endpoints in Freshwater Sediment Toxicity Tests," *Environmental Toxicology and Chemistry*, Vol 13, 1994, pp. 973-977.
- (147) Ankley, G. T., Cook, P. M., Carlson, A. R., Call, D. J., Swenson, J. A., Corcoran, H. F., and Hoke, R. A., "Bioaccumulation of PCBs from Sediments by Oligochaetes and Fishes: Comparison of Laboratory and Field Studies," *Canadian Journal of Fisheries and Aquatic Sciences*, Vol 49, 1992b, pp. 2080-2085.
- (148) Swartz, R. C., Cole, F. A., Lamberson, J. O., Ferraro, S. P., Schults, D. W., DeBen, W. A., Lee II, H., and Ozretich, R. J., "Sediment Toxicity, Contamination and Amphipod Abundance at a DDT and Dieldrin-contaminated Site in San Francisco Bay," *Environmental Toxicology and Chemistry*, Vol 13, 1994, pp. 949-962, Schlekot, C. E., McGee, B. L., Boward, D. M., Reinharz, E., Velinsky, D. J., and Wade, T. L. 1994. Tidal River Sediments in the Washington, D.C. Area. III. Biological Effects Associated With Sediment Contamination. *Estuaries* Vol 17, 1994, pp 334-344; McGee, B. L., Schlekot, C. E., Boward, D. M., and Wade, T. L. 1994. Sediment Contamination and Biological Effects in A Chesapeake Bay Marina. *Ecotoxicology* 3: McGee, B. L., Schlekot, C. E., and Reinharz, E. 1993. Assessing Sublethal Levels of Sediment Contamination Using the Estuarine Amphipod *Leptocheirus plumulosus*. *Environmental Toxicology and Chemistry*, Vol 12, pp. 577-587.
- (149) Ditsworth, G. R., Schults, D. W., and Jones, J. K. P., "Preparation of Benthic Substrates for Sediment Toxicity Testing," *Environmental Toxicology and Chemistry*, Vol 9, 1990, pp. 1523-1529.
- (150) Lamberson, J. O., and Swartz, R. C., Spiked-Sediment Toxicity Test Approach. In: Sediment Classification Methods Compendium, pp. 4-1 to 4-10. EPA 823-R-92-006, Washington, DC, 1992.
- (151) Besser, J.M., Ingersoll, C.G., Leonard, E., and Mount, D.R. Effect of zeolite on toxicity of ammonia freshwater sediments: Implications for sediment toxicity identification evaluation procedures. *Environ. Toxicol. Chem.* 17:2310-2317, 1998.
- (152) Plumb, Jr., R. H., Procedures for handling and Chemical Analysis of Sediment and Water Samples, Technical Committee on Criteria for Dredged

- and Fill Material, USEPA/USACE. EPA-4805572010. USEPA Great Lakes Laboratory. Grosse Ile, MI, 1981.
- (153) Van Rees, K.C.J., E.A. Sudlicky, P. Suresh, C. Rao and K.R. Reddy. Evaluation of laboratory techniques for measuring diffusion coefficients in sediments. *Environ. Sci. Technol.* 25:1605–1611, 1991.
- (154) Bufflap, S.T. and H.E. Allen. Comparison of pore water sampling techniques for trace metals. *Wat. Res.* 29:2051–2054, 1995.
- (155) Ankley, G.T. and Schubauer-Berigan, M.K. Background and overview of current sediment toxicity identification procedures. *J. Aquat. Ecosystem Health.* 4:133–149, 1995.
- (156) Winger, P.V., P.J., Lasier and B.P. Jackson. The influence of extraction procedure on ion concentrations in sediment pore water. *Arch. Environ. Contam. Toxicol.* 35:8–13, 1998.
- (157) Burton, G.A., Jr. Sediment collection and processing factors affecting realism. In: G.A. Burton, Jr. (ed.) *Sediment Toxicity Assessment*. Lewis Publ. Boca Raton, FL, pp. 37–66, 1992.
- (158) Bufflap, S.T. and H.E. Allen. Sediment pore water collection methods for trace metal analysis: a review. *Wat. Res.* 29:165–177, 1995.
- (159) Horowitz, A.J., K.A. Elrick and M.R. Colberg. The effect of membrane filtration artifacts on dissolved trace element concentrations. *Wat. Res.* 26:753–763, 1992.
- (160) USEPA EPA requirements for quality assurance project plans for environmental data operations. Draft interim final. EPA QA/R-5. Office of Research and Development, Washington, D.C. 1994e.
- (161) USEPA. Short-Term Methods for Estimating the Chronic Toxicity of Effluents and Receiving Waters to Marine and Estuarine Organisms, Second edition, EPA 600/4-91/003, Cincinnati, OH, 1993c.
- (162) USEPA. QA/QC guidance for sampling and analysis of sediments, water, and tissues for dredged material evaluations-chemical evaluations, EPA 823-B-95-001, Office of Water, Washington, DC, 1995.
- (163) Food and Drug Administration, Good Laboratory Practices for Non-Clinical Laboratory Studies, Part 58. *Federal Register* 43(247): 60013-60020 (December 22, 1978), 1978.
- (164) USEPA. Handbook for Analytical Quality Assurance in Water and Wastewater Laboratories, EPA-600/4-79-019, Cincinnati, OH, 1979a.
- (165) USEPA. Proposed Good Laboratory Practice Guidelines for Toxicity Testing, Paragraph 163.60-6. *Federal Register* 45:26377-26382 (April 18, 1980), 1980a.
- (166) USEPA. Physical, Chemical, Persistence, and Ecological Effects Testing; Good Laboratory Practice Standards (Proposed Rule). 40 CFR 772, *Federal Register* 45:77353-77365 (November 21, 1980), 1980b.
- (167) USEPA. Short-Term Methods for Estimating the Chronic Toxicity of Effluents and Receiving Waters to Freshwater Organisms, Third edition. EPA 600/4-91/002, Cincinnati, OH, 1993b.
- (168) DeWoskin, R. S., Good Laboratory Practice Regulations: A Comparison, Research Triangle Institute, Research Triangle Park, NC, 1984, p. 63.
- (169) Shoemaker, C.R. 1932. A new amphipod of the genus *Leptocheirus* from Chesapeake Bay. *Wash. Acad. Sci.* 22: 548-551. Sijm, R.T.H., M. Haller, and S.M. Schrap. 1997. Influence of storage on sediment characteristics and of drying sediment on sorption coefficients of organic contaminants. *Bull. Environ. Contam. Toxicol.* 58:961-968.
- (170) McGee, B.L., D.A. Wright and D.J. Fisher. 1998. Biotic factors modifying the acute toxicity of aqueous cadmium to the estuarine amphipod *Leptocheirus plumulosus*. *Archives of Environmental Contamination and Toxicology* 35:34-40.
- (171) Bousfield, E.L. 1973. *Shallow Water Gammaridean Amphipoda of New England*. Ithaca, NY: Cornell University Press, 312 pp.
- (172) McGee, B.L. 1998. Population dynamics of the estuarine amphipod *Leptocheirus plumulosus*: Implications for sediment toxicity tests. Ph.D. dissertation, University of Maryland, College Park, MD.
- (173) Duan, Y., S.I. Guttman, and J.T. Oris. 1997. Genetic differentiation among laboratory populations of *Hyalella azteca*: Implications for toxicology. *Environ. Toxicol. Chem.* 16:691-695.
- (174) Tomasovic, M., Dwyer, F. J., Ingersoll, C. G., and Greer, I. E., “Recovery of *Hyalella azteca* From Sediment,” *Environmental Toxicology and Chemistry*, Vol 14, No. 7, 1995, pp. 1177-1180.
- (175) Lee, D.R. 1980. Reference toxicants in quality control of aquatic bioassays. In, *Aquatic invertebrate bioassays*. ASTM STP 715, A.L. Buikema and J. Cairns, Jr., eds, Philadelphia, PA. pp. 188-199.
- (176) Nichols F.H. and J.K. Thompson. 1985. Persistence of and introduced mudflat community in south San Francisco Bay, California. *Mar. Ecol. Prog. Ser.* 24:83-97.
- (177) Hopkins, D.R. 1986. Atlas of the distributions and abundances of common benthic species in San Francisco Bay, California. *Water Resources Investigations Report* 86- 4003. Menio Park, CA: U.S. Geological Survey. 16 p.
- (178) Richards, S.W. 1963. The demersal fish populations of Long Island Sound: III Food for juveniles from a mud locality. *Bull. Bingham Oceanogr. Coil.*, 18:32-101.
- (179) Hyland, J.L. 1981. Comparative structure and response to (petroleum) disturbance in two nearshore infaunal communities. Ph.D. Thesis, University of Rhode Island, 141p.
- (180) Weisberg, S.B., J.B. Frithsen, A.F. Holland, J.F. Paul, K.J. Scott, J.K. Summers, H.T. Wilson, R. Valente, D.G. Heimbuch, J. Gerritsen, S.C. Schimmel, and R.W. Latimer. 1992. EMAP-Estuaries Virginian Province 1990 Demonstration Project Report EPA 600/R-92/100. USEPA, Environmental Research Laboratory, Narragansett, RI.
- (181) Mills, E.L. 1967. The biology of an amphipod crustacean sibling species pair. *J. Fisheries Res. Board Canada*, 24:305-355.
- (182) Nelson, W.G. 1980. Reproductive patterns of gammaridean amphipods. *Sarsia*, 65:61-71.
- (183) Redmond, M.S., K.J. Scott, R.C. Swartz and J.K.P. Jones. 1994. Preliminary culture and life cycle experiments with the benthic amphipod *Ampelisca abdita*. *Environ. Toxicol. Chem.* 13(8): 1355-1365.
- (184) Environment Canada. 1998. Biological Test Method: Reference Method for Determining Acute Lethality of Sediment to Marine or Estuarine Amphipods. Environment Canada, Conservation and Protection, Ottawa, Ontario, EPS 1/RM/35.
- (185) Holland, A.F., A.T. Shaughnessy, L.C. Scott, V.A. Dickens, J.A. Ranasinghe, and J.K. Summers. 1988. Progress report: Long-term benthic monitoring and assessment program for the Maryland portion of Chesapeake Bay (July 1986-October 1987). PPRP-LTB/EST-88-1. VERSAR, Columbia, MD.
- (186) Schlekot, C.E., B.L. McGee, D.M. Boward, D.J. Velinsky, and T.L. Wade. 1994. Tidal river sediments in the Washington, D.C., area. III. Biological effects associated with sediment contamination. *Estuaries* 17:334-344.
- (187) Lamberson, J.O. and R.C. Swartz. 1988. Use of bioassays in determining the toxicity of sediment to benthic organisms. In *Toxic Contaminants and Ecosystem Health: A Great Lakes Focus.*, M.S. Evans, ed. John Wiley and Sons, New York, NY. pp. 257-279.

- (188) Jordan, R.A. and C.E. Sutton. 1984. Oligohaline benthic invertebrate communities at two Chesapeake Bay power plants. *Estuaries* 7:192-212.
- (189) Marsh, A.G. and K.R. Tenore. 1990. The role of nutrition in regulating the population dynamics of opportunistic, surface deposit feeders in a mesohaline community. *Limnol. Oceanogr.* 35(3):710-724.
- (190) Kemp, P.F., F.A. Code, and R.C. Swartz. 1985. Life history and production of the Phoxocephalid amphipod *Rhepoxynius abronius* (Barnard). *J. Crustacean Biology* 5:449- 464.
- (191) Barnard, J.L., and C.M. Barnard. The genus *Rhepoxynius* (Crustacea: Amphipoda: Phoxocephalidae) in American Seas. *Smithsonian Contributions to Zoology* 357:1-49.
- (192) Puget Sound Estuary Program. 1995. Recommended guidelines for conducting laboratory bioassays on Puget Sound sediments. Prepared for U.S. Env. Prot. Agency, Region 10, Office of Puget Sound, Seattle, WA and Puget Sound Water Quality Authority, P.O. Box 40900, Olympia, WA.
- (193) DeWitt, T.H., G.R. Ditsworth, and R.C. Swartz. 1988. Effects of natural sediment features on survival of the Phoxocephalid amphipod, *Rhepoxynius abronius*. *Marine Envir. Res.*, 25:99-124.
- (194) McLeay, D.J., S. Yee, K. Doe, and S. Wade. 1991. Phase-II and Phase-III studies by Environment Canada laboratories of 10-day tests for sediment toxicity using marine or estuarine infaunal amphipods. Report prepared for Environment Canada (EP, Conservation and Protection) and IGATG by McLeay Associates Ltd., West Vancouver, BC.
- (195) Ramsdell, K.A., J.A. Strand, and V.I. Cullinan. 1989. Amphipod bioassay of selected sediments from Sequim Bay, Washington. In: *Oceans 1989, Volume 2: Ocean Pollution*. IEEE Publication No. 89CH2780-5, pp. 443-448.
- (196) Stein, J.R. 1973. *Handbook of phycological methods: Culture methods and growth measurements*. New York, NY: Cambridge University Press. 445 p.
- (197) Ankley, G. T., Phipps, G. L., Leonard, E. N., Benoit, D. A., Mattson, V. R., Kosian, P. A., Cotter, A. M., Dierkes, J. R., Hansen, D. J., and Mahony, J. D., "Acid-Volatile Sulfide as a Factor Mediating Cadmium and Nickel Bioavailability in Contaminated Sediment," *Environmental Toxicology and Chemistry*, Vol 10, 1991a, pp. 1299-1307.
- (198) Burton, G. A., Jr. and Ingersoll, C. G., "Evaluating the Toxicity of Sediments," *The Assessment of Contaminated Great Lakes Sediment*, EPA 905-B94-002, Chicago, IL, 1994.
- (199) Long, E. R., Buchman, M. F., Bay, S. M., Breteler, R. J., Carr, R. S., Chapman, P. M., Hose, J. E., Lissner, A. L., Scott, J., and Wolfe, D. A., "Comparative Evaluation of Five Toxicity Tests With Sediments from San Francisco Bay and Tomales Bay, California," *Environmental Toxicology and Chemistry*, Vol 9, 1990, pp. 1193-1214.
- (200) Canfield, T. J., Kemble, N. E., Brumbaugh, W. G., Dwyer, F. J., Ingersoll, C. G., and Fairchild, J. F., "Use of Benthic Invertebrate Community Structure and the Sediment Quality Triad to Evaluate Metal-contaminated Sediment in the Upper Clark Fork River, Montana," *Environmental Toxicology and Chemistry*, Vol 13, 1994, pp. 1999-2012.
- (201) Canfield T.J., Brunson, E.L., Dwyer, F.J., Ingersoll, C.G., Kemble N.E. Assessing sediments from the upper Mississippi river navigational pools using a benthic invertebrate community evaluation and the sediment quality triad approach. *Arch. Environ. Contam. Toxicol.* 35:202–212, 1998.
- (202) Marcus, A. H., and Holtzman, A. P., A Robust Statistical Method for Estimating Effects Concentrations in Short-Term Fathead Minnow Toxicity Tests. Manuscript submitted to the Criteria and Standards Division, U.S. Environmental Protection Agency, by Battelle Washington Environmental Program Office, Washington, DC, June 1988, under EPA Contract No. 69-03-3534. 39 p., 1988; Hall, W. S., Patoczka, J. B., Miranda, R. J., Porter, B. A., and Miller, E. Acute toxicity of industrial surfactants to *Mysidopsis bahia*. *Archives of Environmental Contamination and Toxicology*, Vol 18, pp. 765-772.
- (203) Grothe, D. R., and Kimerle, R. A., "Interand Intra-Laboratory Variability in *Daphnia magna* Effluent Toxicity Test Results," *Environmental Toxicology and Chemistry*, Vol 4, 1985, pp. 189-192.
- (204) Pittinger, C. A., Woltering, D. M., and Masters, J. A., "Bioavailability of Sediment-Sorbed and Soluble Surfactants to *Chironomus riparius* (Midge)," *Environmental Toxicology and Chemistry*, Vol 8, 1989, p. 11.
- (205) Mearns, A.J., R.C. Swartz, J.M. Cummins, P.A. Dinnel, P. Plesha, and P.M. Chapman. 1986. Inter-laboratory comparison of a sediment toxicity test using the marine amphipod, *Rhepoxynius abronius*. *Mar. Environ. Res.* 19:13-37.
- (206) USEPA. 2000. Methods for measuring the toxicity and bioaccumulation of sediment-associated contaminants with freshwater invertebrates, second edition. EPA600/R-99/064, Duluth, MN.
- (207) Tomasovic, M. et al. 1994. Recovery of *Hyalella azteca* from sediment. *Environ. Toxicol. Chem.* 13:1311-1314.
- (208) Dillon, T.M. and A.B. Gibson. 1986. Bioassessment methodologies for the regulatory testing of freshwater dredged material. Miscellaneous Paper EL-86-6, U.S. Army Engineer Waterways Experiment Station, Vicksburg, MS.
- (209) Ingersoll, C.G. 1991. Sediment toxicity and bioaccumulation testing. *Standardization News* 19: 28-33.
- (210) Burton, G.A., M.K. Nelson, and C.G. Ingersoll. 1992. Freshwater benthic toxicity tests. In *Sediment Toxicity Assessment*, G.A. Burton, ed.. Lewis Publishers, Chelsea, MI. pp. 213-240.
- (211) Niewolny, L.A., T.H. DeWitt, and B.D. Gruendell. 1997. ALCOA Point Comfort Sediment Quality Triad Project: *Leptocheirus plumulosus* Chronic Survival Sediment Toxicity Test. Prepared for Parametrix, Inc. Kirkland, Washington. PNWD-2381. Battelle Pacific Northwest Division, Richland, WA.
- (212) DeWitt, T.H., M.R. Pinza, and L.A. Niewolny. 1997c. Phase I (Dilution Series Study) and Phase II (Field-Collected Sediment Study) of the PTI Chronic *Leptocheirus plumulosus* Project. Prepared for PTI Environmental, Bellevue, Washington. PNWD-2384. Battelle Pacific Northwest Division, Richland, WA..
- (213) Sims, J.G. and D.W. Moore. 1995. Risk of pore water hydrogen sulfide toxicity in dredged material bioassays. Miscellaneous paper, D-95-4, U.S. Army Engineer Waterways Experiment Station, Vicksburg, MS.
- (214) Moore et al. 1995. Long-term Storage of Sediments: Implications for Sediment Toxicity Testing. *Env. Pollution*, 89, No. 2 147-154.
- (215) Knezovich, J.P., D.J. Steichen, J.A. Jelinski, and S.L. Anderson. 1996. Sulfide tolerance of four marine species used to evaluate sediment and pore-water toxicity. *Bull. Environ. Contam. Toxicol.* 57: 450-457.
- (216) Bagarinao, T. 1992. Sulfide as an environmental factor and toxicant: tolerance and adaptations in aquatic organisms. *Aquatic Tox.* 24: 21-62.
- (217) Scott, J., C. Mueller, W. Starkel, and R. Mayer. 1996. Chronic response of *Leptocheirus plumulosus* in 28-day exposures to contaminated estuarine sediments. Poster presented at the Society of Environmental Toxicology and Chemistry 17th Annual Meeting, November 17-21, 1996, Washington, D.C.
- (218) McGee, B.L. and M. Spencer. 2001. A field-based population model for the sediment toxicity test organism *Leptocheirus plumulosus*. II. Model Application. *Marine Environmental Research* 51:347-363.

- (219) Spencer, M. and B.L. McGee. 2001. A field-based population model for the sediment toxicity test organism *Leptocheirus plumulosus*. I. Model Development. *Marine Environmental Research* 51:327-345.
- (220) Oakden J.M. 1984. Feeding and sediment selection in five species of Central Californian phoxocephalid amphipods. *J. Crustacean Biol.* 4:233-247.
- (221) Environment Canada, Biological Test Methods: Reference Method for Determining Acute Utility of Sediment to Marine or Estuarine Amphipods. *Environment Canada EPS 1/RM/35*, Ottawa, Ontario, 57pp. 1998.

SUMMARY OF CHANGES

The primary technical changes from the previous version of this standard (E 1367-99) are summarized in this section.

- (1) The information on conducting 10-d sediment toxicity tests with estuarine and marine amphipods has been updated based on information presented in USEPA (1994a (1)).
- (2) The information on conducting 28-d sediment toxicity tests with *Leptocheirus plumulosus* has been included based on information presented in USEPA-USACE (2001(2)).
- (3) The general format of the standard has been revised to be consistent with Test Method E 1706.

ASTM International takes no position respecting the validity of any patent rights asserted in connection with any item mentioned in this standard. Users of this standard are expressly advised that determination of the validity of any such patent rights, and the risk of infringement of such rights, are entirely their own responsibility.

This standard is subject to revision at any time by the responsible technical committee and must be reviewed every five years and if not revised, either reapproved or withdrawn. Your comments are invited either for revision of this standard or for additional standards and should be addressed to ASTM International Headquarters. Your comments will receive careful consideration at a meeting of the responsible technical committee, which you may attend. If you feel that your comments have not received a fair hearing you should make your views known to the ASTM Committee on Standards, at the address shown below.

This standard is copyrighted by ASTM International, 100 Barr Harbor Drive, PO Box C700, West Conshohocken, PA 19428-2959, United States. Individual reprints (single or multiple copies) of this standard may be obtained by contacting ASTM at the above address or at 610-832-9585 (phone), 610-832-9555 (fax), or service@astm.org (e-mail); or through the ASTM website (www.astm.org).