



Standard Terminology Relating to Biological Effects and Environmental Fate¹

This standard is issued under the fixed designation E 943; the number immediately following the designation indicates the year of original adoption or, in the case of revision, the year of last revision. A number in parentheses indicates the year of last reapproval. A superscript epsilon (ϵ) indicates an editorial change since the last revision or reapproval.

1. Scope

1.1 This terminology document defines terms commonly used in standards developed by ASTM Committee E-47 on Biological Effects and Environmental Fate. This terminology document is intended to be consistent with the use of terms in ASTM standards related to this field and, to the extent possible, with use by other organizations.

1.1.1 If a specific Committee E-47 standard uses one of these terms in a different context, then the term must be defined in that standard. A term used only in a specific ASTM standard need not be included in this terminology document.

2. Terminology

2.1 Definitions:

acute test—a comparative study in which organisms, that are subjected to different treatments, are observed for a short period usually not constituting a substantial portion of their life span.

DISCUSSION—There is no specific test duration that represents a distinct boundary between acute and chronic test durations for any species. Although acute or chronic test procedures may specify standard duration(s), these durations have not been intended to define an acute:chronic boundary.

Acute tests often utilize mortality as the only measure of effect; chronic tests usually include additional measures of effect such as growth or reproduction.

attraction—a response towards or to facilitate contact with a material or condition.

avoidance—a response away from or to limit contact with a material or condition.

BAF (bioaccumulation factor), n —the quotient obtained by dividing the concentration of a substance in an organism (or specified tissue) by its concentration in a specified exposure medium, for example, air, food, sediment, soil, water, when several media are possible sources (see **bioaccumulation**).

behavior, n —observable, recordable, or measurable actions or activity of an organism.

DISCUSSION—This definition conveys the idea of motion whether motility is involved or not, and excludes physiological responses,

¹ This terminology is under the jurisdiction of ASTM Committee E-47 on Biological Effects and Environmental Fate and is the direct responsibility of Subcommittee E47.02 on Terminology.

Current edition approved April 10, 2000. Published July 2000. Originally published as E 943 – 83. Last previous edition E 943 – 00.

death, and so forth, from the concept. It avoids the issue of internal versus external stimuli.

bioaccumulation—the net accumulation of a substance by an organism as a result of uptake from all environmental sources.

bioassay—an experiment that uses living whole organisms, tissues or cells to measure the presence, the concentration or the relative potency of one or more chemicals.

DISCUSSION—A bioassay must include the appropriate controls(s). There is no intended stipulation of endpoint for such a test; the response may be positive or negative. This term defines a subset of the protocols (methods) referred to by the term “biological assay” (Finney, 1947)

bioconcentration—the net accumulation of a substance by an aquatic organism as a result of uptake directly from aqueous solution.

biomagnification—the increase in tissue concentration of poorly depurated materials in organisms along a series of predator-prey associations, primarily through the mechanism of dietary accumulation.

biomarker, n —a biological measure (within organisms) of exposure to, effects of or susceptibility to, environmental stress using molecular, genetic, biochemical, histological, or physiological techniques.

biomarker assay—an experiment that uses a molecular, genetic, biochemical, histological, anatomical or physiological technique to assess exposure, response or susceptibility of organisms, tissue or cells to environmental stress.

chronic test—a comparative study in which organisms, that are subjected to different treatments, are observed for a long period or a substantial portion of their life span.

DISCUSSION—There is no specific test duration that represents a distinct boundary between acute and chronic test durations for any species. Although acute or chronic test procedures may specify standard duration(s), these durations have not been intended to define an acute:chronic boundary.

Acute tests often utilize mortality as the only measure of effect; chronic tests usually include additional measures of effect such as growth or reproduction.

control sediment—a sediment that is essentially free of contaminants and is used routinely to assess the acceptability of a test.

depuration—loss of a substance from an organism as a result of any active or passive process.

dietary accumulation—the net accumulation of a substance by an organism as a result of ingestion in the diet.

EC50—a statistically or graphically estimated concentration that is expected to cause one or more specified effects in 50 % of a group of organisms under specified conditions.

ED50—a statistically or graphically estimated dose that is expected to cause one or more specified effects in 50 % of a group of organisms under specified conditions.

exposure—contact with a chemical or physical agent.

fate, environmental—the form and location of a material resulting from transport and transformation.

hazard—the adverse effect(s) that may result from exposure(s).

IC50—a statistically or graphically estimated concentration of test material that, under specified conditions, is expected to cause a 50 % inhibition of a biological process (such as growth or reproduction) for which the data are not dichotomous.

indigenous species—a species that is likely, due to historical presence, to occur at a specified site for some portion of its life span.

DISCUSSION—This definition is intended to remove the requirement that the species occur presently at a site. This definition excludes species that have been introduced either intentionally or unintentionally by man whether recently or in the remote past. The terms “indigenous” and “native” are synonymous in this context.

interstitial water—water occupying space between sediment or soil particles (Syn. **pore water**).

key species—a species of special concern for ecological reasons.

LC50—a statistically or graphically estimated concentration that is expected to be lethal to 50 % of a group of organisms under specified conditions.

LD50—a statistically or graphically estimated dose that is expected to be lethal to 50 % of a group of organisms under specified conditions.

life-cycle test—a comparative study in which organisms, that are subjected to different treatments, are observed at least from a life stage in one generation to the same life stage in the next generation.

lowest-observed-effect concentration (LOEC)—in a toxicity test, the tested concentration of one or more chemicals immediately above the highest tested concentration that did not result in a statistically significant change in the particular toxicological variable compared to that value in the control (s).

DISCUSSION—Within a concentration-effect curve at concentrations near the NOEC and LOEC, the following situation can occur: one concentration might induce an effect that is significantly greater than the control or background, while the next higher concentration induces an effect that is not significantly greater than control or background, and all the higher concentrations induce effects that are significantly greater than control or background. In this region of uncertainty, the concentration inducing a significant effect may be inverted one or more times. In order to avoid a situation where the LOEC is less than the NOEC, the NOEC is defined as the concentration immediately below the region of uncertainty, and the LOEC is defined as the concentration immediately above this region. If the region of uncertainty is large, the investigator may not choose to define a NOEC or LOEC.

no-observed-effect concentration (NOEC)—in a toxicity test, the tested concentration of one or more chemicals

immediately below the lowest tested concentration that resulted in a statistically significant change in a particular toxicological variable compared to that value in the control (s).

DISCUSSION—Within a concentration-effect curve at concentrations near the NOEC and LOEC, the following situation can occur: one concentration might induce an effect that is significantly greater than the control or background, while the next higher concentration induces an effect that is not significantly greater than control or background, and all the higher concentrations induce effects that are significantly greater than control or background. In this region of uncertainty, the concentration inducing a significant effect may be inverted one or more times. In order to avoid a situation where the LOEC is less than the NOEC, the NOEC is defined as the concentration immediately below the region of uncertainty, and the LOEC is defined as the concentration immediately above this region. If the region of uncertainty is large, the investigator may not choose to define a NOEC or LOEC.

pore water—water occupying space between sediment or soil particles (Syn. **interstitial water**).

reference sediment—a whole sediment near an area of concern used to assess sediment conditions exclusive of material(s) of interest.

replicate, *n*—each of several experimental units that are tested simultaneously using the same experimental conditions.

DISCUSSION—Replicates are independent executions of treatments upon experimental units, or multiple samples collected from a location. The observations from these multiple units provide the data from which statistics are computed. The purpose of replicates is to determine the variability of the effect of a given treatment within an experiment or to determine the variability among independent samples from a given location. In both field and laboratory experiments, study design and equipment must prevent or minimize exchange of test media between replicates. Repeat of whole experiments at different times are not considered replicates because time is a variable in the case of repeats.

resident species—a species that is regularly present at a specified site for some portion of its life span.

DISCUSSION—This definition is intended to be inclusive of species with small home ranges, species that have foraging ranges distinct from sleeping/reproducing areas, and migratory species. The definition is devoid of any reference to the type of activity within a specified site (feeding, sleeping, reproducing, and so forth), but relates only to normal presence. The concept is inclusive of stocked populations. When stocked populations are considered, it is appropriate to state this explicitly.

risk—the probability or likelihood an adverse effect will occur.

sediment—(1) particulate material that usually lies below water, and (2) formulated particulate material that is intended to lie below water in a test.

spiked sediment—a sediment to which a material has been added for experimental purposes.

surrogate species—a species that is tested to estimate responses of other species, for which direct testing is impractical.

toxicity—the property of a chemical, or combination of chemicals, to adversely affect organisms, tissues or cell.

toxicity test—an experiment used to study the adverse effect (s) of one or more chemicals on whole organisms, tissues, or cells.

DISCUSSION—A toxicity test must include the appropriate control (s).

The only intended stipulation of endpoint for such a test is that the effect must be adverse. This term, defines a subset of the protocols (methods) referred to by the term biological assay. (Finney, 1947).

whole sediment—sediment and associated pore water that have had minimal manipulation.

uptake—acquisition of a substance from the environment by an organism as a result of any active or passive process.

This standard is subject to revision at any time by the responsible technical committee and must be reviewed every five years and if not revised, either reapproved or withdrawn. Your comments are invited either for revision of this standard or for additional standards and should be addressed to ASTM Headquarters. Your comments will receive careful consideration at a meeting of the responsible technical committee, which you may attend. If you feel that your comments have not received a fair hearing you should make your views known to the ASTM Committee on Standards, at the address shown below.

This standard is copyrighted by ASTM, 100 Barr Harbor Drive, PO Box C700, West Conshohocken, PA 19428-2959, United States. Individual reprints (single or multiple copies) of this standard may be obtained by contacting ASTM at the above address or at 610-832-9585 (phone), 610-832-9555 (fax), or service@astm.org (e-mail); or through the ASTM website (www.astm.org).